


Sexually Transmitted Disease Profile

King County 2012



Disease Control and Health Statistics
Infectious Disease Assessment Unit



Sexually Transmitted Disease Profile

King County 2012



For persons with disabilities, this document is available in other formats. To submit a request, please call 1-800-525-0127 (TDD/TYY 1-800-833-6388).

Washington State Department of Health
Disease Control and Health Statistics
Infectious Disease Assessment Unit
Olympia, Washington
(360) 236-3455

Contents

Introduction.....1

Data Sources, Definitions, and Limitations1

County STD Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, 20124

Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates per 100,000 population, 1993-20124

Figure 2. Age-specific Chlamydia Incidence Rates by Gender, 20125

Figure 3. Chlamydia Incidence Rates by Gender, 1993-20125

Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates per 100,000 population, 1993-20126

Figure 5. Age-specific Gonorrhea Incidence Rates by Gender, 20126

Figure 6. Gonorrhea Incidence Rates by Gender, 1993-20127

Primary & Secondary Syphilis

Figure 7. Primary & Secondary Syphilis Cases and Incidence Rates, 1993-20127

Figure 8. Age-specific Primary & Secondary Syphilis Incidence Rates by Gender, 20128

Figure 9. Primary & Secondary Syphilis Incidence Rates by Gender, 1993-20128

Genital Herpes

Figure 10. Genital Herpes Cases and Incidence Rates per 100,000 population 1993-2012.....9

Figure 11. Age-specific Genital Herpes Incidence Rates by Gender, 20129

Figure 12. Genital Herpes Incidence Rates by Gender, 1993-201210

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group11

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group13

Table 4. Primary & Secondary Syphilis Cases and Incidence Rate by Gender/Age Group15

Table 5. Genital Herpes Cases and Incidence Rates by Gender and Age Group17

Introduction

Sexually transmitted infections continue to be the most frequently diagnosed and reported notifiable conditions in Washington State. This report describes the sexually transmitted disease burden in King County. Data are presented for the more commonly reported diseases of primary and secondary syphilis, chlamydial infection, gonorrhea and genital herpes when at least twenty (20) cases of disease were diagnosed in 2012 and the corresponding rates are statistically stable. The incidence rates, including those by age and sex, are presented graphically. The report concludes with tables containing a decade of historical data by age group and gender.

Data Sources, Definitions and Limitations

Cases: Surveillance cases are the number of new episodes of disease (not unique persons) diagnosed in a given year. Cases are identified and submitted by health care providers to local health jurisdictions and entered into the Washington State Department of Health Public Health Information Management System – Sexually Transmitted Diseases (PHIMS-STD) data system. To be included in surveillance reporting, each case must meet disease definitions (see below). Data presented in this report represent new cases of infection diagnosed during a given year and reported as of March 4, 2013.

Confidence Intervals: A confidence interval is a range around a measurement that suggests how precise the measurement is. The confidence interval displays both the possible range around the estimate and also how stable the estimate is. Wider confidence intervals in relation to the estimate itself indicate instability. A confidence interval is always qualified by a particular confidence level, usually expressed as a percentage. In this report, 95% confidence intervals are used.

Disease Definitions:

- | | |
|----------------------|--|
| Chancroid | – A sexually transmitted infection caused by the bacterium <i>Haemophilus ducreyi</i> that may include the symptoms of painful genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of <i>H. ducreyi</i> from a clinical specimen. |
| Chlamydia (CT) | – A sexually transmitted infection caused by the bacterium <i>Chlamydia trachomatis</i> that may include the symptoms of swelling and pain of internal sexual organs, however the infection often has no symptoms in women. Cases are defined by laboratory detection of <i>C. trachomatis</i> from a clinical specimen. |
| Genital Herpes (HSV) | – A sexually transmitted infection caused by the herpes simplex viruses type 1 and type 2 that may include the symptoms of blisters or sores in the genital area. Cases are defined by laboratory detection of herpes simplex virus (HSV1 or HSV2) or positive antibody response from a clinical specimen. Reportable cases include only adult genital initial infection and neonatal infection. |
| Gonorrhea (GC) | – A sexually transmitted infection caused by the bacterium <i>Neisseria gonorrhoeae</i> that may include the symptoms of swelling and pain of |

internal sexual organs, however the infection sometimes has no symptoms. Cases are defined by laboratory detection of the bacterium *N. gonorrhoeae* from a clinical specimen.

Granuloma Inguinale (GI) – A sexually transmitted infection caused by the bacterium *Klebsiella granulomatis* that may include the symptoms of slowly increasing genital sores and swollen pelvic lymph nodes. Cases are defined by microscopic examination of a clinical specimen.

Lymphogranuloma Venereum (LGV) – A sexually transmitted infection caused by three strains of *Chlamydia trachomatis* that may include the symptoms of genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of the L1, L2 and L3 serovars of *C. trachomatis* from a clinical specimen.

Syphilis – A sexually transmitted infection caused by the bacterium *Treponema pallidum* that may include many kinds of symptoms or none at all depending upon the stage of disease. Cases are defined and assigned a stage by a combination of positive blood tests, symptoms, and history of previous treatment. The U.S. Centers for Disease Control and Prevention (CDC) provides guidelines with additional details of surveillance definitions and staging criteria. The stages of primary and secondary (P&S) syphilis are grouped together for analysis in this report.

Primary – identified by the presence of one or many painless sores.

Secondary – identified by the presence of a rash on one or more areas of the body, often with fever, fatigue or other symptoms at the same time.

Other Stages – additional stages of syphilis include, early latent, latent, late latent, and congenital. See CDC guidelines for specific criteria: www.cdc.gov/std/

Incidence Rates: Incidence rates in this report are calculated as the number of new episodes of a disease (not unique persons) diagnosed in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates were not presented for when the calculated relative standard error is greater than thirty percent (>30%) as this indicates statistical instability.

Limitations: The data presented in this report may be subject to a number of limiting factors. Clinically diagnosed cases (without laboratory confirmation) may be missed through public health surveillance systems. Depending upon diagnosing practices, completeness of reporting may vary by the source of health care. In addition, the diagnosing practitioner is responsible for providing the case information including the patient demographic data items of age and gender upon which many of the analyses in this report depend. Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. Also, small increases or decreases in numbers from year to year can look large if the actual number of cases is small. Care should be taken in interpreting these data in light of known limitations.

Population: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM) April 1, 2012.

Tabular Data: The data tables are provided in hopes that community and local partners will use these historical data as a resource for future health planning. Users of these data should keep in mind that small numbers often produce rates that can be unstable from year to year and may not be reliable when comparing across strata within a particular year. Data tables for additional years previous are available upon request.

Anyone with specific questions about how these data should be interpreted is encouraged to contact the Infectious Disease Assessment Unit at 360-236-3455.

King County STD Disease Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, King County, 2012


Disease	County Cases	County Rate§	WA State Rate
Chlamydia	6763	345.6	360.8
Gonorrhea	1527	78.0	48.1
P&S Syphilis	210	10.7	4.4
Genital Herpes	742	37.9	32.2
Chancroid/GI/LGV	0		
Total	9242		

§ Crude incidence rate per 100,000 population.

* Rates are suppressed for counts with RSE >30% due to statistical instability.


Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates* per 100,000 population, King County, 1993-2012




* Crude incidence rate with 95% confidence intervals.

Figure 2. Age-specific Chlamydia Incidence Rates* by Gender, King County, 2012



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 3. Chlamydia Incidence Rates by Gender*, King County, 1993-2012



* Gender-specific crude incidence with 95% confidence intervals.


Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates* per 100,000 population, King County, 1993-2012




* Crude incidence rate with 95% confidence intervals.

Figure 5. Age-specific Gonorrhea Incidence Rates* by Gender, King County, 2012



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 6. Gonorrhea Incidence Rates by Gender*, King County, 1993-2012



* Gender-specific crude incidence with 95% confidence intervals.


Primary and Secondary Syphilis

Figure 7. Primary and Secondary Syphilis Cases and Incidence Rates* per 100,000 population, King County, 1993-2012




* Crude incidence rate with 95% confidence intervals.

Figure 8. Age-specific Primary and Secondary Syphilis Incidence Rates* by Gender, King County, 2012



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 9. Primary and Secondary Syphilis Incidence Rates by Gender*, King County, 1993-2012



* Gender-specific crude incidence with 95% confidence intervals.


Genital Herpes

Figure 10. Genital Herpes Cases and Incidence Rates* per 100,000 population, King County, 1993-2012




* Crude incidence rate with 95% confidence intervals.

Figure 11. Age-specific Genital Herpes Incidence Rates* by Gender, King County, 2012



* Age-specific crude incidence rate with 95% confidence intervals.

Figure 12. Genital Herpes Incidence Rates by Gender*, King County, 1993-2012



* Gender-specific crude incidence with 95% confidence intervals.

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2003	0-9	2	0.9	0	0.0	2	1.9
	10-14	91	80.4	7	12.0	84	152.4
	15-19	1765	1,583.4	318	559.8	1447	2,646.9
	20-24	1964	1,548.5	625	977.8	1339	2,128.3
	25-29	852	620.9	400	569.0	452	675.4
	30-34	466	304.3	248	312.2	218	295.8
	35-39	263	178.8	177	235.1	86	119.7
	40-44	124	79.5	79	99.9	45	58.6
	45+	137	21.8	103	34.6	34	10.3
	Missing	21	0.0	9	0.0	12	0.0
	All Ages	5685	317.9	1966	220.6	3719	414.6
2004	0-9	2	0.9	0	0.0	2	1.9
	10-14	107	94.9	8	13.8	99	180.4
	15-19	1877	1,674.8	318	557.5	1559	2,833.1
	20-24	1930	1,491.4	576	884.0	1354	2,107.5
	25-29	953	683.5	399	558.2	554	815.2
	30-34	484	323.9	272	351.7	212	294.0
	35-39	290	198.9	177	236.5	113	159.3
	40-44	158	101.2	111	139.9	47	61.2
	45+	144	22.5	113	37.2	31	9.2
	Missing	33	0.0	13	0.0	20	0.0
	All Ages	5988	332.5	1987	221.4	3991	441.9
2005	0-9	0	0.0	0	0.0	0	0.0
	10-14	92	82.6	8	14.0	84	154.8
	15-19	1773	1,563.8	345	597.4	1428	2,567.1
	20-24	1935	1,480.5	603	916.8	1332	2,051.5
	25-29	1094	768.0	474	650.1	620	891.7
	30-34	478	329.6	229	305.6	249	355.3
	35-39	261	177.4	160	211.4	101	141.5
	40-44	190	122.9	141	179.2	49	64.5
	45+	128	19.6	94	30.3	34	9.9
	Missing	22	0.0	12	0.0	10	0.0
	All Ages	5982	329.6	2066	228.3	3907	429.4
2006	0-9	4	1.8	0	0.0	4	3.7
	10-14	69	62.3	9	15.8	60	111.2
	15-19	1604	1,390.2	272	462.4	1332	2,355.1
	20-24	1837	1,376.2	545	811.2	1292	1,948.8
	25-29	976	655.9	402	529.5	574	787.6
	30-34	503	353.1	249	338.6	254	368.5
	35-39	320	212.0	169	217.3	151	206.3
	40-44	182	119.5	129	165.9	53	71.0
	45+	170	25.4	120	37.6	50	14.3
	Missing	22	0.0	9	0.0	13	0.0
	All Ages	5692	308.5	1904	206.8	3783	409.1
2007	0-9	4	1.8	3	2.6	1	0.9
	10-14	75	68.3	10	17.7	65	121.4
	15-19	1615	1,378.3	281	470.0	1334	2,324.4
	20-24	1801	1,339.4	494	731.5	1307	1,952.6
	25-29	1066	691.2	392	498.5	674	891.5
	30-34	491	345.7	247	337.2	244	354.7
	35-39	317	206.9	174	220.0	143	193.0
	40-44	150	100.4	100	131.0	50	68.4
	45+	189	27.6	127	38.8	62	17.3
	Missing	24	0.0	9	0.0	15	0.0
	All Ages	5744	307.0	1837	196.7	3895	415.5

Continued Table 2. Chlamydia

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2008	0-9	4	1.7	1	0.9	3	2.7
	10-14	86	78.8	9	16.1	77	144.7
	15-19	1710	1,450.3	323	536.2	1387	2,405.1
	20-24	1996	1,488.5	633	941.9	1363	2,037.8
	25-29	1048	659.5	395	487.0	653	839.4
	30-34	511	356.0	246	332.9	265	380.6
	35-39	256	167.8	151	191.9	105	142.2
	40-44	176	119.6	120	159.2	56	78.0
	45+	205	29.3	148	44.3	57	15.6
	Missing	2	0.0	1	0.0	1	0.0
All Ages		5994	317.0	2027	214.7	3967	418.8
2009	0-9	1	0.4	1	0.8	0	0.0
	10-14	58	53.0	6	10.7	52	97.5
	15-19	1690	1,436.3	339	564.0	1351	2,347.2
	20-24	1980	1,483.3	598	894.8	1382	2,073.3
	25-29	1101	680.2	398	482.0	703	886.5
	30-34	439	300.6	225	299.2	214	302.0
	35-39	255	169.5	150	193.7	105	143.9
	40-44	147	100.5	101	134.3	46	64.7
	45+	194	27.2	135	39.5	59	15.9
	Missing	2	0.0	0	0.0	2	0.0
All Ages		5867	307.3	1953	204.9	3914	409.4
2010	0-9	2	0.9	0	0.0	2	1.8
	10-14	63	56.9	4	7.0	59	109.3
	15-19	1509	1,284.1	276	459.1	1233	2,148.2
	20-24	2003	1,542.9	581	885.3	1422	2,215.2
	25-29	1101	685.3	443	542.2	658	833.5
	30-34	561	368.9	251	323.7	310	416.0
	35-39	314	210.5	181	238.3	133	181.7
	40-44	203	137.5	138	182.5	65	90.2
	45+	227	31.1	178	51.0	49	12.9
	Missing	1	0.0	1	0.0	0	0.0
All Ages		5985	309.9	2053	213.4	3931	405.6
2011	0-9	1	0.4	0	0.0	1	0.9
	10-14	61	54.8	8	14.1	53	97.3
	15-19	1527	1,314.9	280	469.5	1247	2,207.6
	20-24	2226	1,787.1	642	1,024.4	1584	2,559.4
	25-29	1113	699.4	483	597.4	630	804.8
	30-34	634	407.5	313	393.8	321	421.7
	35-39	333	228.0	206	277.1	127	177.0
	40-44	234	154.6	167	215.4	67	90.7
	45+	281	37.8	225	63.1	56	14.5
	Missing	4	0.0	3	0.0	1	0.0
All Ages		6416	330.3	2327	240.4	4087	419.3
2012	0-9	0	0.0	0	0.0	0	0.0
	10-14	42	37.6	1	1.7	41	75.0
	15-19	1516	1,305.4	321	540.9	1195	2,104.4
	20-24	2302	1,860.9	716	1,149.0	1586	2,583.5
	25-29	1257	817.7	579	741.1	678	896.9
	30-34	689	434.6	378	466.3	311	401.5
	35-39	387	266.6	245	331.7	142	199.2
	40-44	244	159.7	179	228.6	65	87.2
	45+	322	42.4	251	68.8	71	18.0
	Missing	0	0.0	0	0.0	0	0.0
All Ages		6763	345.6	2670	273.7	4089	416.6

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2003	0-9	0	0.0	0	0.0	0	0.0
	10-14	15	13.2	0	0.0	15	27.2
	15-19	186	166.9	54	95.1	132	241.5
	20-24	309	243.6	180	281.6	129	205.0
	25-29	262	190.9	195	277.4	67	100.1
	30-34	228	148.9	178	224.1	50	67.8
	35-39	205	139.3	177	235.1	28	39.0
	40-44	151	96.9	131	165.7	20	26.0
	45+	147	23.4	136	45.7	11	3.3
	Missing	7	0.0	6	0.0	1	0.0
2004	All Ages	1510	84.4	1057	118.6	453	50.5
	0-9	1	0.5	1	0.9	0	0.0
	10-14	17	15.1	2	3.5	15	27.3
	15-19	237	211.5	58	101.7	179	325.3
	20-24	277	214.1	165	253.2	112	174.3
	25-29	199	142.7	146	204.3	53	78.0
	30-34	172	115.1	150	194.0	22	30.5
	35-39	198	135.8	165	220.5	33	46.5
	40-44	132	84.6	114	143.7	18	23.5
	45+	153	23.9	141	46.4	12	3.6
2005	Missing	6	0.0	3	0.0	3	0.0
	All Ages	1396	77.5	945	105.3	447	49.5
	0-9	0	0.0	0	0.0	0	0.0
	10-14	17	15.3	0	0.0	17	31.3
	15-19	328	289.3	108	187.0	220	395.5
	20-24	428	327.5	239	363.4	189	291.1
	25-29	298	209.2	223	305.8	75	107.9
	30-34	255	175.9	196	261.6	59	84.2
	35-39	237	161.1	210	277.4	27	37.8
	40-44	209	135.2	186	236.4	23	30.3
2006	45+	189	29.0	172	55.4	17	5.0
	Missing	4	0.0	2	0.0	2	0.0
	All Ages	1968	108.4	1336	147.6	629	69.1
	0-9	0	0.0	0	0.0	0	0.0
	10-14	18	16.3	1	1.8	17	31.5
	15-19	352	305.1	113	192.1	239	422.6
	20-24	467	349.9	250	372.1	217	327.3
	25-29	372	250.0	264	347.7	108	148.2
	30-34	256	179.7	190	258.4	66	95.8
	35-39	246	162.9	183	235.3	63	86.1
2007	40-44	213	139.8	173	222.5	40	53.6
	45+	252	37.6	217	68.0	35	10.0
	Missing	8	0.0	7	0.0	1	0.0
	All Ages	2186	118.5	1398	151.9	786	85.0
	0-9	0	0.0	0	0.0	0	0.0
	10-14	14	12.7	3	5.3	11	20.5
	15-19	253	215.9	70	117.1	183	318.9
	20-24	311	231.3	166	245.8	145	216.6
	25-29	281	182.2	183	232.7	98	129.6
	30-34	153	107.7	112	152.9	41	59.6
	35-39	136	88.8	114	144.1	22	29.7
	40-44	131	87.7	99	129.7	32	43.8
	45+	119	17.4	102	31.2	17	4.7
	Missing	10	0.0	4	0.0	6	0.0
	All Ages	1414	75.6	853	91.4	555	59.2

Continued Table 3. Gonorrhea

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2008	0-9	1	0.4	0	0.0	1	0.9
	10-14	18	16.5	1	1.8	17	31.9
	15-19	239	202.7	74	122.8	165	286.1
	20-24	317	236.4	183	272.3	134	200.3
	25-29	219	137.8	145	178.8	74	95.1
	30-34	125	87.1	92	124.5	33	47.4
	35-39	136	89.2	107	136.0	29	39.3
	40-44	93	63.2	75	99.5	18	25.1
	45+	156	22.3	138	41.3	18	4.9
	Missing	0	0.0	0	0.0	0	0.0
2009	All Ages	1304	69.0	815	86.3	489	51.6
	0-9	0	0.0	0	0.0	0	0.0
	10-14	7	6.4	0	0.0	7	13.1
	15-19	139	118.1	56	93.2	83	144.2
	20-24	294	220.2	188	281.3	106	159.0
	25-29	208	128.5	166	201.0	42	53.0
	30-34	117	80.1	99	131.6	18	25.4
	35-39	114	75.8	102	131.7	12	16.4
	40-44	90	61.5	83	110.4	7	9.8
	45+	125	17.5	114	33.4	11	3.0
2010	Missing	0	0.0	0	0.0	0	0.0
	All Ages	1096	57.4	808	84.8	286	29.9
	0-9	0	0.0	0	0.0	0	0.0
	10-14	2	1.8	0	0.0	2	3.7
	15-19	171	145.5	51	84.8	120	209.1
	20-24	303	233.4	207	315.4	96	149.5
	25-29	280	174.3	211	258.2	69	87.4
	30-34	255	167.7	200	257.9	55	73.8
	35-39	148	99.2	132	173.8	16	21.9
	40-44	165	111.8	144	190.5	21	29.2
2011	45+	255	34.9	223	63.9	32	8.4
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	1581	81.9	1168	121.4	411	42.4
	0-9	0	0.0	0	0.0	0	0.0
	10-14	5	4.5	0	0.0	5	9.2
	15-19	143	123.1	39	65.4	104	184.1
	20-24	331	265.7	193	308.0	138	223.0
	25-29	298	187.3	235	290.7	63	80.5
	30-34	184	118.3	146	183.7	38	49.9
	35-39	124	84.9	108	145.3	16	22.3
2012	40-44	112	74.0	96	123.8	16	21.7
	45+	181	24.3	170	47.7	11	2.8
	Missing	1	0.0	1	0.0	0	0.0
	All Ages	1381	71.1	988	102.1	391	40.1
	0-9	0	0.0	0	0.0	0	0.0
	10-14	3	2.7	0	0.0	3	5.5
	15-19	133	114.5	44	74.1	89	156.7
	20-24	346	279.7	247	396.4	99	161.3
	25-29	313	203.6	250	320.0	63	83.3
	30-34	225	141.9	199	245.5	26	33.6

Table 4. P&S Syphilis Cases and Incidence Rate by Gender/Age Group

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2003	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	3	2.7	3	5.3	0
	20-24	5	3.9	5	7.8	0
	25-29	8	5.8	7	10.0	1
	30-34	16	10.4	16	20.1	0
	35-39	15	10.2	15	19.9	0
	40-44	9	5.8	9	11.4	0
	45+	5	0.8	5	1.7	0
	Missing	0	0.0	0	0.0	0
All Ages		61	3.4	60	6.7	1
2004	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	0.9	0	0.0	1
	20-24	13	10.0	13	20.0	0
	25-29	15	10.8	13	18.2	2
	30-34	28	18.7	28	36.2	0
	35-39	32	22.0	32	42.8	0
	40-44	26	16.7	24	30.2	2
	45+	9	1.4	9	3.0	0
	Missing	0	0.0	0	0.0	0
All Ages		124	6.9	119	13.3	5
2005	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	4	3.5	4	6.9	0
	20-24	18	13.8	16	24.3	2
	25-29	13	9.1	13	17.8	0
	30-34	13	9.0	13	17.4	0
	35-39	35	23.8	34	44.9	1
	40-44	34	22.0	34	43.2	0
	45+	18	2.8	17	5.5	1
	Missing	0	0.0	0	0.0	0
All Ages		136	7.5	131	14.5	4
2006	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	2	1.7	2	3.4	0
	20-24	9	6.7	8	11.9	1
	25-29	13	8.7	13	17.1	0
	30-34	18	12.6	18	24.5	0
	35-39	24	15.9	24	30.9	0
	40-44	30	19.7	30	38.6	0
	45+	31	4.6	30	9.4	1
	Missing	2	0.0	2	0.0	0
All Ages		129	7.0	127	13.8	2
2007	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	2	1.7	2	3.3	0
	20-24	9	6.7	9	13.3	0
	25-29	14	9.1	14	17.8	0
	30-34	22	15.5	22	30.0	0
	35-39	28	18.3	28	35.4	0
	40-44	20	13.4	20	26.2	0
	45+	28	4.1	27	8.3	1
	Missing	122	0.0	122	13.1	0
All Ages		123	6.6	122	13.1	1

Continued Table 4. P&S Syphilis

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2008	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	3	2.5	3	5.0	0
	20-24	11	8.2	11	16.4	0
	25-29	16	10.1	16	19.7	0
	30-34	16	11.1	16	21.7	0
	35-39	16	10.5	16	20.3	0
	40-44	25	17.0	25	33.2	0
	45+	38	5.4	37	11.1	1
	Missing	0	0.0	0	0.0	0
All Ages		125	6.6	124	13.1	1
2009	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	0.8	1	1.7	0
	20-24	10	7.5	10	15.0	0
	25-29	12	7.4	12	14.5	0
	30-34	26	17.8	25	33.2	1
	35-39	15	10.0	15	19.4	0
	40-44	21	14.4	20	26.6	1
	45+	27	3.8	26	7.6	1
	Missing	0	0.0	0	0.0	0
All Ages		112	5.9	109	11.4	3
2010	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	2	1.7	2	3.3	0
	20-24	12	9.2	12	18.3	0
	25-29	28	17.4	28	34.3	0
	30-34	27	17.8	27	34.8	0
	35-39	31	20.8	31	40.8	0
	40-44	46	31.2	45	59.5	1
	45+	68	9.3	67	19.2	1
	Missing	0	0.0	0	0.0	0
All Ages		214	11.1	212	22.0	2
2011	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	2	1.7	2	3.4	0
	20-24	12	9.6	12	19.1	0
	25-29	29	18.2	29	35.9	0
	30-34	40	25.7	40	50.3	0
	35-39	35	24.0	34	45.7	1
	40-44	44	29.1	44	56.8	0
	45+	74	9.9	72	20.2	2
	Missing	0	0.0	0	0.0	0
All Ages		237	12.2	233	24.1	3
2012	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	5	4.3	5	8.4	0
	20-24	16	12.9	15	24.1	1
	25-29	36	23.4	35	44.8	1
	30-34	32	20.2	31	38.2	1
	35-39	31	21.4	31	42.0	0
	40-44	41	26.8	41	52.4	0
	45+	48	6.3	48	13.2	0
	Missing	0	0.0	0	0.0	0
All Ages		210	10.7	206	21.1	3

Table 5. Genital Herpes Cases and Incidence Rate by Gender/Age Group

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2003	0-9	0	0.0	0	0.0	0	0.0
	10-14	2	1.8	0	0.0	2	3.6
	15-19	93	83.4	8	14.1	85	155.5
	20-24	199	156.9	43	67.3	156	248.0
	25-29	139	101.3	49	69.7	90	134.5
	30-34	77	50.3	35	44.1	42	57.0
	35-39	57	38.7	20	26.6	37	51.5
	40-44	30	19.2	10	12.6	20	26.0
	45+	52	8.3	25	8.4	27	8.2
	Missing	3	0.0	2	0.0	1	0.0
2004	All Ages	652	36.5	192	21.5	460	51.3
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	118	105.3	15	26.3	103	187.2
	20-24	197	152.2	48	73.7	149	231.9
	25-29	137	98.3	39	54.6	98	144.2
	30-34	80	53.5	42	54.3	38	52.7
	35-39	54	37.0	26	34.7	28	39.5
	40-44	43	27.5	22	27.7	21	27.4
	45+	63	9.9	19	6.3	44	13.1
2005	Missing	6	0.0	0	0.0	6	0.0
	All Ages	698	38.8	211	23.5	487	53.9
	0-9	2	0.9	1	0.9	1	0.9
	10-14	3	2.7	0	0.0	3	5.5
	15-19	100	88.2	11	19.0	89	160.0
	20-24	233	178.3	56	85.1	177	272.6
	25-29	151	106.0	61	83.7	90	129.4
	30-34	86	59.3	31	41.4	55	78.5
	35-39	65	44.2	28	37.0	37	51.8
	40-44	45	29.1	23	29.2	22	29.0
2006	45+	73	11.2	23	7.4	50	14.6
	Missing	10	0.0	3	0.0	7	0.0
	All Ages	768	42.3	237	26.2	531	58.4
	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	0.9	0	0.0	1	1.9
	15-19	82	71.1	7	11.9	75	132.6
	20-24	226	169.3	47	70.0	179	270.0
	25-29	159	106.8	46	60.6	113	155.0
	30-34	79	55.5	27	36.7	52	75.4
	35-39	60	39.7	25	32.1	35	47.8
2007	40-44	33	21.7	12	15.4	21	28.1
	45+	93	13.9	33	10.3	60	17.1
	Missing	9	0.0	4	0.0	5	0.0
	All Ages	742	40.2	201	21.8	541	58.5
	0-9	2	0.9	0	0.0	2	1.8
	10-14	2	1.8	1	1.8	1	1.9
	15-19	70	59.7	10	16.7	60	104.5
	20-24	159	118.2	45	66.6	114	170.3
	25-29	127	82.3	37	47.1	90	119.0

Continued Table 5. Genital Herpes

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2008	0-9	0	0.0	0	0.0	0	0.0
	10-14	2	1.8	1	1.8	1	1.9
	15-19	77	65.3	9	14.9	68	117.9
	20-24	118	88.0	29	43.2	89	133.1
	25-29	98	61.7	25	30.8	73	93.8
	30-34	61	42.5	19	25.7	42	60.3
	35-39	38	24.9	15	19.1	23	31.1
	40-44	37	25.1	11	14.6	26	36.2
	45+	80	11.4	18	5.4	62	17.0
	Missing	4	0.0	2	0.0	2	0.0
2009	All Ages	515	27.2	129	13.7	386	40.8
	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	0.9	0	0.0	1	1.9
	15-19	59	50.1	9	15.0	50	86.9
	20-24	136	101.9	32	47.9	104	156.0
	25-29	122	75.4	37	44.8	85	107.2
	30-34	68	46.6	31	41.2	37	52.2
	35-39	42	27.9	19	24.5	23	31.5
	40-44	33	22.6	10	13.3	23	32.4
	45+	80	11.2	25	7.3	55	14.8
2010	Missing	0	0.0	0	0.0	0	0.0
	All Ages	541	28.3	163	17.1	378	39.5
	0-9	1	0.4	0	0.0	1	0.9
	10-14	3	2.7	0	0.0	3	5.6
	15-19	64	54.5	9	15.0	55	95.8
	20-24	155	119.4	35	53.3	120	186.9
	25-29	123	76.6	34	41.6	89	112.7
	30-34	70	46.0	26	33.5	44	59.1
	35-39	48	32.2	23	30.3	25	34.2
	40-44	39	26.4	12	15.9	27	37.5
2011	45+	98	13.4	32	9.2	66	17.3
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	601	31.1	171	17.8	430	44.4
	0-9	1	0.4	0	0.0	1	0.9
	10-14	1	0.9	0	0.0	1	1.8
	15-19	70	60.3	11	18.4	59	104.4
	20-24	164	131.7	43	68.6	121	195.5
	25-29	128	80.4	41	50.7	87	111.1
	30-34	86	55.3	32	40.3	54	70.9
	35-39	53	36.3	21	28.2	32	44.6
2012	40-44	48	31.7	19	24.5	29	39.3
	45+	91	12.2	28	7.9	63	16.3
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	642	33.0	195	20.1	447	45.9
	0-9	1	0.4	0	0.0	1	0.9
	10-14	0	0.0	0	0.0	0	0.0
	15-19	80	68.9	7	11.8	73	128.6
	20-24	189	152.8	41	65.8	148	241.1
	25-29	167	108.6	46	58.9	121	160.1
	30-34	92	58.0	35	43.2	57	73.6