



**Dental Quality Assurance Commission
General Anesthesia Permit Office
On-Site Inspection Form**

Name of Practitioner Inspected: _____

General Anesthesia Permit Number: _____

Location of Inspection (Address):

Telephone Number: _____

Date of Inspection: _____

Time of Inspection: _____

Note: All office equipment and records related to patient care should be available for inspection by the evaluating practitioner. **This inspection is not intended to be punitive; it is an educational tool.**

Part I

Office Equipment, Records, and Emergency Medications

A. Dental Practitioner

1. YES/NO/NA -- BLS Certificate Expiration Date: _____
2. YES/NO/NA -- ACLS Certificate Expiration Date: _____
3. YES/NO/NA -- PALS Certificate Expiration Date: _____
4. YES/NO/NA -- Documentation of Advanced Sedation or Anesthesia Training
5. YES/NO/NA -- Board Certification

B. Staff - Verify appropriate valid BLS/ACLS/PALS/Dental Assistant Registration/Expanded Function Dental Auxiliary License/Dental Hygiene License/Dental Anesthesia Assistant Certification/Anesthesia Monitor Training (14 hours) as applicable for each employee that assists and/or monitors in general anesthesia cases.

Name	Licenses/Certifications Verified	Type
_____	Yes/No/NA	_____

(Attach additional list if necessary)

C. Records

Review three charts of patients who have been treated with general anesthesia in the inspected practice, previously selected by the inspected practitioner (note compliance by checking the entry line).

YES/NO -- Medical history of the patients taken.

YES/NO -- Physical evaluation of the patients completed.

YES/NO -- Anesthesia records show continuous heart rate, blood pressure, respiration, oxygen saturation, and expired CO2 monitoring using electrocardiographic monitoring, pulse oximetry, and capnography as mandated by [WAC 246-817-770](#).

YES/NO -- Recording of blood pressure, heart rate, and respiration rate monitoring every five minutes.

YES/NO -- Documentation of compliance with [WAC 246-817-776](#) discharge criteria (to include vital signs and mental condition, with documentation noting that condition).

YES/NO -- Accurate recording of medications administered, including amounts and times administered.

YES/NO -- Documentation of length of procedure.

YES/NO -- Documentation of any complications of anesthesia.

D. Office Facility and Equipment

YES/NO -- Non-invasive Blood Pressure Monitors

Test/Calibration Dates: _____

YES/NO -- Electrocardiographs/AED or Defibrillator

Test/Calibration Dates: _____

YES/NO -- Pulse Oximeters

Test/Calibration Dates: _____

YES/NO -- End-tidal Carbon Dioxide Monitors

Test/Calibration Dates: _____

YES/NO -- Emergency Oxygen

Test/Calibration Dates: _____

Operating Theater

YES/NO -- Operating theater is large enough to adequately accommodate the patient on a table or in an operating dental chair.

YES/NO -- The operating theater permits an operating team to move freely about the patient.

Operating Chair or Table

YES/NO -- The operating chair or table permits the patient to be positioned so the operating team can maintain the airway.

YES/NO -- The operating chair or table permits the team to alter the patients position quickly in an emergency.

YES/NO -- The operating chair or table provides a firm platform for the management of cardiopulmonary resuscitation.

Lighting System

YES/NO -- The lighting system permits evaluation of the patient's skin and mucosal color.

YES/NO -- There is a battery powered backup light system in place.

YES/NO -- The backup lighting system is of sufficient intensity to permit completion of any operation underway at the time of general power failure.

Gas Storage

YES/NO -- Medical gases are stored in a secure, well-ventilated area and are properly labeled.

YES/NO -- Gas cylinders have steadying devices/brackets in place where applicable.

Suction Equipment

YES/NO -- The suction equipment permits aspiration of the oral and pharyngeal cavities.

YES/NO -- There is a back-up suction device available.

Oxygen Delivery System

YES/NO -- The oxygen delivery system has adequate full-face masks and appropriate connectors and is capable of delivering oxygen to the patient under positive pressure.

YES/NO -- There is an AMBU-bag, self-inflating resuscitator available for emergency use.

YES/NO -- There is an adequate backup-oxygen delivery system.

Recovery Area (can be the operating theater)

YES/NO -- The recovery area has available oxygen.

YES/NO -- The recovery area has available, adequate suction.

YES/NO -- The recovery area has adequate lighting.

YES/NO -- The recovering patient can be observed by a member of the staff at all times during the recovery period.

Ancillary Equipment

YES/NO -- There are oral and nasopharyngeal airways.

YES/NO -- There are laryngeal mask airways.

YES/NO -- There is a tonsillar or pharyngeal type suction tip available to all office outlets.

YES/NO -- There is a sphygmomanometer and stethoscope.

YES/NO -- There is equipment for the establishment and maintenance of an intravenous infusion.

YES/NO -- There are endotracheal tube forceps.

YES/NO -- There are endotracheal tubes and appropriate connectors.

YES/NO -- There is a working laryngoscope complete with an adequate selection of blades, spare batteries, and bulbs.

E. Drugs

YES/NO -- Emergency drug/crash cart is quickly accessible for emergencies. Drugs are stored in a secure location per DEA regulations, and there is an area to aseptically prepare medications.

YES/NO -- Vasopressor drug (alpha- and beta- adrenergic stimulant)

Expiration Date: _____

YES/NO -- Corticosteroid drug

Expiration Date: _____

YES/NO -- Bronchodilator drug

Expiration Date: _____

YES/NO -- Muscle relaxant drug

Expiration Date: _____

YES/NO -- ACLS/PALS medications/drugs for treatment of cardiopulmonary arrest

(expiration dates):

YES/NO -- Adenosine _____

YES/NO -- Amiodarone _____

YES/NO -- Atropine _____

YES/NO -- Epinephrine _____

YES/NO -- Lidocaine _____

YES/NO -- Magnesium _____

YES/NO -- ASA Aspirin _____

YES/NO -- Narcotic antagonist drug

Expiration Date: _____

YES/NO -- Benzodiazepine antagonist drug

Expiration Date: _____

YES/NO -- Antihistamine drug

Expiration Date: _____

YES/NO -- Antiarrhythmic drug

Expiration Date: _____

YES/NO -- Anticholinergic drug

Expiration Date: _____

YES/NO -- Coronary artery vasodilator drug

Expiration Date: _____

YES/NO -- Antihypertensive drug

Expiration Date: _____

YES/NO -- Malignant hyperthermia drug-Dantrolene

Expiration Date: _____

YES/NO -- Antihypoglycemic agent

Expiration Date: _____

YES/NO -- Anticonvulsant drug

Expiration Date: _____

Additional Offices (Satellite Offices):

(Attach additional list if necessary)

I, _____ (name of practitioner being inspected),
declare under penalty of perjury under the laws of the state of Washington that the above listed offices
have equipment, drugs, facilities, and systems that are equally prepared to safely treat patients and
respond to emergency situations as the office location that is being inspected today:

Signature of practitioner being inspected: _____

Date: _____

Overall Equipment and Facility

Comments:

Recommendations/Remediation:

Part II

Simulated Emergencies

The inspectors and the anesthesia team should not just talk about emergency situations and how they should be managed. The practitioner being inspected, and their team should demonstrate their methods for managing the following situations:

YES/NO -- Laryngospasm

YES/NO -- Bronchospasm

YES/NO -- Emesis and aspiration

YES/NO -- Airway obstruction

YES/NO -- Angina/Acute myocardial infarction

YES/NO -- Cardiopulmonary resuscitation (CPR)-(ACLS/PALS protocols)

- Bradycardia
- Ventricular tachycardia (VT) Ventricular fibrillation (VF) Asystole
- Pulseless electrical activity (PEA)

YES/NO -- Hypotension

YES/NO -- Hypertension

YES/NO -- Venipuncture complications

YES/NO -- Neurocardiogenic (vasovagal) syncope

YES/NO -- Allergic reaction

YES/NO -- Local anesthetic toxicity

YES/NO -- Hyperventilation syndrome

YES/NO -- Seizures

YES/NO -- Malignant hyperthermia (MH).

An exact simulation of the emergency situation should be demonstrated in the surgery area, with full participation of the office staff. The “patient” should be positioned and draped, and all the equipment (such as mouth prop, anesthetic machine, and suction apparatus) that may be used should be demonstrated. A simulated IV line should be taped into position, and all emergency equipment should be present, including syringes and medications.

Protocols for Inspection of Simulated Emergencies

Simulated Emergencies and their Algorithms

Respiratory

Laryngospasm

YES/NO -- Administer 100% oxygen via nasal mask.

YES/NO -- Suction the oropharynx, hypopharynx, and nasopharynx with a tonsil suction tip.

YES/NO -- Suction/remove all blood, saliva, and foreign material from the oral cavity.

YES/NO -- Pack the surgical site to prevent bleeding into the hypopharynx.

YES/NO -- Draw the tongue and/or mandible forward.

YES/NO -- Depress the patient's chest while listening with the ear close to the patient's mouth for a rush of air. If a clear "huff" of air is heard, the airway is patent and the spasm probably is resolved.

If a clear "huff" is not heard:

YES/NO -- Try to break the spasm mechanically by attempting to ventilate the patient with a full-face mask and 100% oxygen.

YES/NO -- Administer an IV dose of succinylcholine (partial spasm: 10-20 mg IV; complete spasm 20-40 mg IV) immediately and oxygen under pressure (alternative: rocuronium 0.6-1.2 mg/kg IV; may require prolonged ventilator support).

YES/NO -- Administer intubating IV dose of succinylcholine, intubate, and secure the airway (Alternative: rocuronium 0.6-1.2 mg/kg IV and be prepared to ventilate patient for prolonged period). Consider atropine with repeat dose of succinylcholine to prevent bradycardia.

Bronchospasm

Awake patient:

YES/NO -- Administer 4-8 puffs of a beta-agonist inhalant either via inhaler or nebulizer (2-4 puffs for pediatric patient).

YES/NO -- Administer supplemental oxygen via face mask.

Obtunded non-intubated patient:

YES/NO -- Consider albuterol nebulizer via face mask prior to administration of epinephrine.

YES/NO -- Administer 0.3 to 0.5 mg epinephrine (1:1,000 solution) intramuscularly.

If hypotension is present and believed to be a sign of acute anaphylaxis:

YES/NO -- Administer intravenous bolus dose of 10-20 mcg (0.1-0.2mg) of a 1:10,000 solution of epinephrine titrated to response.

Intubated patient:

YES/NO -- Administer 100% oxygen with positive pressure ventilation.

YES/NO -- Administer 5-10 puffs of a beta agonist inhalant from an inhaler into the open end of the endotracheal tube or through a medication elbow (chamber) added to the circuit designed for inhalant medication delivery.

YES/NO -- Consider deepening plane of anesthesia.

YES/NO -- Check breathing circuit.

YES/NO -- If persistent, consider epinephrine as above for non-intubated patient.

Hypoxemia in spontaneously breathing patient:

YES/NO -- Administer 100% oxygen by face mask.

YES/NO -- Consider reversal of sedative medications.

YES/NO -- Perform chest auscultation and treat findings.

YES/NO -- Intubate and administer positive-pressure ventilation with 100% oxygen if the patient continues to deteriorate with hypoxemia, impending respiratory muscle exhaustion, or worsening obtundation.

Emesis and Aspiration

YES/NO -- Place the patient in the Trendelenburg position with the head down at least 15 degrees and rolled onto the right side.

YES/NO -- Clear any vomitus in the oropharynx with finger sweeps and large-bore suction.

YES/NO -- If no improvement, place the patient in the supine position and intubate (may be necessary to use muscle relaxants prior to intubation).

YES/NO -- Remove any solid particles in the laryngeal region with Magill forceps at time of intubation.

YES/NO -- Oxygenate the patient as soon as possible and manage any bronchospasm as previously described.

YES/NO -- Tracheobronchial lavage is not performed except for small volumes (5-10 mL) of saline to facilitate suctioning.

YES/NO -- Maintain a patient airway and continue ventilation with 100% oxygen in route to hospital.

Difficult Airway Algorithm

YES/NO -- Chin lift/jaw thrust.

YES/NO -- Pull tongue forward, reposition airway.

YES/NO -- Full-face mask positive pressure ventilation.

YES/NO -- Consider nasal/oral pharyngeal airways.

YES/NO -- Consider laryngeal mask airway (LMA) or other adjunct airway devices.

- YES/NO -- Consider endotracheal intubation.
- YES/NO -- Consider needle cricothyrotomy.
- YES/NO -- Consider surgical cricothyrotomy.
- YES/NO -- Consider tracheostomy.

Cardiovascular

Angina/Acute Myocardial Infarction

- YES/NO -- Terminate the surgery and all stimulation of the patient.
- YES/NO -- Place the patient in a comfortable position and loosen tight clothing.
- YES/NO -- Administer oxygen at 4 L/min via mask or nasal cannula.
- YES/NO -- Apply pulse oximeter and ECG monitor.
- YES/NO -- Obtain vital signs with automatic or standard blood pressure cuff.
- YES/NO -- Administer nitroglycerin sublingually or via spray every five minutes if the systolic blood pressure is greater than 90mm/Hg.
- YES/NO -- Establish IV access, if not already present.
- YES/NO -- Administer non-enteric aspirin (160-325 mg) and direct patient to chew one and swallow one, except in cases of allergy to this drug.
- YES/NO -- Employ the mnemonic MONA but not in that order; use ONAM because use of chewed aspirin may decrease absorption of subsequent nitroglycerin.
- YES/NO -- Administer morphine, 1-3 mg IV every five minutes to control pain if systolic blood pressure is greater than 90mm/Hg. Watch carefully so the patient does not become overly sedated or lose respiratory drive.
- YES/NO -- Monitor vital signs closely. Watch for hypotension, hypoxemia, bradycardia, ventricular dysrhythmias, and cardiac arrest and manage appropriately per ACLS protocols.

Management of Perioperative Hypotension

- YES/NO -- Treatment of hypotension is directed toward its cause.
- YES/NO -- Decrease anesthetic depth if patient is under anesthesia.
- YES/NO -- Terminate the surgery.
- YES/NO -- Place the patient in a supine position with legs elevated.
- YES/NO -- Administer 100% oxygen.
- YES/NO -- Re-evaluate blood pressure, heart rate, and rhythm and dysrhythmias.
- YES/NO -- Administer a bolus of isotonic fluid.
- YES/NO -- Titrate ephedrine or phenylephrine or other suitable vasopressor to preserve adequate systemic pressures (Ephedrine is generally used unless there is tachycardia and hypotension, in which case phenylephrine may be a better choice).
- YES/NO -- Dosing of sympathomimetic agents for treatment of hypotension.

Emergencies

YES/NO -- Ephedrine: Dilute a 50 mg/mL vial in 9mL of saline to make a 5mg/mL solution and administer at a dose of 2.5 to 5 mg IV, which can be repeated until blood pressure is stabilized. The drug's effects occur in approximately one minute, peak at fifteen minutes, and last approximately one hour.

YES/NO -- Phenylephrine: Dilute single dose vial (10 mg/mL) in 9mL of saline; discard 9mL and dilute with an additional 9mL of saline to create a 0.1mg/mL concentration; administer IV in 0.1 mg (1 mL) increments until the desired effects are achieved. The effects are seen within one minute and last for approximately twenty minutes.

Hypertension

Therapeutic approaches to treating perioperative hypertension are directed at producing vasodilation or altering cardiac output by beta-adrenergic receptor blockers. Among the drugs that may be useful are beta-blockers such as esmolol (Brevibloc) and metoprolol (Lopressor), the selective alpha blocker labetalol (Normodyne, Trandate), and the vasodilator hydralazine.

YES/NO -- Esmolol dosing.

Immediate control dosing: 80 mg (approximately 1 mg/kg) over 30 seconds followed with a 150 mcg/kg/min infusion (0.15/mg/kg/min) that is adjusted as required to a maximum of 300 mcg/kg/min (0.30 mg/kg/min) to maintain the desired heart rate and/or blood pressure.

Graduate control dosing (when there is time to titrate): loading dose infusion of 500 mcg/kg/min for one minute followed with a 50 mcg/kg/min over four minutes. If an adequate therapeutic effect is not seen in five minutes, loading dose infusion of 500 mcg/kg/min for one minute followed with a 100 mcg/kg/min.

YES/NO -- Labetalol dosing (a selective alpha 1 blocker and non-selective beta blocker):

Initial adult dose 5-20 mg IV over two minutes followed by 2 mg/min (maximum dose 300 mg) IV infusion. Onset five minutes, duration three to six hours.

YES/NO -- Hydralazine dosing (a direct vasodilator for a patient who can tolerate an increase in heart rate and cardiac output but who has a history of asthma):

Initial adult dose 5-20mg IV, can titrate up to 25 mg. Onset five minutes; duration two hours.

Venipuncture Complications

Hematoma

YES/NO -- Apply pressure to the venipuncture site.

Extravasation

YES/NO -- Apply moist heat and elevate extremity above level of heart (in many cases this simple treatment is all that is required).

If irritating drugs have extravasated, then consider:

YES/NO -- Infiltrate 1% plain lidocaine (Xylocaine) at the site, if needed for pain.

YES/NO -- Administer ibuprofen; possible use of steroids and antibiotics may be necessary to prevent or lessen the sloughing.

Phlebitis

YES/NO -- Apply heat to area.

YES/NO -- Administer ibuprofen or other nonsteroidal drugs, steroids, and antibiotics.

YES/NO -- Limit motion to alleviate symptoms.

YES/NO -- Seek appropriate consultation if severe symptoms persist beyond three to four days.

Intra-Arterial Injection

Prevention is the best treatment. No single treatment regimen has been found to be completely effective.

YES/NO -- Leave needle in place.

YES/NO -- Inject 10 mL 1% plain lidocaine into the artery.

YES/NO -- Consider transferring patient to a hospital for further therapy and consider vascular consultation.

Neurocardiogenic (Vasovagal) Syncope

YES/NO -- Place patient in reclining position with legs elevated.

YES/NO -- Maintain airway.

YES/NO -- Deliver 100% oxygen.

YES/NO -- Support respiration if needed.

YES/NO -- Take vital signs.

YES/NO -- Apply cool compress to forehead.

YES/NO -- Administer spirits of ammonia, if necessary.

YES/NO -- Administer IV atropine 0.5 mg every three to five minutes up to 3 mg in the presence of bradycardia.

Hyperventilation Syndrome

YES/NO -- Maintain adequate oxygen levels while reducing carbon dioxide elimination.

If using oxygen delivery system with full-face mask, closed circuit, and no carbon dioxide absorber:

YES/NO -- Deliver oxygen at 600 mL/min.

YES/NO -- Have patient breathe into the circuit until the rebreathing bag is distended.

YES/NO -- Continue until episode subsides.

If using anesthesia reservoir or paper bag:

YES/NO -- Have patient exhale and inhale into the paper bag 6-10 times/min.

If a non-sedated patients fails to respond:

YES/NO -- Administer suitable sedative to abort the hyperventilation episode.

Seizures

YES/NO -- Prevent injury to the uncontrolled unconscious patient.

YES/NO -- Loosen clothing about the neck.

YES/NO -- Place pillow under the head.

YES/NO -- Place padded tongue blade between the teeth if evidence that the patient's tongue is being traumatized (do not force the teeth apart).

YES/NO -- Consider checking blood sugar.

Local Anesthetic Toxicity

YES/NO -- Immediately discontinue any further administration of the local drug.

YES/NO -- Call 911.

YES/NO -- Place patient in supine position.

YES/NO -- Administer oxygen.

YES/NO -- Maintain airway and initiate CPR.

YES/NO -- Confirm or establish IV access.

YES/NO -- Administer IV diazepam 5 mg over one minute OR IV midazolam 2.5 mg over one minute titrated to effect.

YES/NO -- Continue to closely monitor vital signs.

Allergic Reaction

Mild reactions usually manifest as skin reaction without other systemic signs or symptoms:

YES/NO -- Identify causative agent, if time permits.

YES/NO -- Remove, dilute, negate, or antagonize, if possible.

YES/NO -- Administer diphenhydramine adult dose 25-50 mg PO, 10-50 mg IV at a rate generally not exceeding 25 mg/min, deep intramuscular injection of 100 mg if required (maximum daily dosage is 400 mg) **OR**

YES/NO -- Chlorpheniramine maleate adult dose 5-20 mg, IV, IM, or SQ injection as a single dose; tablets or syrup, 4 mg every four to six hours.

Severe Reactions (Anaphylaxis) manifest with all signs of symptoms of allergic reaction with skin rash, watery eyes and nose, abdominal cramps, wheezing, tachycardia, and hypotension.

It is important to have a plan or algorithm to guide treatment of a severe allergic reaction and impending cardiovascular collapse.

YES/NO -- Stop administration of the antigen.

Follow the ABCs:

YES/NO -- Maintain an airway with supplemental oxygen.

YES/NO -- Support respiration and breathing.

YES/NO -- Support circulation by providing an IV route for volume loading with crystalloid solution. In the adult patient, rapidly infuse 1 L lactated Ringer's solution. For children, administer fluid boluses of 20 mL/kg of lactated Ringer's solution or normal saline.

YES/NO -- Administer epinephrine.

Intravenous: Use a premixed solution of 1:10,000 (1 mg of epinephrine in 10 mL).

- Adults: Titrate 0.2 mg (2 mL) to 0.5 mg (5 mL) to effect and repeat every two to five minutes as needed (use with caution in patient with cardiovascular disease).
- Children: 0.01 mg/kg.

Intramuscular or Subcutaneous: If an IV route is unavailable, the drug can be administered IM or subcutaneously using a 1:1,000 (1 mg/mL).

- Adults: 0.3 to 0.5 mg of a 1:1,000 concentration. This dose may be repeated at 10 to 20 minute intervals based on patient response.
- Children: 0.01 mg/kg. This dose may be repeated at 10 to 20 minute intervals based on patient response.

YES/NO -- As epinephrine administration continues, H1-antihistamine may be administered IM or IV. Diphenhydramine (Benadryl) can be given IV in a dose of 50 mg for adults and older children. A child six to twelve years of age receives 25 mg. An H2 blocker, such as famotidine (Pepcid), can be administered to adults IV starting at 20 mg. The dose for children older than one year is 0.5mg/kg IV up to 40 mg per day.

YES/NO -- Albuterol inhaler can be administered at 4-5 puffs as needed (2-4 puffs for pediatric patient).

YES/NO -- Administer corticosteroids: The use of corticosteroids has been advocated in the past to treat asthma, edema, or intense pruritus. Although these drugs act too slowly to be lifesaving in an acute anaphylactic episode, they are important in combating immunologic reactions and regaining homeostasis. Because of their slow action, they are the last drugs administered in the therapeutic regimen. Options include the following:

- Hydrocortisone sodium succinate (Solu-Cortef) at a dose of 100 mg IV given slowly over one minute.
- Dexamethasone (Decadron) at a dose of 4 to 12 mg given slowly over one minute IV or IM.
- Doses can be repeated at intervals as determined by the severity of symptoms. Because anaphylaxis is an acute process, corticosteroid use normally can be stopped after 36 hours without adverse sequelae.

YES/NO -- Follow-up Measures:

Pharyngeal edema and upper airway obstruction can occur and persist after the administration of epinephrine. Endotracheal intubation or tracheotomy may be necessary to secure an airway. If shock continues, the patient should be placed in a slight Trendelenburg position to provide circulatory support. If the patient develops dyspnea and wheezing, a semi-reclining position may be more helpful. If a significant decrease in blood pressure persists despite appropriate volume expansion and the use of epinephrine, an adjunctive vasopressor should be considered (see section on treatment of hypotension for management of the problem).

Seizures may occur during allergic reactions and usually are due to circulatory and/or respiratory inadequacy. If convulsions persist after corrective measures have been taken to ensure cardiorespiratory sufficiency, diazepam, 1 mL (5 mg), can be administered IV for one minute, with midazolam (2.5 mg over one minute) used as an alternative. It is important to wait two minutes for the effect before giving another dose.

After successful treatment of a severe allergic reaction, the patient should be hospitalized and observed for 24 hours, receiving prophylactic antihistamines. Anaphylactic reactions have been known to recur after the initial effects of the counteracting drugs have worn off.

Malignant Hypothermia

The following are the primary signs of impending MH:

- Increase in end-tidal carbon dioxide (doubling or tripling may occur rapidly or during 10 to 20 minutes).
- Unexplained tachycardia, tachypnea, hypercarbia.
- Generalized muscle rigidity.
- Masseter muscle rigidity.
- Hyperthermia (often a late sign).
- Respiratory and/or metabolic acidosis.
- Sudden/unexpected cardiac arrest (consider as secondary to hyperkalemia and treat as such).

Contact telephone number **1-800-MHHYPER (1-800-644-9737)**.

Emergency Treatment (Caution: These protocols may not apply to every patient and may require alteration according to specific patient needs).

Acute Phase:

YES/NO -- Immediately activate the EMS for patient transport and notify receiving hospital to prepare for continuing treatment of an MH patient.

YES/NO -- Immediately discontinue use of all volatile inhalation anesthetics and succinylcholine. Hyperventilate with 100% oxygen at high gas flows, at least 10 L/min. The circle system and carbon dioxide absorbent need not be changed.

YES/NO -- Solubilize the Dantrolene according to manufacturer's instructions. If using the older formula (Dantrolene), it takes time to mix the solution, so it is recommended that one person be designated for this task.

YES/NO -- Administer Dantrolene, 2 to 3 mg/kg initial bolus rapidly, with increments up to a total of 10 mg/kg. Continue to administer Dantrolene until signs of MH (e.g., tachycardia, rigidity, increased end-tidal carbon monoxide, and temperature elevation) are controlled. Occasionally, a total dose greater than 10 mg/kg may be needed.

YES/NO -- Administer bicarbonate to correct metabolic acidosis, as guided by blood gas analysis. In the absence of blood gas analysis, administer 1 to 2 mEq/kg.

YES/NO -- Simultaneously with the previous steps, actively cool the hyperthermic patient with IV cold saline (not Ringer's lactate) 15 mL/kg every fifteen minutes, three times.

- Surface cool with ice and hypothermia blanket. A supply of ice or an ice machine is helpful for this purpose.
- Monitor closely because excessively vigorous treatment may lead to hypothermia.

YES/NO -- Arrhythmias usually respond to treatment of acidosis and hyperkalemia. If they

persist or are life-threatening, use standard antiarrhythmic agents, except calcium channel blockers, which may cause hyperkalemia and cardiovascular collapse.

- YES/NO -- Determine and monitor end-tidal carbon dioxide concentrations; arterial, central, or femoral venous blood gases; serum potassium and calcium levels; clotting studies; and urine output.
- YES/NO -- Treat hyperkalemia with hyperventilation, bicarbonate, and IV glucose and insulin (10 U of regular insulin in 50 mL of 50% glucose titrated to potassium level or 0.15 U/kg of regular insulin in 1 mL/kg of 50% glucose). Life-threatening hyperkalemia also may be treated with calcium (e.g., 2-5 mg/kg of calcium chloride).
- YES/NO -- Ensure urine output of greater than 2 mL/kg/h by hydration and/or administration of furosemide and additional mannitol, if necessary. Consider central venous catheter placement for fluid resuscitation and electrolyte monitoring.
- YES/NO -- Sudden unexpected cardiac death (SUCCD) is a leading cause of potential fluid shifts and electrolyte instability. Children younger than 1 year old experience sudden cardiac arrest after succinylcholine administration in the absence of hypoxemia and anesthetic overdose should be treated initially for acute hyperkalemia. In this situation, calcium chloride should be administered with other means to reduce serum potassium levels. Affected children should be presumed to have subclinical muscular dystrophy, and a neurologist should be consulted.

Part III

Discussion

This part of the inspection should be conducted in private with the practitioner being inspected.

YES/NO -- Completed Simulated Emergencies

Comments: _____

Recommendations/Remediation: _____

Signatures of inspectors:

1. _____
2. _____

Printed names of inspectors:

1. _____
2. _____

License/Permit number of inspectors:

1. _____
2. _____

Date: _____

Signature of practitioner being inspected:

Printed name of practitioner being inspected:

Date: _____

The practitioner being inspected must maintain an original or copy of this inspection for **five years**, per [WAC 246-817-774](#).