


Sexually Transmitted Disease Profile

Yakima County 2014



Disease Control and Health Statistics
Infectious Disease Assessment Unit



Sexually Transmitted Disease Profile

Yakima County 2014



For persons with disabilities, this document is available in other formats. To submit a request, please call 1-800-525-0127 (TDD/TYY 1-800-833-6388).

Washington State Department of Health
Disease Control and Health Statistics
Infectious Disease Assessment Unit
Olympia, Washington
(360) 236-3455

Contents

Introduction.....1

Data Sources, Definitions, and Limitations1

County STD Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, 2014.4

Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates per 100,000 population, 1995-20144

Figure 2. Age-specific Chlamydia Incidence Rates by Gender, 20145

Figure 3. Chlamydia Incidence Rates by Gender, 1995-20145

Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates per 100,000 population, 1995-20146

Figure 5. Age-specific Gonorrhea Incidence Rates by Gender, 20146

Figure 6. Gonorrhea Incidence Rates by Gender, 1995-20147

Primary & Secondary Syphilis

Figure 7. Primary & Secondary Syphilis Cases and Incidence Rates, 1995-2014.7

Figure 8. Age-specific Primary & Secondary Syphilis Incidence Rates by Gender, 20148

Figure 9. Primary & Secondary Syphilis Incidence Rates by Gender, 1995-2014.....8

Genital Herpes

Figure 10. Genital Herpes Cases and Incidence Rates per 100,000 population 1995-2014.....9

Figure 11. Age-specific Genital Herpes Incidence Rates by Gender, 20149

Figure 12. Genital Herpes Incidence Rates by Gender, 1995-2014.10

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group, 2005-201411

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group, 2005-201413

Table 4. Primary & Secondary Syphilis Cases and Incidence Rates by Gender and Age
Group, 2005-201415

Table 5. Genital Herpes Cases and Incidence Rates by Gender and Age Group, 2005-2014 ..17

Introduction

Sexually transmitted infections continue to be the most frequently diagnosed and reported notifiable conditions in Washington State. This report describes the sexually transmitted infection burden in Yakima County. Data are presented for the more commonly reported diseases of chlamydial infection, gonorrhea, primary and secondary syphilis, and genital herpes when at least twenty (20) cases of disease were diagnosed in 2014 and the corresponding rates are statistically stable. The incidence rates, including those by age and sex, are presented graphically. The report concludes with tables containing a decade of historical data by age group and gender.

Data Sources, Definitions and Limitations

Cases: Surveillance cases are the number of new episodes of disease (not unique persons) diagnosed in a given year. Cases are identified and submitted by health care providers to local health jurisdictions and entered into the Washington State Department of Health Public Health Information Management System – Sexually Transmitted Diseases (PHIMS-STD) data system. To be included in surveillance reporting, each case must meet disease definitions (see below). Data presented in this report represent new cases of infection diagnosed during a given year and reported as of March 2, 2015.

Confidence Intervals: A confidence interval is a range around a measurement that suggests how precise the measurement is. The confidence interval displays both the possible range around the estimate and also how stable the estimate is. Wider confidence intervals in relation to the estimate itself indicate instability. A confidence interval is always qualified by a particular confidence level, usually expressed as a percentage. In this report, 95% confidence intervals are used.

Disease Definitions:

- | | |
|----------------------|--|
| Chancroid | – A sexually transmitted infection caused by the bacterium <i>Haemophilus ducreyi</i> that may include the symptoms of painful genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of <i>H. ducreyi</i> from a clinical specimen. |
| Chlamydia (CT) | – A sexually transmitted infection caused by the bacterium <i>Chlamydia trachomatis</i> that may include the symptoms of swelling and pain of internal sexual organs, however the infection often has no symptoms in women. Cases are defined by laboratory detection of <i>C. trachomatis</i> from a clinical specimen. |
| Genital Herpes (HSV) | – A sexually transmitted infection caused by the herpes simplex viruses type 1 and type 2 that may include the symptoms of blisters or sores in the genital area. Cases are defined by laboratory detection of herpes simplex virus (HSV1 or HSV2) or positive antibody response from a clinical specimen. Reportable cases include only adult genital initial infection and neonatal infection. |

- Gonorrhea (GC) – A sexually transmitted infection caused by the bacterium *Neisseria gonorrhoeae* that may include the symptoms of swelling and pain of internal sexual organs, however the infection sometimes has no symptoms. Cases are defined by laboratory detection of the bacterium *N. gonorrhoeae* from a clinical specimen.
- Granuloma Inguinale (GI) – A sexually transmitted infection caused by the bacterium *Klebsiella granulomatis* that may include the symptoms of slowly increasing genital sores and swollen pelvic lymph nodes. Cases are defined by microscopic examination of a clinical specimen.
- Lymphogranuloma Venereum (LGV) – A sexually transmitted infection caused by three strains of *Chlamydia trachomatis* that may include the symptoms of genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of the L1, L2 and L3 serovars of *C. trachomatis* from a clinical specimen.
- Syphilis – A sexually transmitted infection caused by the bacterium *Treponema pallidum* that may include many kinds of symptoms or none at all depending upon the stage of disease. Cases are defined and assigned a stage by a combination of positive blood tests, symptoms, and history of previous treatment. The U.S. Centers for Disease Control and Prevention (CDC) provides guidelines with additional details of surveillance definitions and staging criteria. The stages of primary and secondary (P&S) syphilis are grouped together for analysis in this report.
Primary – identified by the presence of one or many painless sores.
Secondary – identified by the presence of a rash on one or more areas of the body, often with fever, fatigue or other symptoms at the same time.
Other Stages – additional stages of syphilis include, early latent, latent, late latent, and congenital. See CDC guidelines for specific criteria: www.cdc.gov/std/

Incidence Rates: Incidence rates in this report are calculated as the number of new episodes of a disease (not unique persons) diagnosed in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates are not presented when there were fewer than 20 cases of disease reported and the calculated relative standard error was greater than thirty percent (>30%) as this indicates statistical instability.

Limitations: The data presented in this report may be subject to a number of limiting factors. Clinically diagnosed cases (without laboratory confirmation) may be missed through public health surveillance systems. Depending upon diagnosing practices, completeness of reporting may vary by the source of health care. In addition, the diagnosing practitioner is responsible for providing the case information including the patient demographic data items of age and gender upon which many of the analyses in this report depend. Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. Also, small

increases or decreases in numbers from year to year can look large if the actual number of cases is small. Care should be taken in interpreting these data in light of known limitations.

Population: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM) April 1, 2014. This data was accessed March 3, 2015 from <http://www.ofm.wa.gov/pop/>

Tabular Data: The data tables are provided in hopes that community and local partners will use these historical data as a resource for future health planning. Users of these data should keep in mind that small numbers often produce rates that can be unstable from year to year and may not be reliable when comparing across strata within a particular year. Data tables for additional years previous are available upon request.

Anyone with specific questions about how these data should be interpreted is encouraged to contact the Infectious Disease Assessment Unit at 360-236-3455.

Yakima County STD Disease Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, Yakima County, 2014


Disease	County Cases	County Rate§	WA State Rate
Chlamydia	1,504	604.5	376.7
Gonorrhea	406	163.2	88.1
P&S Syphilis	15	60.3	4.8
Genital Herpes	60	24.1	29.9
Chancroid/GI/LGV	0		
Total	1,985		

§ Crude incidence rate per 100,000 population.

* Rates are suppressed for counts under 20 with a corresponding RSE >30% due to statistical instability.


Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates* per 100,000 population, Yakima County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 2. Age-specific Chlamydia Incidence Rates* by Gender, Yakima County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 3. Chlamydia Incidence Rates by Gender*, Yakima County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.


Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates* per 100,000 population, Yakima County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 5. Age-specific Gonorrhea Incidence Rates* by Gender, Yakima County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 6. Gonorrhea Incidence Rates by Gender*, Yakima County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.


Primary and Secondary Syphilis

Figure 7. Primary and Secondary Syphilis Cases and Incidence Rates* per 100,000 population, Yakima County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 8. Age-specific Primary and Secondary Syphilis Incidence Rates* by Gender, Yakima County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 9. Primary and Secondary Syphilis Incidence Rates by Gender*, Yakima County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.


Genital Herpes

Figure 10. Genital Herpes Cases and Incidence Rates* per 100,000 population, Yakima County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 11. Age-specific Genital Herpes Incidence Rates* by Gender, Yakima County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.

Figure 12. Genital Herpes Incidence Rates by Gender*, Yakima County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2005	0-9	2	5.0	1	4.9	1	5.1
	10-14	24	120.3	3	29.2	21	217.2
	15-19	347	1,780.7	48	472.0	299	3,209.2
	20-24	341	2,035.8	81	933.9	260	3,219.0
	25-29	149	1,003.4	34	454.1	115	1,562.1
	30-34	63	432.8	17	228.0	46	647.7
	35-39	24	160.5	9	118.0	15	204.7
	40-44	11	70.5	2	25.4	9	116.5
	45+	14	18.4	8	22.1	6	15.1
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	975	420.4	203	174.9	772	666.3
2006	0-9	0	0.0	0	0.0	0	0.0
	10-14	8	40.4	0	0.0	8	83.4
	15-19	346	1,751.9	62	601.1	284	3,009.7
	20-24	430	2,522.9	80	906.6	350	4,257.9
	25-29	195	1,265.5	36	465.1	159	2,073.3
	30-34	71	495.1	17	232.0	54	770.2
	35-39	37	242.2	12	154.0	25	334.0
	40-44	17	111.4	4	51.8	13	172.6
	45+	13	16.8	4	10.8	9	22.3
	Missing	3	0.0	1	0.0	2	0.0
	All Ages	1120	477.8	216	184.1	904	772.3
2007	0-9	2	4.9	1	4.8	1	5.0
	10-14	14	71.3	2	19.8	12	125.9
	15-19	391	1,955.0	62	593.4	329	3,444.3
	20-24	393	2,294.9	76	859.2	317	3,828.5
	25-29	222	1,398.2	43	539.7	179	2,263.0
	30-34	89	620.4	20	273.1	69	982.3
	35-39	34	220.1	14	177.5	20	264.4
	40-44	18	120.9	9	119.2	9	122.6
	45+	16	20.3	8	21.2	8	19.5
	Missing	3	0.0	1	0.0	2	0.0
	All Ages	1182	498.9	236	198.9	946	800.0
2008	0-9	1	2.4	0	0.0	1	4.9
	10-14	11	56.2	2	19.9	9	94.7
	15-19	334	1,657.4	42	398.3	292	3,039.5
	20-24	404	2,363.7	77	873.7	327	3,949.8
	25-29	231	1,416.5	53	647.2	178	2,192.4
	30-34	97	664.7	24	323.0	73	1,019.1
	35-39	47	305.5	11	140.3	36	477.3
	40-44	15	102.6	6	80.8	9	125.1
	45+	18	22.4	10	25.9	8	19.1
	Missing	2	0.0	0	0.0	2	0.0
	All Ages	1165	486.4	225	187.5	935	782.4
2009	0-9	0	0.0	0	0.0	0	0.0
	10-14	10	50.9	0	0.0	10	104.6
	15-19	365	1,817.2	44	418.4	321	3,354.2
	20-24	438	2,571.2	68	774.8	370	4,480.5
	25-29	213	1,286.3	33	397.3	180	2,181.3
	30-34	83	555.1	24	315.4	59	803.5
	35-39	39	257.6	14	182.0	25	335.8
	40-44	21	145.0	8	108.4	13	183.1
	45+	11	13.4	4	10.1	7	16.4
	Missing	1	0.0	0	0.0	1	0.0
	All Ages	1181	488.6	195	160.9	986	817.9

Continued Table 2. Chlamydia

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2010	0-9	0	0.0	0	0.0	0	0.0
	10-14	11	55.4	0	0.0	11	113.8
	15-19	336	1,683.0	42	401.8	294	3,091.5
	20-24	383	2,317.0	63	733.2	320	4,031.8
	25-29	205	1,256.4	45	551.7	160	1,960.5
	30-34	99	634.6	22	280.1	77	994.1
	35-39	40	267.6	16	213.3	24	322.3
	40-44	15	103.6	6	81.7	9	126.1
	45+	22	26.3	2	5.0	20	46.0
	Missing	0	0.0	0	0.0	0	0.0
2011	All Ages	1111	456.8	196	161.1	915	752.7
	0-9	1	2.4	0	0.0	1	4.9
	10-14	12	60.4	0	0.0	12	124.1
	15-19	418	2,121.5	45	436.1	373	3,974.4
	20-24	423	2,555.0	75	871.9	348	4,375.2
	25-29	199	1,224.5	54	663.5	145	1,787.3
	30-34	82	514.4	19	236.8	63	795.9
	35-39	52	356.7	12	164.2	40	550.0
	40-44	19	128.8	7	93.5	12	165.2
	45+	19	22.3	4	9.8	15	33.9
2012	Missing	0	0.0	0	0.0	0	0.0
	All Ages	1225	500.6	216	176.4	1009	825.2
	0-9	4	9.2	1	4.5	3	14.2
	10-14	13	65.5	3	29.2	10	104.4
	15-19	392	2,186.5	66	696.5	326	3,857.1
	20-24	493	3,037.7	101	1,175.8	392	5,131.3
	25-29	200	1,288.5	43	562.8	157	1,991.9
	30-34	104	608.2	28	336.7	76	865.2
	35-39	49	325.7	8	107.9	41	537.4
	40-44	23	153.6	5	67.6	18	237.5
2013	45+	24	28.0	5	12.1	19	42.6
	Missing	1	0.0	0	0.0	1	0.0
	All Ages	1303	529.7	260	211.8	1043	846.2
	0-9	0	0.0	0	0.0	0	0.0
	10-14	12	60.5	2	19.5	10	104.4
	15-19	368	2,130.3	49	530.4	319	3,969.4
	20-24	560	3,508.4	122	1,450.2	438	5,802.2
	25-29	220	1,466.4	44	596.7	176	2,307.0
	30-34	95	542.4	21	249.5	74	813.2
	35-39	58	376.2	15	199.6	43	544.2
2014	40-44	34	226.0	18	245.9	16	207.1
	45+	33	37.9	10	24.0	23	50.7
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	1380	558.1	281	228.2	1099	885.4
	0-9	0	0.0	0	0.0	0	0.0
	10-14	14	69.9	0	0.0	14	144.6
	15-19	424	2,471.3	71	770.8	353	4,442.7
	20-24	547	3,486.7	128	1,545.4	419	5,658.1
	25-29	271	1,850.7	70	968.4	201	2,710.9
	30-34	122	702.0	32	386.3	90	989.5

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2005	0-9	0	0.0	0	0.0	0
	10-14	3	15.0	1	9.7	2
	15-19	38	195.0	9	88.5	29
	20-24	54	322.4	23	265.2	31
	25-29	25	168.4	13	173.6	12
	30-34	11	75.6	5	67.1	6
	35-39	4	26.8	0	0.0	4
	40-44	4	25.6	3	38.0	1
	45+	4	5.3	4	11.0	0
	Missing	0	0.0	0	0.0	0
2006	All Ages	143	61.7	58	50.0	85
	0-9	1	2.5	0	0.0	1
	10-14	1	5.1	0	0.0	1
	15-19	46	232.9	13	126.0	33
	20-24	51	299.2	15	170.0	36
	25-29	31	201.2	12	155.0	19
	30-34	12	83.7	3	40.9	9
	35-39	14	91.7	7	89.9	7
	40-44	8	52.4	7	90.6	1
	45+	3	3.9	1	2.7	2
2007	Missing	2	0.0	0	0.0	2
	All Ages	169	72.1	58	49.4	111
	0-9	0	0.0	0	0.0	0
	10-14	2	10.2	0	0.0	2
	15-19	37	185.0	12	114.9	25
	20-24	31	181.0	8	90.4	23
	25-29	21	132.3	5	62.8	16
	30-34	10	69.7	4	54.6	6
	35-39	4	25.9	0	0.0	4
	40-44	1	6.7	0	0.0	1
2008	45+	6	7.6	5	13.2	1
	Missing	1	0.0	0	0.0	1
	All Ages	113	47.7	34	28.7	79
	0-9	1	2.4	1	4.7	0
	10-14	0	0.0	0	0.0	0
	15-19	20	99.2	3	28.4	17
	20-24	26	152.1	11	124.8	15
	25-29	18	110.4	8	97.7	10
	30-34	8	54.8	2	26.9	6
	35-39	4	26.0	1	12.8	3
2009	40-44	4	27.4	3	40.4	1
	45+	3	3.7	2	5.2	1
	Missing	1	0.0	1	0.0	0
	All Ages	85	35.5	32	26.7	53
	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	13	64.7	2	19.0	11
	20-24	17	99.8	7	79.8	10
	25-29	7	42.3	3	36.1	4
	30-34	1	6.7	0	0.0	1
	35-39	1	6.6	1	13.0	0
	40-44	0	0.0	0	0.0	0
	45+	0	0.0	0	0.0	0
	Missing	0	0.0	0	0.0	0
	All Ages	39	16.1	13	10.7	26

Continued Table 3. Gonorrhea

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2010	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	7	35.1	3	28.7	4	42.1
	20-24	11	66.5	5	58.2	6	75.6
	25-29	9	55.2	6	73.6	3	36.8
	30-34	1	6.4	0	0.0	1	12.9
	35-39	2	13.4	0	0.0	2	26.9
	40-44	3	20.7	0	0.0	3	42.0
	45+	1	1.2	1	2.5	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
2011	All Ages	34	14.0	15	12.3	19	15.6
	0-9	1	2.4	0	0.0	1	4.9
	10-14	0	0.0	0	0.0	0	0.0
	15-19	15	76.1	4	38.8	11	117.2
	20-24	31	187.2	13	151.1	18	226.3
	25-29	21	129.2	9	110.6	12	147.9
	30-34	16	100.4	9	112.2	7	88.4
	35-39	4	27.4	4	54.7	0	0.0
	40-44	3	20.3	2	26.7	1	13.8
	45+	8	9.4	5	12.2	3	6.8
2012	Missing	0	0.0	0	0.0	0	0.0
	All Ages	99	40.5	46	37.6	53	43.3
	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	5.0	0	0.0	1	10.4
	15-19	11	61.4	4	42.2	7	82.8
	20-24	26	160.2	12	139.7	14	183.3
	25-29	19	122.4	9	117.8	10	126.9
	30-34	5	29.2	3	36.1	2	22.8
	35-39	7	46.5	7	94.4	0	0.0
	40-44	7	46.7	3	40.5	4	52.8
2013	45+	5	5.8	2	4.9	3	6.7
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	81	32.9	40	32.6	41	33.3
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	19	110.0	7	75.8	12	149.3
	20-24	60	375.9	34	404.1	26	344.4
	25-29	43	286.6	22	298.4	21	275.3
	30-34	22	125.6	12	142.6	10	109.9
	35-39	13	84.3	6	79.8	7	88.6
2014	40-44	11	73.1	6	82.0	5	64.7
	45+	12	13.8	5	12.0	7	15.4
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	180	72.8	92	74.7	88	70.9
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	54	314.7	17	184.6	37	465.7
	20-24	104	662.9	54	652.0	50	675.2
	25-29	99	676.1	49	677.9	50	674.3
	30-34	48	276.2	21	253.5	27	296.8

Table 4. P&S Syphilis Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2005	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	1	6.7	1	13.4	0
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	1	6.4	1	12.7	0
	45+	0	0.0	0	0.0	0
	Missing	0	0.0	0	0.0	0
2006	All Ages	2	0.9	2	1.7	0
	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	1	6.5	1	12.9	0
	30-34	1	7.0	0	0.0	1
	35-39	0	0.0	0	0.0	0
	40-44	1	6.6	1	12.9	0
	45+	0	0.0	0	0.0	0
2007	Missing	0	0.0	0	0.0	0
	All Ages	3	1.3	2	1.7	1
	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	0	0.0	0	0.0	0
2008	45+	0	0.0	0	0.0	0
	Missing	0	0.0	0	0.0	0
	All Ages	0	0.0	0	0.0	0
	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	1	6.5	1	12.8	0
2009	40-44	0	0.0	0	0.0	0
	45+	0	0.0	0	0.0	0
	Missing	0	0.0	0	0.0	0
	All Ages	1	0.4	1	0.8	0
	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	2	13.2	0	0.0	2
	40-44	0	0.0	0	0.0	0
	45+	0	0.0	0	0.0	0
	Missing	0	0.0	0	0.0	0
	All Ages	2	0.8	0	0.0	2
						1.7

Continued Table 4. P&S Syphilis

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2010	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	1	6.0	1	11.6	0
	25-29	2	12.3	2	24.5	0
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	0	0.0	0	0.0	0
	45+	3	3.6	3	7.5	0
	Missing	0	0.0	0	0.0	0
All Ages		6	2.5	6	4.9	0
2011	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	5.1	0	0.0	1
	20-24	1	6.0	1	11.6	0
	25-29	2	12.3	2	24.6	0
	30-34	0	0.0	0	0.0	0
	35-39	2	13.7	2	27.4	0
	40-44	0	0.0	0	0.0	0
	45+	3	3.5	3	7.3	0
	Missing	0	0.0	0	0.0	0
All Ages		9	3.7	8	6.5	1
2012	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	1	6.2	1	11.6	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	1	6.6	1	13.5	0
	40-44	1	6.7	1	13.5	0
	45+	3	3.5	3	7.3	0
	Missing	0	0.0	0	0.0	0
All Ages		6	2.4	6	4.9	0
2013	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	5.8	1	10.8	0
	20-24	6	37.6	6	71.3	0
	25-29	0	0.0	0	0.0	0
	30-34	1	5.7	1	11.9	0
	35-39	1	6.5	1	13.3	0
	40-44	2	13.3	2	27.3	0
	45+	3	3.4	2	4.8	1
	Missing	0	0.0	0	0.0	0
All Ages		14	5.7	13	10.6	1
2014	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	5.8	1	10.9	0
	20-24	4	25.5	4	48.3	0
	25-29	4	27.3	4	55.3	0
	30-34	2	11.5	2	24.1	0
	35-39	1	6.3	1	13.1	0
	40-44	0	0.0	0	0.0	0
	45+	3	3.4	3	7.1	0
	Missing	0	0.0	0	0.0	0
All Ages		15	6.0	15	12.1	0

Table 5. Genital Herpes Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2005	0-9	0	0.0	0	0.0	0
	10-14	1	5.0	0	0.0	1
	15-19	12	61.6	2	19.7	10
	20-24	21	125.4	5	57.7	16
	25-29	12	80.8	7	93.5	5
	30-34	18	123.6	6	80.5	12
	35-39	7	46.8	2	26.2	5
	40-44	6	38.4	3	38.0	3
	45+	15	19.7	3	8.3	12
	Missing	1	0.0	0	0.0	1
All Ages		93	40.1	28	24.1	65
2006	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	8	40.5	1	9.7	7
	20-24	14	82.1	2	22.7	12
	25-29	13	84.4	3	38.8	10
	30-34	13	90.7	5	68.2	8
	35-39	5	32.7	2	25.7	3
	40-44	3	19.7	1	12.9	2
	45+	11	14.2	2	5.4	9
	Missing	2	0.0	0	0.0	2
All Ages		69	29.4	16	13.6	53
2007	0-9	0	0.0	0	0.0	0
	10-14	1	5.1	0	0.0	1
	15-19	6	30.0	0	0.0	6
	20-24	16	93.4	7	79.1	9
	25-29	7	44.1	2	25.1	5
	30-34	6	41.8	0	0.0	6
	35-39	8	51.8	2	25.4	6
	40-44	2	13.4	0	0.0	2
	45+	4	5.1	0	0.0	4
	Missing	0	0.0	0	0.0	0
All Ages		50	21.1	11	9.3	39
2008	0-9	0	0.0	0	0.0	0
	10-14	1	5.1	0	0.0	1
	15-19	14	69.5	0	0.0	14
	20-24	20	117.0	4	45.4	16
	25-29	10	61.3	2	24.4	8
	30-34	10	68.5	3	40.4	7
	35-39	3	19.5	2	25.5	1
	40-44	3	20.5	2	26.9	1
	45+	2	2.5	1	2.6	1
	Missing	1	0.0	0	0.0	1
All Ages		65	27.1	14	11.7	50
2009	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	12	59.7	2	19.0	10
	20-24	14	82.2	3	34.2	11
	25-29	9	54.4	7	84.3	2
	30-34	7	46.8	3	39.4	4
	35-39	2	13.2	1	13.0	1
	40-44	7	48.3	1	13.6	6
	45+	5	6.1	1	2.5	4
	Missing	1	0.0	0	0.0	1
All Ages		57	23.6	18	14.9	39

Continued Table 5. Genital Herpes

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2010	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	8	40.1	2	19.1	6	63.1
	20-24	12	72.6	3	34.9	9	113.4
	25-29	10	61.3	2	24.5	8	98.0
	30-34	11	70.5	3	38.2	8	103.3
	35-39	1	6.7	0	0.0	1	13.4
	40-44	2	13.8	0	0.0	2	28.0
	45+	6	7.2	1	2.5	5	11.5
	Missing	0	0.0	0	0.0	0	0.0
2011	All Ages	50	20.6	11	9.0	39	32.1
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	6	30.5	1	9.7	5	53.3
	20-24	19	114.8	4	46.5	15	188.6
	25-29	14	86.1	5	61.4	9	110.9
	30-34	13	81.6	4	49.8	9	113.7
	35-39	6	41.2	4	54.7	2	27.5
	40-44	6	40.7	2	26.7	4	55.1
	45+	9	10.6	1	2.4	8	18.1
2012	Missing	1	0.0	0	0.0	1	0.0
	All Ages	74	30.2	21	17.2	53	43.3
	0-9	1	2.3	0	0.0	1	4.7
	10-14	0	0.0	0	0.0	0	0.0
	15-19	5	27.9	1	10.6	4	47.3
	20-24	21	129.4	5	58.2	16	209.4
	25-29	12	77.3	4	52.4	8	101.5
	30-34	10	58.5	3	36.1	7	79.7
	35-39	2	13.3	1	13.5	1	13.1
	40-44	2	13.4	2	27.0	0	0.0
2013	45+	7	8.2	4	9.7	3	6.7
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	60	24.4	20	16.3	40	32.5
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	5	28.9	0	0.0	5	62.2
	20-24	14	87.7	2	23.8	12	159.0
	25-29	9	60.0	4	54.2	5	65.5
	30-34	10	57.1	3	35.6	7	76.9
	35-39	6	38.9	3	39.9	3	38.0
2014	40-44	6	39.9	4	54.7	2	25.9
	45+	6	6.9	1	2.4	5	11.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	56	22.6	17	13.8	39	31.4
	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	5.0	0	0.0	1	10.3
	15-19	6	35.0	0	0.0	6	75.5
	20-24	17	108.4	4	48.3	13	175.6
	25-29	9	61.5	5	69.2	4	53.9
	30-34	9	51.8	5	60.4	4	44.0