

Specimen Collection and Submission Instructions Carbapenem-resistant Enterobacteriaceae (CRE) Colonization Surveillance (Version 2) Page 1 of 2

Specimen Type	Collection Time	Collection Frequency	Collection Procedures	Transport Media	Shipping & Handling (S&H)*
Rectal Swab	NA	NA	Follow instructions included with the collection kit. Patient swabs should be mailed immediately following collection. Avoid shipping Fridays. Copan dual swabs are viable for testing <5 days after patient collection. Stool specimens may be considered on a case by case basis; Contact HAI coordinator for additional information.	Copan Liquid Stuart Transystem	 Transport device: Provided in collection kit. Rejection Criteria: Using media not provided with collection kit. No approval from LHJ and/or DOH Communicable Disease Epidemiology (CDE). Broken transport device. Unacceptable amount of sample on swabs. Specimens received at testing laboratory collected >5 days prior. Requisition form/isolate discrepancies. Transport: Ship at ambient temperature. Ship as Category B. Storage: Store at ambient temperature.



Specimen Collection and Submission Instructions Carbapenem-resistant Enterobacteriaceae (CRE) Colonization Surveillance (Version 2) Page 2 of 2

General Rejection Criteria (for additional details, see S&H)

• Failure to follow specific S&H requirements.

NOTE:

Specimens will not be processed until ALL of the following information are provided on requisition form:

- Patient name, patient's county of residence, and second identifier.
- Two matching identifiers on both the tube and the laboratory requisition form.
- Specimen type, date of collection, and test requested.
- Submitter name, address, and telephone/FAX numbers.
- Reason for testing at WAPHL.

Collection kit availability

Available. Please contact the Local Health Jurisdiction (LHJ) of the patient's county of residence.

Comments

Pre-approval required from Local Health Jurisdiction (LHJ) and/or DOH Communicable Disease Epidemiology (CDE) before submitting.