# **BIRTH DEFECTS**

# CASE DEFINITION FOR SURVEILLANCE

### Case definition

Case definition for Washington State Birth Defects Surveillance System is based on ICD-9-CM diagnostic and procedure codes as they appear in the hospital medical records. Any child up to age one year, diagnosed or treated, with a reportable birth defect who was a Washington State resident at the time of birth, or treated in a Washington facility is reportable. Information for all stillbirths over 20 weeks gestation diagnosed with a reportable birth defect should also be reported. Currently, we receive a completed hard copy form including the following data elements: Child's name, medical record number, date of birth, sex, admission date, zip code, discharge date, ICD-9-CM code for diagnosis, diagnosis, ICD code for procedure, and procedure.

Table. List of Currently Reportable Birth Defects and the Corresponding ICD-9-CM and ICD-10-CM Codes

Birth Defect	ICD-9-CM	ICD-10-CM	<b>Procedure Code</b>
			(used with ICD9)
1. Anencephaly and similar anomalies	740.0-740.2	Q00	
Acrania	740.0	Q00.0	
Amyelencephalus	740.0	Q00.0	
Hemianencephaly	740.0	Q00.0	
Hemicephaly	740.0	Q00.0	
Craniorachischisis	740.1	Q00.1	
Iniencephaly	740.2	Q00.2	
2. Spina Bifida	741.0, 741.9	<i>Q05, Q07</i>	
• •			
With Hydrocephalus	741.0	Q05.4	
Arnold-Chiari syndrome,	741.0	Q07.0	
type II			
Without mention of		Must select region of	
Hydrocephalus	741.9	defect	
Hydromeningocele	741.9	Q05.9	
Hydromyelocele	741.9	Q05.9	
Meningocele (spinal)	741.9	Q05.9	
3. Cleft Palate	749.0	Q35	
Cleft Palate, Unspecified	749.00	No longer valid; must	
		select bilateral,	
		unilateral or medial	
Cleft Palate, Unilateral,	749.01	Q35.5	
Complete			
Cleft Palate, Unilateral,	749.02	Hard = Q35.1	
Incomplete		<i>Soft</i> = Q35.3	
Cleft Uvula	749.02	Q35.7	

Birth Defect	ICD-9-CM	ICD-10-CM	Procedure Code
			(used with ICD9)
Cleft Palate, Bilateral, Complete	749.03	Q35.4	
Cleft Palate, Bilateral,	749.04	Hard = Q35.0	
Incomplete		Soft = Q35.2	
4. Cleft Lip	749.1	Q36	
Cheiloschisis	749.1	Q36.9	
Congenital fissure of lip	749.1		
Harelip	749.1		
Labium Leporinum	749.1		
Cleft Lip, Unspecified	749.10	Q36.9	
Cleft Lip, Unilateral, Complete	749.11	Q36.9	
Cleft Lip, Unilateral, Incomplete	749.12	Q36.9	
Cleft Lip, Bilateral,	749.13	Q36.0	
Complete Cleft Lip, Bilateral, Incomplete	749.14	Q36.0	
5 Cleft palate with cleft lin	749 2	037	
<i>Cheilonalatoschisis</i>	749.2	Term no longer used	
<i>Cleft palate with cleft lip</i>	749.20	No longer valid must	
unspecified		select bilateral or	
		unilateral	
Unilateral, complete	749.21	Q37.5	
Unilateral, incomplete	749.22	Hard = Q37.1	
		<i>Soft</i> = Q37.3	
Bilateral, complete	749.23	Q37.4	
Bilateral, incomplete	749.24	Hard = Q37.0	
		Soft = Q37.2	
6. Abnormalities of Abdominal Wall	756.70	<i>Q</i> 79.2, <i>Q</i> 79.3	
Omphalocele	756.79	Q79.2	53.41
Gastroschisis	756.79	Q79.3	54.71
7. Limb reduction defects	755.20-755.4	<i>Q71-Q73</i>	
Unspecified reduction deformity of	755.20	Q71.9	
Transverse deficiency of upper limb	755.21	Q71.0 - Q71.3	
Longitudinal deficiency of upper limb	755.22	Q71.4 – Q71.5	
Longitudinal deficiency	755.23	Must select specific	
combined.		bone(s)	
Involving Humerus, Radius,		Q71.4, Q71.5, Q71.6	
and Ulna			
(Complete or incomplete)			

Birth Defect	ICD-9-CM	ICD-10-CM	<b>Procedure Code</b> (used with ICD9)
Longitudinal deficiency. Humeral, complete or Partial(with or without distal deficiencies Incomplete)	755.24	Q71.8	(used with ICD3)
Longitudinal deficiency, Radioulnar, complete or partial (with or without distal deficiencies, incomplete)	755.25	Must select specific bone(s) Q71.4, Q71.5, Q71.6	
Longitudinal deficiency, Radial, Complete or partial (with or without distal deficiencies, Incomplete)	755.26	Q71.4	
Longitudinal deficiency ulnar, complete or partial (with or without distal deficiencies, incomplete)	755.27	Q71.5	
Longitudinal deficiency Carpals or Metacarpals, complete or partial (with or without incomplete phalangeal deficiencies)	755.28	Q71.8	
Longitudinal deficiency, Phalanges Complete or Partial	755.29	Q71.8	
Unspecified reduction deformity of Lower Limbs	755.30	Q72	
Transverse deficiency of Lower Limbs	755.31	Q72.0 – Q72.3	
Longitudinal deficiency of Lower Limb	755.32	Q72.4 – Q72.6	
Longitudinal deficiency, combined, involving Femur, Tibia, and Fibula (complete or incomplete)	755.33	Q72.8	
Longitudinal deficiency Femoral, complete or partial (with or without distal deficiencies, incomplete)	755.34	Q72.4	
Longitudinal deficiency, Tibiofubular complete or partial (with or without distal deficiencies, incomplete)	755.35	Must select specific bone(s) Q72.5, Q72.6	

Birth Defect	ICD-9-CM	ICD-10-CM	<b>Procedure Code</b>
			(used with ICD9)
Longitudinal deficiency Tibia, complete or partial(with or without distal deficiencies, incomplete)	755.36	Q72.5	
Longitudinal deficiency, Fibular complete or partial, (with or without distal deficiencies, incomplete)	755.37	Q72.6	
Longitudinal deficiency, Tarsals or Metatarsals,complete or Partial (with or without incomplete phalangeal deficiency)	755.38	Q72.8	
Longitudinal deficiency, Phalanges, complete or partial	755.39	Q72.8	
<b>Reduction deformities unspecified limb</b>	755.4	Q73.0	
Ectromelia	755.4	Q73.0 Q73.8	
Hemimelia	755.4	Q73.8	
Phocomelia	755.4	Q73.1	
Other		Q73.8	
8. Hypospadias and Epispadias	752.6	<i>Q54, Q64</i>	50.45
Hypospadias	752.61	Report specific degree/location when available Q54.0 -Q54.3 <u>or</u> Other Q54.8 <u>or</u> Unspecified Q54.9	58.45
Epispadias	752.62	Q64.0	58.45
9. Down Syndrome	758.0	<b>Q90.9</b> for NOS Use specific condition third digits when possible(Q90.0 – Q90.2)	
10. Cerebral Palsy	343	<b>G80.9</b> for NOS Use specific condition third digits when possible(G80.0 – G80.8)	
11. Fetal Alcohol Syndrome/Alcohol Related Birth Defects*	760.71	Q86.0, P04.3	
Fetal Alcohol Syndrome		Q86.0	
<i>Fetus/Newborn affected by maternal alcohol</i>		P04.3	

Birth Defect	ICD-9-CM	ICD-10-CM	<b>Procedure Code</b>
			(used with ICD9)
12. Autism Spectrum Disorder**	299.0	F84	
Autism	299.0	F84.0 for Childhood or	
		F84.1 for Atypical	
Asperger's		F84.5	
Pervasive Developmental Disorder		<i>Other</i> F84.8 <u>or</u>	
		Unspecified F84.9	

# A. Description

# 1. General

Birth defects are inborn syndromes, diseases, disorders and malformations that occur before birth. They can affect the organs, senses, limbs, physical and mental development. They also cause pregnancy loss through miscarriage and stillbirth. Some conditions are recognized at birth, others become apparent later in life.

# 2. Cause(s)

The cause(s) of about 70% of all birth defects are unknown. Environmental pollutants may cause birth defects, developmental disabilities, or other adverse reproductive outcomes. Similarly, occupational hazards, dietary factors, medications, infections, and personal behaviors may cause or contribute to birth defects. The causes of birth defects may be a defect in any part of the genome, an interaction between genes or between genes and the environment.

#### 3. World Wide Occurrence

Congenital abnormalities may occur any where in the world. It is possible to see random or systematic geographical variation in the occurrence of some forms of birth defects. This could be due to certain genetic factor(s) or due to a specific exposure(s).

#### 4. High risk factors

Factors that can cause an adverse outcome of pregnancy exist in the environment, where some are known and most of them are not. In addition to what genetic factors may contribute, exposure before or during pregnancy to factors such as smoking, alcohol, medications, illicit drugs, or dietary factors may predispose a pregnant woman to have a child with one or more abnormalities.

# B. Methods of Control

# 1. Prevention Activities

Education of health professionals and the public regarding factors that may be related to birth defects is crucial for prevention. The use of folic acid, a B-vitamin (B9) found in fortified foods and vitamin pills has been shown to be effective in reducing the incidence of neural tube defects. The U.S. Public Health Service recommends that all women who could possibly become pregnant get 400 micrograms (or 0.4mg) of folic acid every day. This could prevent up to 70% of some types of serious birth defects. But to do this, women need folic acid before they get pregnant. Since over 50% of births in Washington are unintended, it is important to promote daily consumption of folic acid to all women of childbearing age. In addition, early identification of birth defects promotes care coordination and secondary prevention activities to improve the quality of life.

# 2. Occurrence of Clusters

The occurrence of a new or an already known birth defect in a cluster is an indicator of exposure to a known or an unknown factor(s) or may be by chance. Suspected clusters (3) or more cases of the same or developmentally similar conditions) may be reported to Jeanette Robbins, at (360) 236-3591 or to Juliet VanEenwyk, the state epidemiologist for non-communicable conditions at (360) 236-4250.

# 3. International Measures

Once a teratogen or a risk factor proves to be causal, that knowledge should be communicated internationally and appropriate measure should be taken to avoid exposures. Efforts to prevent exposures of pregnant women to Thalidomide, Aminopterin, Isotretinoin all over the world are examples of this kind of international effort.

# 4. Control Measures

Environmental protection, surveillance and research

# 5. Public Health Importance

Birth defects are the leading cause of infant mortality in the United States, accounting for more than 20% of all infant deaths. Of about 120,000 U.S. babies born each year with a birth defect, 8,000 die during their first year of life. Birth defects are the 5<sup>th</sup> leading cause of years of potential life lost and contribute substantially to childhood morbidity and long-term disability. The health care cost associated with birth defects is also immense.



DOH 420-454 October 2006

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email doh.information@doh.wa.gov.