

12 Months

Well-Child Visits

Your child is 1!

Your child's next well-child visit is due just after their first birthday. During this visit, ask your child's doctor or nurse if you have questions about your child's growth and development. You can also check with your child's doctor or nurse to make sure your child's vaccine records are in the Washington State Immunization Information System (WAIIS).

Oral Health

Time for your child's first dental checkup.

Your child should see the dentist (or a doctor trained to check children's teeth) by age 1 even if their teeth have not grown in yet. This visit is usually a short and simple exam with your child on your lap. Your dentist will suggest ways to prevent tooth decay. To find a dentist, call the Help Me Grow Washington Hotline at 1-800-322-2588 or call 211.

Vaccines

Vaccines between 12 and 18 months.

Childhood vaccines can protect your child from 17 different diseases. Vaccines strengthen the immune system by preparing it to defend against viruses and bacteria that cause serious disease.

More than 1 dose of a vaccine is often needed for the best protection against specific diseases. Talk with your child's doctor or nurse about which of the following vaccines your child needs to be up to date:

- Hepatitis B
- Diphtheria, tetanus, pertussis (DTaP)
- Haemophilus influenzae type b (Hib)
- Pneumococcal conjugate vaccine (PCV)
 (PCV)
- Inactivated polio vaccine (IPV)
- Flu, every year

Growth and Development

No screen time (TV, video, or computer) until 18 months.

Babies and young children do not need any screen time. The American Academy of Pediatrics recommends no screen time, other than video-chatting, until at least 18 months. Instead, talk, sing, read, and play with your child to help their brain develop best. Make a screen time plan for your family.

Consider each child's age, health, personality, and developmental stage. Share your plan with your children's other caregivers to be consistent. Find screen time guidelines at **bit.ly/stguide2**.

Very young children can feel stress.

Starting child care or changing child care providers can be hard for your child. They may show signs like needing more attention or not sleeping or eating well. It is normal for children to miss people they have spent time with.

Be patient. If you see a change in their behavior, try to think about what routines may have changed. Comfort your child and talk about it. Even though your child may not be able to talk back to you, your child will know they can depend on you when things are difficult or scary. For more help, call the Family Help Line at 1-800-932-4673.



Comfort me before and after I get vaccinated

- Measles, mumps, and rubella (MMR)
- Chickenpox (varicella)

and Physical

Your child needs nutritious foods

To help your child grow strong and have

plenty of energy, it is important for them

Good examples of these foods are fruits,

eggs, beans, and tofu. While your child's

stomach is small, they still need 3 meals

If you're feeding your child human milk,

it has the fat, protein, and most vitamins

and minerals they need for growth and

brain development. Follow your child's

need. Talk to your doctor about a vitamin D supplement and adding other foods to

your child's diet that are rich in vitamin D.

cues and nurse them as often as they

to eat a variety of foods rich in fiber,

protein, vitamins and minerals. Your

child also needs iron and calcium.

vegetables, grains, chopped meats,

along with 2-3 snacks daily.

Milk for 1 year olds:

- Hepatitis A
- COVID-19

Nutrition

Activity

every day.

If your child is formula-fed, it is recommended to switch them to whole cow's milk when they turn 1 year of age. Whole cow's milk is a better source of fat than lower-fat milk or non-dairy milks for your growing child. Vitamins, including vitamin D, are added to cow's milk to meet your child's needs.

Give your child a total of 4 to 5 servings of whole milk a day. A serving of milk is 4 ounces. Giving more milk than that can make your child too full to eat other foods. Wait until your child is at least 2 years old to give them non-fat or low-fat milk.

If your child has trouble digesting cow's milk, talk to your doctor or WIC staff (Women, Infants, and Children Nutrition Program) about other foods to give them that have enough calcium, such as lactose-free milk or calcium- and vitamin D-fortified milk alternatives.





Trust your insticts. You know your baby best!

To comfort your child during and after they get vaccines, try these tips:

- Stay calm. Your child can sense when you are anxious. If you don't think you can stay calm, find someone else who can be with your child while they get vaccinated.
- Bring your child's favorite toy or blanket.
- Hold your child on your lap. Talk or sing to them.
- Breast, chest, or bottle feed your child while they get vaccinated.
- Put a cool, clean, wet washcloth over the area where the vaccine was given.

Ask your doctor about medicine for pain or fever. Find more after vaccination care tips at **bit.ly/vax-after-care-tips**.

Safety

Encourage your child to explore safely.

Never leave your child in a room alone. Always make sure there are safe places to play when you can't give them your full attention.

If the room you are in is not child proofed, put them in a play seat (without wheels) or a playpen for a short period of time.

Make sure your home and other caregivers' homes are safe for your child to explore. Lock up things like:

- Household chemicals
- E-cigarette and vaping products
- Marijuana products
- Medications
- Sharp objects
- Firearms
- Alcohol
- Illegal substances

Do not expect your child not to play or touch these things just because you say "No!" Putting them away is easier and safer.

More ways to keep your home safe for your child.

Here are other things you can do to keep your home safe for your child to explore:

- Keep furniture away from windows so your child cannot climb up and fall out.
- Window screens are not strong enough to hold your child inside. Consider adding window stops for upper-level windows and set them to stop at 4 inches or keep these windows closed and locked.

• Bolt or strap furniture, like dressers, flat panel TVs, or TV cabinets, to the wall from the top. Dressers as short as 24 inches have killed children.

Choosing safe toys.

When you buy a toy, look at what age child it is recommended for. Your child is not old enough for toys with small pieces or sharp edges.

Be especially careful with toys containing button batteries, magnets or water beads, which can be dangerous for your child if swallowed.

Your child may enjoy simpler toys, such as plastic cups in the bath or a large cardboard box.

Keep your child in a car seat.

Keep your child in the backseat of your car and buckled into a car seat. A car seat that is used correctly can save your child's life and is required by law.

Your child must sit rear-facing until at least age 2 or until they exceed the weight or height maximum for the car seat. This protects your child's head, neck, and spine from serious injury and it is the law in Washington.

During cold months, take off your child's heavy clothes before putting them in the car seat. Bulky jackets or coats make it hard to get the car seat straps fitting snugly. First secure the harness on your child then cover them with a coat or blanket to keep them warm.

As your child gets more active, they may not want to stay in the car seat. Have soft toys or talk and sing to them in the car. Never leave babies or children in the car without an adult, even for a short time. It is very dangerous.

For more information to keep your family safe in the car or about Washington's child passenger safety laws visit **wacarseats.com**.



Keep me sitting rear-facing as long as possible.

Prevent burns (scalds).

Your child can get burned by hot water, drinks, and food. Watch them closely in the kitchen. Here are some tips to prevent burns:

- Keep pots, bowls, and cups away from the edge of the table or counter.
- Use the back burners and always turn pot handles toward the back of the stove.
- Avoid using tablecloths. Children can pull hot things off the table and onto themselves.
- Do not hold hot drinks when carrying your child, when your child is on your lap, or when they are in the stroller.
- Use travel mugs with lids when drinking hot drinks at home. Children can pull hot things off coffee tables and end tables.

Family Support and Routines

Pregnancy planning.

Waiting at least 18 months between pregnancies is best for you and your baby. This allows your body to heal and reduces the risk for premature birth.

A full-term pregnancy is best for a baby's brain, lungs, eyes, ears, and other organs. To find a family planning provider, call the Help Me Grow Washington Hotline at 1-800-322-2588 or visit **helpmegrowwa.org**

To learn more on how long to wait between pregnancies, visit **bit.ly/MD-IDI**.

Take care of your body.

Eat meals and snacks that include a variety of fruits, vegetables, protein, and whole grains. Choose foods rich in calcium, vitamin D, iron, and folic acid.

Check with your doctor to ask if you should continue taking prenatal vitamins or switch to a multivitamin. All people who are planning to get pregnant are encouraged to get at least 400 micrograms of folic acid daily to help reduce risk of birth defects.

Environmental Health

Lock up things that can poison your child.

Older babies and toddlers are curious. They learn to open containers (even ones that are child resistant) and may eat things that will harm them, like:

- all chemicals including cleaning supplies and pest products
- medications and supplements,
- alcohol,
- firearms,

- tobacco and tobacco products,cannabis and cannabis products,
- illegal substances,
- and other small items.
- Keep these items on a high shelf in a closet or cupboard with a door that locks. If you think your child has eaten or drank something poisonous, stay calm and call the Washington Poison Center at 1-800-222-1222. If your child is unconscious, convulsing, or having trouble breathing, call 911. For information on poisons around the home and environmental toxins, visit **wapc.org**.



Lock up anything that can poison me

For the Health of All Our Children



To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers needing assistance, please call 711 (Washington Relay) or email **doh.information@doh.wa.gov**. DOH 920-929E March 2025

fin a clocot or cupboard with a door that l

Thanks for reading!

Share this letter with other caregivers in your child's life. You'll hear from us again before your child is 15 months old with information on:

- Hand washing.
- Helping your child learn about emotions.
- How dangerous batteries and magnets can be to your child.