


# Multisystem Inflammatory Syndrome in Children Associated with COVID-19 in Washington State

November 9, 2022





To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email [civil.rights@doh.wa.gov](mailto:civil.rights@doh.wa.gov).

**Publication Number**

420-290

For more information or additional copies of this report:

Disease Control and Health Statistics

Communicable Disease Epidemiology

1610 NE 150th Street, MS: K17-9

Shoreline, WA 98155Address 2

206-418-5500 (24-hour contact for local health jurisdictions only)

[CommDisEpi@doh.wa.gov](mailto:CommDisEpi@doh.wa.gov)



# Overview

This report contains information on cases reported through October 31, 2022.

Multisystem inflammatory syndrome in children (MIS-C) is a condition that causes inflammation in different body parts, including the heart, lungs, kidneys, brain, skin, eyes, or gastrointestinal organs. Children with MIS-C may have a fever and symptoms such as abdominal (gut) pain, vomiting, diarrhea, neck pain, rash, bloodshot eyes, or feeling extra tired. We do not yet know what causes MIS-C, but it is a syndrome associated with COVID-19. Children with MIS-C had the virus that causes COVID-19 or had been around someone with COVID-19.

The case definition of MIS-C is:

- Under the age of 21, with a fever, laboratory evidence of inflammation, and severe illness involving more than two organs that requires hospitalization; AND
- No other plausible diagnoses; AND
- Positive COVID-19 test (PCR, antigen, or serology) or exposure to a confirmed case within the four weeks prior to the onset of symptoms.

See the US Centers for Disease Control website for [national MIS-C case reporting](#).

Healthcare providers should report patients meeting MIS-C criteria to their [local public health agency](#).

In August 2022, Washington State Department of Health enhanced MIS-C surveillance by looking for potential cases in the Rapid Health Information Network (RHINO) database of healthcare visits. Previously, only direct reports from healthcare providers and facilities were available to us to help identify cases. Going forward, we will use both methods to identify potential MIS-C cases.


We monitor the vaccine status of all MIS-C cases. No MIS-C cases reported in Washington in 2020 or 2021 had received COVID vaccine. In 2022, five cases of MIS-C have been identified in children who had received COVID vaccine prior to the onset of their COVID-related illness. No deaths have occurred among children who had MIS-C.

Please note that the data presented in this report may change as we get additional information on cases.

# Cases identified in Washington

A total of 165 confirmed cases of MIS-C have been identified in Washington as of October 31, 2022.

The chart below displays all MIS-C cases identified in Washington by month of illness onset.



The table below shows MIS-C cases in Washington by county of residence.

**Table 1 MIS-C cases in WA by county**

<b>County in Washington</b>	<b>2020</b>	<b>2021</b>	<b>2022 YTD</b>	<b>Total</b>
<b>Benton</b>	0	2	1	3
<b>Chelan</b>	1	1	1	3
<b>Clark</b>	0	2	0	2
<b>Cowlitz</b>	0	1	0	1
<b>Douglas</b>	1	1	0	2
<b>Franklin</b>	2	5	0	7
<b>Grant</b>	0	6	1	7
<b>Island</b>	0	2	1	3
<b>King</b>	10	20	12	42
<b>Kitsap</b>	2	2	1	5
<b>Kittitas</b>	0	1	0	1
<b>Lewis</b>	2	0	0	2
<b>Lincoln</b>	0	0	1	1
<b>Mason</b>	1	0	0	1
<b>Pend Oreille</b>	0	0	1	1
<b>Pierce</b>	4	17	6	27
<b>Skagit</b>	2	0	1	3
<b>Snohomish</b>	5	8	5	18
<b>Spokane</b>	1	14	2	17
<b>Stevens</b>	0	1	0	1
<b>Thurston</b>	0	3	2	5
<b>Whatcom</b>	1	2	1	4
<b>Yakima</b>	3	5	1	9
<b>Total</b>	<b>35</b>	<b>93</b>	<b>37</b>	<b>165</b>

The table below shows MIS-C cases in Washington by patient ethnicity.

**Table 2 MIS-C cases in WA by patient ethnicity**

Patient ethnicity	Number of cases of MIS-C (percent)			
	2020	2021	2022 YTD	Total
<b>Hispanic</b>	15 (42.9%)	24 (25.8%)	4 (10.8%)	43 (26.1%)
<b>Non-Hispanic</b>	18 (51.4%)	62 (66.7%)	31 (83.8%)	111 (67.3%)
<b>Unknown</b>	2 (5.7%)	7 (7.5%)	2 (5.4%)	11 (6.7%)
<b>Total</b>	<b>35 (100.0%)</b>	<b>93 (100.0%)</b>	<b>37 (100.0%)</b>	<b>165 (100.0%)</b>

The table below shows MIS-C cases in Washington by patient race.

**Table 3 MIS-C cases in WA by patient race**

Patient race	Number of cases of MIS-C (percent)			
	2020	2021	2022 YTD	Total
<b>American Indian and Alaska Native</b>	1 (2.9%)	0	1 (2.7%)	2 (1.2%)
<b>Asian</b>	2 (5.7%)	4 (4.3%)	3 (8.1%)	9 (5.5%)
<b>Black or African American</b>	3 (8.6%)	15 (16.1%)	3 (8.1%)	21 (12.7%)
<b>Native Hawaiian and Pacific Islander</b>	2 (5.7%)	2 (2.2%)	0	4 (2.4%)
<b>White</b>	18 (51.4%)	36 (38.7%)	21 (56.8%)	75 (45.5%)
<b>Other</b>	2 (5.7%)	12 (12.9%)	2 (5.4%)	16 (9.7%)
<b>Multiracial</b>	0	5 (5.4%)	1 (2.7%)	6 (3.6%)
<b>Unknown</b>	7 (20.0%)	19 (20.4%)	6 (16.2%)	32 (19.4%)
<b>Total</b>	<b>35 (100.0%)</b>	<b>93 (100.0%)</b>	<b>37 (100.0%)</b>	<b>165 (100.0%)</b>



Ages of MIS-C cases in Washington range from 0-19 years (median 8 years, mean 8 years).  
 The table below shows MIS-C cases in Washington by age range.

**Table 4 MIS-C cases in WA by age range**

Patient age range	Number of cases of MIS-C (percent)			
	2020	2021	2022 YTD	Total
<b>0-9 years</b>	19 (54.3%)	60 (64.5%)	27 (73.0%)	106 (64.2%)
<b>10-20 years</b>	16 (45.7%)	33 (35.5%)	10 (27.0%)	59 (35.8%)
<b>Total</b>	<b>35 (100.0%)</b>	<b>93 (100.0%)</b>	<b>37 (100.0%)</b>	<b>165 (100.0%)</b>

The table below shows MIS-C cases in Washington by patient sex at birth.

**Table 5 MIS-C cases in WA by patient sex at birth**

Patient sex at birth	Number of cases of MIS-C (percent)			
	2020	2021	2022 YTD	Total
<b>Female</b>	15 (42.9%)	46 (49.5%)	17 (45.9%)	78 (47.3%)
<b>Male</b>	20 (57.1%)	47 (50.5%)	20 (54.1%)	87 (52.7%)
<b>Total</b>	<b>35 (100.0%)</b>	<b>93 (100.0%)</b>	<b>37 (100.0%)</b>	<b>165 (100.0%)</b>

