

FIRST QUARTER 2022 – March Update

Statewide High-Level Analysis of Forecasted Behavioral Health Impacts from COVID-19

Purpose

This document provides a brief overview of the potential statewide behavioral health impacts from the COVID-19 pandemic. The intent of this document is to communicate potential behavioral health impacts to response planners and organizations or individuals who are responding to or helping to mitigate the behavioral health impacts of the COVID-19 pandemic.

Bottom Line Up Front

- The COVID-19 pandemic has strongly influenced behavioral health symptoms and behaviors across the state due to far-reaching medical, economic, social, and political consequences. This forecast is heavily informed by disaster research and the latest data and findings specific to this pandemic. Updates will be made monthly to reflect changes in baseline data.
- As we transition into the third year of the global pandemic, we shift from an emergency response perspective into a long-term, recovery-management perspective on behavioral health outcomes related to COVID-19. Various outcome pathways are beginning to emerge. Adaptability and resilience will be the most-likely outcome for many, but for others impacted by factors such as age, access to resources, or protective aspects, there may be an increase in acuity of existing symptoms, the development of new psychological conditions (Major Depressive Disorder, Generalized Anxiety Disorder, and PTSD as examples), or an outcome of chronic dysfunction. These varied responses over time, in addition to other behavioral health trajectories and outcomes, are typical of large-scale layered events in the long run.^{1,2}
- There are three behavioral health areas of focus for 2022:

1. **Anxiety and excitement dichotomy regarding the end of mask mandates and vaccine requirements in certain sectors, including many public spaces and schools.**

Anxiety and excitement activate very similar and often overlapping neurological pathways in the brain. It is likely that many people may experience a mixed reaction, or a clear but rapid fluctuation in feelings about these changes to the state's guidelines.



DOH 821-103-23 March 2022
To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov.

Sheer exhaustion with the complexity and duration of this experience may contribute to emotion dysregulation and increases in hostility or aggression. This may be evident in both behaviors and perceptions related to “us versus them” thinking. Families with young children and parents supporting adolescents and youth may be under considerable and disproportionate strain related to their students’ need to navigate previously unknown social opportunities and challenges with peers when it comes to mask guideline changes.

2. **Children, youth, and young adults.** Behavioral health concerns for children, youth, and young adults may increase as seasonal challenges overlap with pandemic related issues. It is also notable that there has been a significant increase in tic and tic-like³ behavior amongst adolescent females in recent months. This is likely in part a reflection of the degree to which this demographic group is currently experiencing distress, as well as the influence of social media, causing some to self-diagnose disorders.
3. **Additional world-wide stressors.** The mental and emotional toll of items, such as the recently released [UN climate report](#),^{4,a} in addition to the war in Ukraine, may have significant impacts on specific groups and communities within the population.

Areas of Focus for First Quarter 2022

General Trends

Long-term outcomes for large-scale disasters typically are characterized by resilience, but there are certainly groups and individuals who experience cascade effects, including increased behavioral health symptoms and substance use, chronic dysfunction, and other problematic long-term effects.^{1,2}

The stressors on an already overwhelmed and exhausted health care workforce have created a breaking point for many.

For health care workers, those in public health services, and educators, the need for the establishment and maintenance of clear boundaries around work and home time to promote resilience and reduce burnout is essential. Burnout is a concern across agencies and work functions for many,^{5,6,7,8} and this may be even more so for women and people of color.⁹ Staffing shortages nationwide have contributed to additional job stress. The politicized nature of the way the pandemic has evolved has also contributed to adverse mental health experiences for many responders.

As the pandemic continues, behavioral choices about compliance and caution may be heavily influenced by personal, primary impacts (illness and hospitalization, or the loss of a loved one from the virus), with more cautious behaviors being associated with more significant and negative experiences of the pandemic or the virus itself. However, the longer the pandemic continues and is conceptualized and managed as an ongoing disaster or threat, the more complacent many may become in following safety guidelines and recommendations for non-pharmaceutical interventions.¹⁰

Anxiety and excitement dichotomy about the ending of mask and vaccine mandates

As we transition away from “pandemic response” towards endemic management of COVID-19, some restrictions will be eased that were previously necessary. There is likely to be a predictable level of

^a https://report.ipcc.ch/ar6wg2/pdf/IPCC_AR6_WGII_FinalDraft_FullReport.pdf

emotional intensity that arises. For many, the easing of some mandates is celebratory and may result in significant excitement, energy, and happiness. For others, the same changes may result in a sense of uncertainty, fear, and anxiety or dread about potential risks or harm.

For some demographic groups, such as students in the K – 12 system, reactions to changes specific to mask mandates are likely to be very intense in either direction. Many students have long awaited the opportunity to go mask-free, while others have appreciated the sense of social and personal security that masks have provided. Some students may fear social reprisal from peers if they choose to continue to wear a mask, and the potential that attention to this may increase-bullying or harassment amongst students. This will need to be a point of vigilance for educators for the remainder of this academic year.

Children, Youth, and Young Adults

Concerning behavioral health trends for children, youth, and young adults will very likely continue and potentially increase into the second quarter of 2022.

Children and Families

The behavioral health crisis that was addressed in the Governor’s [emergency proclamation](#) on March 15, 2021 may be compounded by typical seasonal challenges in behavioral health.^{11,12,13} It is typical for behavioral health symptoms to increase in youth at this time of year, when the longer holidays are behind them and the academic demands tend to increase. Suicide attempts and rates also typically rise in spring,¹⁴ which also coincides with the anniversary of the start of the COVID-19 pandemic, and the recognition that it is not yet “over.” Emergency Department visits for behavioral health concerns related to psychological distress, suicidal ideation, and suicide attempts have been increasing throughout the state in recent weeks. Please see the latest [Youth Situational Report](#)^b for more detail. Also of note is the dramatic nationwide increase in tic and tic-like disorders and symptoms, particularly amongst adolescent females, according to the Northwest Healthcare Response Network (twice weekly hospital huddle, 2022).³ It is likely that these increases are related in some way to the significant degree of distress, social pressure, confusion, and general upheaval that has been a hallmark of the last several years for this demographic group. Caregivers, educators, and parents may benefit from helping youth focus on active anxiety symptom management to help mitigate more problematic behavioral health symptoms from developing. This might include a calm and matter-of-fact approach to symptoms, not encouraging peer or adult attention to tics, and encouraging students exhibiting tics to use the tools provided to calm themselves. These can include psycho-education, relaxation, and behavioral interventions, such as habit reversal training.³ These students may also benefit from having places designated within the schools for them to go briefly to regroup and use their self-calming tools before returning to the classroom.

Classroom Considerations & Academics

As the 2021-2022 academic year progresses, many students and their caregivers may be confronted with concerns that their academic attainment is not where they would want it to be after the experience of the last 18 months. Some may need intensive tutoring to catch up on missed academic skills. Students who experienced the loss of a key transition year (between elementary and middle school or middle and high school) during the pandemic may be experiencing more extreme psychological or developmental disruptions in the current year. Parents, caregivers, and educators may need to modify expectations and focus on helping

^b<https://doh.wa.gov/sites/default/files/2022-03/821-135-YouthBehavioralHealthSitRep-March2022.pdf>

children and youth re-engage by first learning how to be a successful student again by socializing and participating with peers in a classroom context before focusing on academic success. Resources for parents, caregivers, educators, and other school staff can be found in both [the COVID-19 Behavioral Health Toolbox for Families: Supporting Children and Teens During the COVID-19^c Pandemic](#) and the [COVID-19 Back-to-Classroom THINK Toolbox: Teaching with Healthcare Informed Neurological Strategies for Kids^d](#)

Depression and Suicide

Depression is a common response throughout the disaster recovery cycle. Many children, youth, and young adults are experiencing significant symptoms of depression during the pandemic.¹⁵ In Washington, trends indicate that emergency department visits for suicidal ideation, attempts, and psychological distress may be increasing, and these are data that will continue to be monitored as emergency department visits are increasing generally. The most recent reporting from hospitals in Washington that admit pediatric patients indicates that the surge of youth presenting to emergency departments for suicidal ideation and suspected suicide attempts remains an ongoing increasing issue. Lack of outpatient behavioral health services and psychiatric beds has led to increasing numbers of youth who are “boarding” in emergency rooms and med/surge beds, sometimes for extended periods of time, and without treatment while waiting. Youth with additional complexities, such as autism with aggressive behaviors or significant developmental delay, have even more restricted access to appropriate treatment and some have boarded for months awaiting placement. Other youth are boarding for extended periods of time not because of ongoing mental health needs, but because exhausted parents are refusing to bring them home, citing safety concerns, and agencies who might take them into care are unable to find placement for them.

Active suicide prevention should be promoted through sharing information on recognizing [warning signs^e](#) and other related resources, and checking in with colleagues, friends, family members, and neighbors. When someone is expressing thoughts of self-harm, [access to dangerous means of harm should be removed^f](#), and medications, poisons, and firearms should be stored safely. Suicides consistently account for approximately 75% of all firearm-related fatalities in Washington.¹⁶ [Storing firearms safely](#) and [temporarily removing them from the home](#) of an at-risk person during a crisis can save lives.

Additional Resources

- Anyone concerned about depression or other behavioral health symptoms should talk with their **healthcare provider**.
- [Washington Listens^g](#): Call 833-681-0211 to talk to a support specialist who will listen and help you cope with the stress of COVID-19.
- **Health Care Authority: [Mental health crisis lines^h](#)**

^c <https://www.doh.wa.gov/Portals/1/Documents/1600/coronavirus/BHG-COVID19-FamilyToolbox.pdf>

^d <https://www.doh.wa.gov/Portals/1/Documents/1600/coronavirus/821-148-BackToClassroomToolbox.pdf>

^e <https://www.doh.wa.gov/YouandYourFamily/InjuryandViolencePrevention/SuicidePrevention/HelpSomeoneElse#common>

^f <https://www.seattlechildrens.org/health-safety/keeping-kids-healthy/prevention/home-checklist/>

^g <https://www.walistsens.org/>

^h <https://www.hca.wa.gov/health-care-services-supports/behavioral-health-recovery/mental-health-crisis-lines>

- [National Suicide Prevention Lifeline](https://suicidepreventionlifeline.org/):ⁱ Call 800-273-8255 (English) or 1-888-628-9454 (Español).
- [Crisis Connections](https://www.crisisconnections.org/24-hour-crisis-line/):^j Call 866-427-4747.
- [Crisis Text Line](https://www.crisistextline.org/):^k Text HEAL to 741741.
- Department of Health: [Crisis lines for specific groups](https://www.doh.wa.gov/YouandYourFamily/InjuryandViolencePrevention/SuicidePrevention/HotlinesTextandChatResources)^l
- [TeenLink](https://www.crisisconnections.org/teen-link/):^m Call or text 866-833-6546
- A Mindful Stateⁿ: <https://amindfulstate.org/>
- [Washington Warm Line](https://www.crisisconnections.org/wa-warm-line/):^o Call 877-500-9276
- Washington State COVID-19 Response: [Mental and emotional well-being webpage](https://coronavirus.wa.gov/wellbeing)^p

Additional World-Wide Stressors

Current events that affect the global population also add layers of concern to our collective recovery from the COVID-19 pandemic. The recently released [UN climate report](#)⁴ may contribute to a sense of foreboding, additional grief, or even acute distress for individuals, communities, and regions who have been more directly affected by climate related disasters (fires, floods, etc.) over the past two years. In addition to climate-related anxiety and distress, the current war in Ukraine may also be a significant source of concern for many in Washington. Many Washingtonians have close family or professional ties to Ukraine and Russia, and may be experiencing additional trauma-related concerns for the safety of loved ones, or the war-related images and stories that are flooding the media.

Focusing on control over the *process* we use to engage with the world around us is one active coping technique that may be helpful during this time. Looking for ways to **express compassion for others, lend a helping hand in our own communities, and respond with empathy** when possible are all examples of how we can engage with personal empowerment right now. It may also be beneficial to establish and maintain healthy boundaries around personal exposure to media (including social media) and news generally.

Other Considerations

Behavioral health symptoms will continue to present in phases.^{17,18} The unique characteristics of this pandemic trend towards anxiety and depression as a significant behavioral health outcome for many in Washington. These outcomes have been shown throughout the Behavioral Health Impact Situation Reports published by DOH, which are available on the [Behavioral Health Resources & Recommendations webpage](#)^q under the “Situation Reports” dropdown. Behavioral health symptoms of anxiety, impulsivity, reduced frustration tolerance, anger, depression, and post-traumatic stress disorder (PTSD) are likely to increase with any significant increases in infection and hospitalization rates.^{19,20}

ⁱ <https://suicidepreventionlifeline.org/>

^j <https://www.crisisconnections.org/24-hour-crisis-line/>

^k <https://www.crisistextline.org/>

^l <https://www.doh.wa.gov/YouandYourFamily/InjuryandViolencePrevention/SuicidePrevention/HotlinesTextandChatResources>

^m <https://www.crisisconnections.org/teen-link/>

ⁿ <https://amindfulstate.org/>

^o <https://www.crisisconnections.org/wa-warm-line/>

^p coronavirus.wa.gov/wellbeing

^q <https://www.doh.wa.gov/Emergencies/COVID19/HealthcareProviders/BehavioralHealthResources>

Reactions and Behavioral Health Symptoms in Disasters – COVID-19




Figure 1: Phases of reactions and behavioral health symptoms in disasters. The dotted graph line represents the response and recovery pattern that may occur if the full force of a disaster cascade is experienced by a majority of the population (i.e., the disaster cascade pathway). Protective factors are characteristics, conditions, or behaviors that reduce the effects of stressful life events. They also increase a person’s ability to avoid risks or hazards, recover, and grow stronger. Adapted from the Substance Abuse and Mental Health Services Administration (SAMHSA).²¹

Acknowledgements

This document was developed by the Washington State Department of Health’s Behavioral Health Strike Team for the COVID-19 response. The strike team is a group of clinical psychologists, psychiatrists, and therapists who are professionals in disaster relief and behavioral health. Lead authors from the Behavioral Health Strike Team are Kira Mauseth, Ph.D., Tona McGuire, Ph.D., and Stacy Cecchet, Ph.D., ABPP. Research support for this report was provided by undergraduate psychology students at Seattle University.

References

1. Fran H. Norris, F.H., Tracy, M., & Galea, S. (2009). Looking for resilience: Understanding the longitudinal trajectories of responses to stress, *Social Science & Medicine*, Volume 68, Issue 12, Pgs 2190-2198, ISSN 0277-9536, <https://doi.org/10.1016/j.socscimed.2009.03.043>.
2. Adams, V., VAN Hattum, T., & English, D. (2009). Chronic disaster syndrome: Displacement, disaster capitalism, and the eviction of the poor from New Orleans. *American ethnologist*, 36(4), 615–636. <https://doi.org/10.1111/j.1548-1425.2009.01199.x>
3. Ganos, C., Martino, D., & Pringsheim, T. (2017). Tics in the Pediatric Population: Pragmatic Management. *Movement disorders clinical practice*, 4(2), 160–172. <https://doi.org/10.1002/mdc3.12428>
4. H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.). IPCC, 2022: *Climate Change 2022: Impacts, Adaptation, and Vulnerability*. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press. https://report.ipcc.ch/ar6wg2/pdf/IPCC_AR6_WGII_FinalDraft_FullReport.pdf.
5. Wiesman, J., & Baker, E. L. (2022). The Public Health Worker Mental Health Crisis—A Major Leadership Challenge. *Journal of Public Health Management and Practice*, 28(1), 95-98.
6. Sarah E. Scales, Elizabeth Patrick, Kahler W. Stone, Kristina W. Kintziger, Meredith A. Jagger, and Jennifer A. Horney. Health Security. ahead of <http://doi.org/10.1089/hs.2021.0132>
7. Chen, R., Sun, C., Chen, J.-J., Jen, H.-J., Kang, X.L., Kao, C.-C. and Chou, K.-R. (2021), A Large-Scale Survey on Trauma, Burnout, and Posttraumatic Growth among Nurses during the COVID-19 Pandemic. *Int. J. Mental Health Nurs.*, 30: 102-116. <https://doi.org/10.1111/inm.12796>
8. Pressley, T. (2021). *Factors Contributing to Teacher Burnout During COVID-19*. *Educational Researcher*, 50(5), 325–327. <https://doi.org/10.3102/0013189X211004138>
9. Prasad .K, Colleen McLoughlin, Martin Stillman, Sara Poplau, Elizabeth Goelz, Sam Taylor, Nancy Nankivil, Roger Brown, Mark Linzer, Kyra Cappelucci, Michael Barbouche, Christine A. Sinsky. *Prevalence and correlates of stress and burnout among U.S. healthcare workers during the COVID-19 pandemic: A national cross-sectional survey study*. *EClinicalMedicine*, Volume 35, 2021, 100879, ISSN 2589-5370, <https://doi.org/10.1016/j.eclinm.2021.100879>. (<https://www.sciencedirect.com/science/article/pii/S2589537021001590>)
10. Chen, F., Griffith, A., Cottrell, A., & Wong, Y. L. (2013). *Behavioral responses to epidemics in an online experiment: using virtual diseases to study human behavior*. *PloS one*, 8(1), e52814. <https://doi.org/10.1371/journal.pone.0052814>
11. Eastwood MR, Peacocke J. *Seasonal patterns of suicide, depression, and electroconvulsive therapy*. *Br J Psychiatry*. 1976 Nov;129:472-5. <https://pubmed.ncbi.nlm.nih.gov/990662/>
12. Ayers, J. W., Althouse, B. M., Allem, J.-p., Rosenquist, J. N., & Ford, D. E. (2-13). Seasonality in Seeking Mental Health Information on Google. *American Journal of Preventive Medicine*, 44(5) 520-525. <https://pubmed.ncbi.nlm.nih.gov/23597817/>
13. Sullivan, B., & Payne, T. W. (2007). Affective Disorders and Cognitive Failures: A Comparison of Seasonal and Nonseasonal Depression. *American Journal of Psychiatry*, 164(11), 1663-1667. <https://pubmed.ncbi.nlm.nih.gov/17974930/>
14. Woo, M., Okusaga, O., Postoache, T.T. 2012 Seasonality of Suicidal Behavior. *Int J Environ Res Public Health* Feb; 9(2): 531-547. <https://doi.org/10.3390/ijerph9020531>
15. Czeisler, M. É., Lane, R. I., Petrosky, E., et al. (2020). Mental Health, Substance Use, and Suicidal Ideation During the COVID-19 Pandemic — United States, June 24–30, 2020. *MMWR Morb Mortal Wkly Rep*, 69, 1049–1057. <http://dx.doi.org/10.15585/mmwr.mm6932a1>
16. Washington State Department of Health. (2019). *Annual Report: Firearm Fatality and Suicide Prevention – A Public Health Approach*. <https://www.doh.wa.gov/Portals/1/Documents/8390/346-087-SuicideFirearmPrevention.pdf>
17. Substance Abuse and Mental Health Services Administration (SAMHSA). (2015). *Supplemental research bulletin - Issue 5: Traumatic stress and suicide after disasters*. https://www.samhsa.gov/sites/default/files/dtac/srb_sept2015.pdf
18. Centers for Disease Control and Prevention. (2018). The continuum of pandemic phases. CDC. <https://www.cdc.gov/flu/pandemic-resources/planning-preparedness/global-planning-508.html>

19. Anesi, G. L. & Manaker, S. (2020). *Coronavirus disease 2019 (COVID-19): Critical care issues*. <https://www.uptodate.com/contents/coronavirus-disease-2019-covid-19-critical-care-issues>
20. Bhatraju, P. K., Ghassemieh, B. J., Nichols, M., Kim, R., Jerome, K. R., Nalla, A. K., Greninger, A. L., Pipavath, S., Wurfel, M. M., Evans, L., Kritek, P. A., West, R. E., et al. (2020). Covid-19 in Critically Ill Patients in the Seattle Region. *New England Journal of Medicine*. 10.1056/NEJMoa2004500. <https://www.nejm.org/doi/full/10.1056/nejmoa2004500>
21. Substance Abuse and Mental Health Services Administration (SAMHSA). (2020). *Phases of Disaster*. <https://www.samhsa.gov/dtac/disaster-behavioral-health-resources>
https://store.samhsa.gov/sites/default/files/SAMHSA_Digital_Download/pep21-02-01-001.pdf