

# COVID-19 THERAPIES QUICK GUIDE

Type of COVID-19 Therapies	Pre-Exposure Prophylaxis	Monoclonal Antibodies	Oral Antivirals	IV Antivirals
What is available	Evusheld	Bebtelovimab	Lagevrio (molnupiravir) Paxlovid	VEKLURY (remdesivir)
Who is it for?	<p>Adult and pediatric patients (at least 12 years of age and older weighing at least 40 kg) who:</p> <ul style="list-style-type: none"> <li>are not currently infected with SARS-CoV-2 (COVID-19).</li> <li>have not had a known recent exposure to an individual infected with SARS CoV-2 (COVID-19).</li> <li>are moderately to severely immunocompromised.</li> <li>are not recommended to receive the COVID-19 vaccine.</li> </ul>	<p>Adult and pediatric patients (at least 12 years of age and older weighing at least 40 kg) who:</p> <ul style="list-style-type: none"> <li>have high risk for progressing to severe COVID-19, including hospitalization or death.</li> </ul>	<p><b>Paxlovid:</b></p> <ul style="list-style-type: none"> <li>Adults and pediatric patients (12 years of age and older weighing at least 40 kg).</li> </ul> <p><b>Lagevrio:</b></p> <ul style="list-style-type: none"> <li>Adults at high risk for progressing to severe COVID-19, including hospitalization or death.</li> <li>Adults who are not recommended alternative COVID-19 treatment options or do not have accessibility to other COVID-19 treatment options.</li> </ul>	<p>Adults and pediatric patients (12 years of age and older weighing at least 40 kg) who:</p> <ul style="list-style-type: none"> <li>are hospitalized</li> <li>are not hospitalized and at high risk for progression to severe COVID-19, including hospitalization or death.</li> </ul> <p>Pediatric patients weighing 3.5 kg to less than 40 kg or pediatric patients less than 12 years of age weighing at least 3.5 kg who:</p> <ul style="list-style-type: none"> <li>are hospitalized</li> <li>are not hospitalized and at high risk for progression to severe COVID-19, including hospitalization or death.</li> </ul>
When to access?	2 weeks after vaccine	Within 7 days or less after testing positive for COVID-19	Within 5 days or less after testing positive for COVID-19	Within 7 days or less after testing positive for COVID-19
How to access?	Healthcare Providers	Healthcare Providers	Healthcare Providers, Local pharmacies	Inpatient healthcare facilities

More information can be found at [hhs.gov](https://www.hhs.gov)



DOH 825-035 May 2022

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email [civil.rights@doh.wa.gov](mailto:civil.rights@doh.wa.gov).