

The best way for your patient to get up-to-date information about when and how to get vaccinated is by visiting the Washington State Department of Health website *CovidVaccineWA.org* or by calling the Department of Health phone line **1-800-525-0127**, then press #. (For interpretive services, say your language when the call is answered.)

Concerns & Example Responses

They expressed concern that Novavax is a new COVID-19 vaccine and they don't want to be a research experiment.

"I understand your hesitation, but vaccine developers didn't skip any testing steps following the appropriate safety measures including a rigorous three phase clinical trials process with very clear outcomes. The technology for the Novavax vaccine has been used in other common vaccines such as some flu, hepatitis B, HPV, and shingles vaccines. The Novavax COVID-19 vaccine has already been authorized in more than 40 countries and the EU and there have been over 1 million COVID-19 Novavax vaccine doses administered safely worldwide, and there are very strong vaccine safety systems to catch any warning signs early."

They've experienced racism or faced another form of discrimination when receiving medical care.

"Your experience sounds frustrating and hurtful. I've heard from other patients about negative experiences they've had when getting medical care, and I realize that discrimination exists in health care. It upsets me that this is still happening."

After listening and further exploring the patient's concerns, consider: "The vaccine is strongly recommended, but completely voluntary. I only want to make sure that you get the protection/care you need."

They mention harmful medical practices or experiments on Black and Indigenous people in the past.

"You're absolutely right. The history of medical harm should not be ignored. I acknowledge that scientists engaged in many harmful and racist practices in the name of medicine. We are trying to handle the COVID-19 vaccine differently and trying to be as transparent as possible. I want to make sure this feels like the right decision for you."

They shared that they believe there is no point in getting vaccinated if they can still get infected.

"I understand. Although there is still a chance of COVID-19 infection after vaccination, the vaccines were designed to prevent severe illness, hospitalization, and death from COVID-19 and are still successful at doing so. If you do get sick after you're fully vaccinated, you will still have some benefit from the vaccine because you are more likely to only get a mild case instead of a serious case."

They share concerns about politics or the government's involvement.

"You are right, it has gotten political. Would it by OK if I shared my medical perspective on this issue?" If granted permission: "I've looked at the results, and this vaccine does really protect people from COVID. I want you to have the benefit of it."



Frequently Asked Questions & Example Responses

Do I still need to get vaccinated? Is COVID-19 still a threat?

"Yes and yes. COVID-19 is still infecting people every day and I strongly recommend you continue to stay up to date on vaccinations. The updated vaccine will help protect against serious illness and hospitalization from COVID-19 and may also protect people around you."

Should I get vaccinated even if I already had COVID-19?

"Yes, I strongly recommend you get vaccinated despite previously contracting COVID-19 but you should wait to receive the vaccine until after you recover and complete your isolation period. For maximum effectiveness, individuals who recently had COVID-19 may consider delaying COVID-19 vaccination by 3 months from the start of their symptoms or positive test. Getting a COVID-19 vaccine after recovering from COVID-19 infection provides added protection to your immune system. People who already had COVID-19 and do not get vaccinated after their recovery are more likely to get COVID-19 again than those who get vaccinated after their recovery."

What are the side effects?

"Like other vaccines, the most common side effects are a sore arm, tiredness, headache, and muscle pain, which were generally reported to be mild to moderate in severity and occurred within two days after vaccination, and most went away within one to two days. These symptoms are a sign that the vaccine is prompting an immune response as intended. The health risks if you are infected with COVID-19 are much higher than the risk of vaccine side effects."

What's in the Novavax COVID-19 vaccine?

"The Novavax COVID-19 vaccine is a protein subunit based vaccine that contains an additive, along with fats and sugars to help the vaccine work better in the body. This vaccine does not use mRNA. The ingredients are all safe and needed for the vaccine to do its job. It does not contain human cells (including fetal cells), the virus that causes COVID-19, any animal by-products including pork products, latex, preservatives, or tracking devices. It was not grown in eggs and contains no egg products."

What is Myocarditis and is there a connection to COVID-19 vaccination?

"Myocarditis (and pericarditis) are terms to describe inflammation in or around the heart. The body's immune system can often cause this inflammation in response to an infection, such as viruses. Myocarditis is not very common and rarely fatal, with less than 200,000 cases in the United States each year and less than 2% of these cases fatal. The risk of myocarditis after COVID-19 vaccination is extremely rare, but there have been a few reported cases. There have been no deaths from myocarditis determined to be caused by COVID-19 vaccination in the United States. An individual is more likely to develop myocarditis after infection with COVID-19 than from the vaccine."

Can I get the vaccine if I am pregnant or am planning to have a baby one day?

"COVID-19 vaccination is strongly recommended for anyone 6 months and older, including people who are pregnant, breastfeeding, or trying to get pregnant. Pregnant and recently pregnant people are more likely to get severely ill with COVID-19 compared with non-pregnant people. There is no evidence that COVID-19 vaccines cause female or male fertility problems and the ingredients and antibodies developed following COVID-19 vaccination have not been found to cause any problems with becoming pregnant. In fact, some studies show that a vaccinated parent may pass protective antibodies against COVID-19 along to their baby through pregnancy and lactation."

Is it safe to receive other vaccines at the same time as the COVID-19 vaccine?

"Yes. The Advisory Committee on Immunization Practices (ACIP) changed their recommendations on May 12, 2021. You can now get a COVID-19 vaccine at the same time that you get other vaccines.

You do not need to schedule other recommended vaccines separately from COVID-19 vaccination. A COVID-19 vaccine appointment is another opportunity to get caught up on all recommended vaccines."

For other frequently asked questions, visit: www.doh.wa.gov/covidvaccine/faq

Adapted from:

- 1. Communication skills for the COVID vaccine. VitalTalk. vitaltalk.org/guides/communication-skills-for-the-covid-vaccine/. Published 1/8/2021.
- Gagneur A. Motivational interviewing: A powerful tool to address vaccine hesitancy. Can Commun Dis Rep. 2020;46(4):93-97. Published 2020 Apr 2. doi:10.14745/ccdr.v46i04a06
- 3. Opel DJ, Lo B, Peek ME. Addressing Mistrust About COVID-19 Vaccines Among Patients of Color. Ann Intern Med. Published online February 9, 2021 at <u>acpjournals.org/doi/10.7326/M21-0055</u>. doi:10.7326/m21-0055

DOH 825-051 CS March 2024 To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov.