

# Washington State Influenza Summary 2021-2022 Season

Washington State Department of Health, Communicable Disease Epidemiology

The Department of Health (DOH), in collaboration with local health jurisdictions and the Centers Disease Control and Prevention (CDC), performed surveillance for influenza during the 2021-2022 season using several different systems. This report summarizes data collected through key systems from October 3, 2021 to October 1, 2022 (week 40 of 2021 through week 39 of 2022).

*Due to the COVID-19 pandemic, data reported from the various influenza surveillance systems may not represent an accurate reflection of influenza activity. Results should be interpreted with caution, especially where comparisons are made to previous influenza seasons.*

## National Summary

The severity of the 2021–22 influenza season was low. The season was characterized by two waves of influenza A activity. The first wave of influenza activity peaked in mid-December throughout the country, but the timing of peak activity during the second wave varied by region, ranging from mid-March to May. Notably, the second wave peaked and influenza activity remained elevated nationally later than in any previous seasonal influenza epidemic. The predominant influenza virus throughout both waves was influenza A(H3N2) virus. Influenza activity continued from October 2021 through mid-June 2022.

<https://www.cdc.gov/mmwr/volumes/71/wr/mm7129a1.htm>

## Washington State Summary


In Washington State, twenty-six laboratory-confirmed influenza-associated deaths and sixteen influenza-like illness outbreaks in long-term care facilities were reported for the 2021-2022 season. Illness attributed to influenza A viruses predominated, with very little influenza B activity. The adoption of COVID-19–related mitigation measures such as masking, staying home, and limiting gatherings might have had an impact on the timing or severity of influenza activity.

## Influenza Laboratory Surveillance Data


### Laboratory Data: World Health Organization (WHO) & National Respiratory and Enteric Virus Surveillance System (NREVSS) Data Reported to CDC

For the 2021-2022 influenza season, CDC has generated separate graphs of data reported to CDC by public health laboratories (Figure 1) and commercial laboratories (Figure 2).

**Figure 1: Influenza Positive Tests Reported to CDC, WA Public Health Laboratories**



**Figure 2: Influenza Positive Tests Reported to CDC, WA Commercial Laboratories**




## Outpatient Influenza-like Illness Surveillance

### Outpatient Influenza-like Illness Surveillance Network (ILINet) Data

Information on patient visits to health care providers for influenza-like illness is collected through the U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet). Each week, healthcare providers in Washington reported data to CDC on the total number of patients seen and the number of those patients with influenza-like illness (ILI) by age group. For the purposes of ILINet, ILI is defined as fever (temp 100°F/37.8°C or higher) plus cough and/or sore throat. More information about ILINet is available [here](#).

*It should be noted that in addition to the overarching impacts of COVID-19 on influenza surveillance systems, interpretation of ILINet data for the 2021-2022 influenza season should take into account the following COVID-19 impacts: changes in the health seeking behavior at ILINet sentinel sites, changes to provider swabbing at ILINet sentinel sites due to the availability of telehealth and respiratory clinics, and limited ability to distinguish between ILI and COVID-19 symptoms.*

**Figure 3: Percentage of ILI Visits Reported by Sentinel Providers, Washington, 2021-2022**



## Influenza-like Illness Syndromic Surveillance Data


### ESSENCE Syndromic Surveillance Data

Figure 4 shows the proportion of visits at a subset of emergency departments across Washington for a chief complaint of influenza-like illness, or discharge diagnosis of influenza, by CDC week. For this purpose, ILI is defined as “influenza” or fever with cough or fever with sore throat.

*It should be noted that in addition to the overarching impacts of COVID-19 on influenza surveillance systems, interpretation of syndromic surveillance data for the 2021-2022 influenza season should take into account the following COVID-19 impacts: changes in the health seeking behavior at syndromic surveillance sites and limited ability to distinguish between ILI and COVID-19 symptoms.*

For more information about Syndromic Surveillance in Washington State, see [here](#).

**Figure 4: Syndromic Surveillance, Percentage of Hospital Visits for a Chief Complaint of ILI, or Discharge Diagnosis of Influenza, by CDC Week, Washington, 2018-2022**



## Influenza-like Illness Outbreaks in Long Term Care Facilities

Long term care facilities are required to report all suspected and confirmed outbreaks to their [local health jurisdiction](#) per Washington Administrative Code (WAC) [246-101-305](#). Long-term care facilities are required to report the following:

- A sudden increase in acute febrile respiratory illness over the normal background rate (e.g., 2 or more cases of acute respiratory illness occurring within 72 hours of each other) OR
- Any resident who tests positive for influenza

This count of Influenza-like Illness Outbreaks does not include lab-confirmed COVID-19 outbreaks. For more information on COVID-19 outbreaks, see the WA DOH [COVID-19 Long Term Care Monthly Report](#)

Recommendations for prevention and control of influenza outbreaks in long-term care facilities are available [here](#).

Local health jurisdictions in turn report long-term care facility influenza-like illness outbreaks to the Washington State Department of Health.


Since Week 40 of 2021, 16 influenza-like illness outbreaks in long-term care facilities have been reported to the Washington State Department of Health.

### Other Causes of Respiratory Infections

During the 2021-2022 season, the following respiratory viruses were reported to the National Respiratory and Enteric Surveillance System (NREVSS). NREVSS does not capture COVID-19 testing data. For more information on COVID-19, see [here](#).

For more information about NREVSS, see [here](#).

**Figure 5: Respiratory and Enteric Viruses, Washington, 2021-2022 Season to Date**



## Laboratory Confirmed Influenza-Associated Deaths

### Reported Laboratory-Confirmed Influenza Associated Deaths

Note that these counts reflect only deaths officially reported to the Washington State Department of Health and are likely underreported for a variety of reasons. Influenza may not be listed as a cause of death, influenza testing may not have been performed, and lab-confirmed influenza deaths may not have been appropriately reported to public health. CDC has published information about estimating seasonal influenza-associated deaths: <https://www.cdc.gov/flu/about/burden/how-cdc-estimates.htm>

Twenty-six laboratory-confirmed influenza-associated deaths have been reported since week 40 of 2021, 26 influenza A, 0 influenza B, and 0 type unknown.

**Table 1: Count and rate of reported laboratory-confirmed influenza-associated deaths by age group, Washington, 2021-2022 season**

Age Group (in years)	Count of Deaths	Death Rate (per 100,000 population)
0-4	0	0.00
5-17	1	0.08
18-29	0	0.00
30-49	1	0.05
50-64	3	0.21
65+	21	1.79
Total	26	0.35

### Reported Laboratory-Confirmed Influenza-Associated Deaths, Past Seasons

Lab-confirmed influenza-associated death totals reported to the Department of Health for past seasons are presented below in Table 2. Note that for the purposes of tables 1 and 2, each influenza season runs from week 40 of one year to week 39 of the next (roughly October to October). Past season summaries are available on the [Department of Health website](#)

**Table 2: Count of Reported Laboratory-Confirmed Influenza-Associated Deaths, Past Seasons to Week 39 and Total**

Season	Count of Deaths Reported for the Entire Season (week 40 to week 39)
2021-2022, to date	26
2020-2021	0
2019-2020	114
2018-2019	245
2017-2018	296
2016-2017	278
2015-2016	67
2014-2015	156
2013-2014	81

## Additional Resources

International Influenza Data: <https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-updates/current-influenza-update>

National Influenza Surveillance Report: <http://www.cdc.gov/flu/weekly/>

Washington DOH Influenza Information for Public Health and Healthcare Providers:

<http://www.doh.wa.gov/ForPublicHealthandHealthcareProviders/PublicHealthSystemResourcesandServices/Immunization/InfluenzaFluInformation>

Washington Local Health Department Influenza Surveillance Reports:

Clark County: <https://clark.wa.gov/sites/default/files/media/document/2021-05/Clark%20County%20Weekly%20Influenza%20Update.pdf>

King County: <https://kingcounty.gov/depts/health/communicable-diseases/disease-control/influenza.aspx>

Kitsap County: <https://kitsappublichealth.org/Respiratory.pdf>

Pierce County: <https://www.tpchd.org/healthy-people/provider-resources/disease-information-for-providers/influenza/influenza-reports>

Whatcom County: <https://www.whatcomcounty.us/3532/Whatcom-County-Weekly-Influenza-Report>

Yakima County: <http://www.yakimacounty.us/365/RSV-Flu-Stats>

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email [civil.rights@doh.wa.gov](mailto:civil.rights@doh.wa.gov).