

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-005 Radiation protection programs. (1) Each specific licensee shall develop, document, and implement a radiation protection program sufficient to ensure compliance with the provisions of this chapter.

(2) The licensee shall use, to the extent practical, procedures and engineering controls based upon sound radiation protection principles to achieve occupational doses and doses to members of the public that are as low as is reasonably achievable (ALARA).

(3) The licensee shall review the radiation protection program content and implementation at ~~((the frequency specified in the license))~~ least annually.

(4) To implement the ALARA requirements of subsection (2) of this section, and notwithstanding the requirements of WAC 246-221-060, a constraint on air emission of radioactive material to the environment, excluding radon-220, radon-222 and their daughters, shall be established by licensees such that the individual member of the public likely to receive the highest dose will not be expected to receive a total effective dose equivalent in excess of 0.1 mSv (10 mrem) per year from these emissions. This dose constraint does not apply to sealed sources or to accelerators less than 200MeV. If a licensee subject to this requirement exceeds this dose constraint, the licensee shall report the exceedance as provided in WAC 246-221-260 and promptly take appropriate corrective action to ensure against recurrence.

(5) Each licensee shall maintain records of the radiation protection program, including:

(a) The provisions of the program; and

(b) Audits, where required, and other reviews of program content and implementation.

AMENDATORY SECTION (Amending WSR 18-21-020, filed 10/4/18, effective 11/4/18)

WAC 246-221-010 Occupational dose limits for adults. (1) The licensee or registrant shall control the occupational dose to individual adults, except for planned special exposures pursuant to WAC 246-221-030, to the following dose limits:

(a) An annual limit, which is the more limiting of:

(i) The total effective dose equivalent being equal to 0.05 Sv ~~((5))~~ five rem); or

(ii) The sum of the deep dose equivalent and the committed dose equivalent to any individual organ or tissue other than the lens of the eye being equal to 0.50 Sv (50 rem).

(b) The annual limits to the lens of the eye, to the skin of the whole body, and to the skin of the extremities which are:

(i) A lens dose equivalent of 0.15 Sv (15 rem); and

(ii) A shallow dose equivalent of 0.50 Sv (50 rem) to the skin of the whole body or to the skin of any extremity.

(2) Doses received in excess of the annual limits, including doses received during accidents, emergencies, and planned special exposures, must be subtracted from the limits specified in WAC

246-221-030 for planned special exposures that the individual may receive during the current year and during the individual's lifetime.

(3) When the external exposure is determined by measurement with an external personal monitoring device, the deep-dose equivalent must be used in place of the effective dose equivalent, unless the effective dose equivalent is determined by a dosimetry method approved by the NRC or the department. The assigned deep-dose equivalent must be for the part of the body receiving the highest exposure. The assigned shallow dose equivalent shall be the dose averaged over the contiguous (~~ten~~) 10 square centimeters of skin receiving the highest exposure. The deep dose equivalent, lens dose equivalent, and shallow dose equivalent may be assessed from surveys or other radiation measurements for the purpose of demonstrating compliance with the occupational dose limits, if the individual monitoring device was not in the region of highest potential exposure, or the results of the individual monitoring are unavailable.

(4) Derived air concentration (DAC) and annual limit on intake (ALI) values are specified in WAC 246-221-290 and may be used to determine the individual's dose and to demonstrate compliance with the occupational dose limits.

(5) Notwithstanding the annual dose limits, the licensee shall limit the soluble uranium intake by an individual to 10 milligrams in a week in consideration of chemical toxicity.

(6) The licensee or registrant shall reduce the dose that an individual may be allowed to receive in the current year by the amount of occupational dose received while employed by any other person during the current year as determined in accordance with WAC 246-221-020.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-015 Compliance with requirements for summation of external and internal doses. (1) If the licensee is required to monitor under both WAC 246-221-090 and 246-221-100, the licensee shall demonstrate compliance with the dose limits by summing external and internal doses. If the licensee is required to monitor only under WAC 246-221-090 or only under WAC 246-221-100, then summation is not required to demonstrate compliance with the dose limits. The licensee may demonstrate compliance with the requirements for summation of external and internal doses under subsections (2), (3), and (4) of this section. The dose equivalents for the lens of the eye, the skin, and the extremities are not included in the summation, but are subject to separate limits.

(2) **Intake by inhalation.** If the only intake of radionuclides is by inhalation, the total effective dose equivalent limit is not exceeded if the sum of the deep dose equivalent divided by the total effective dose equivalent limit, and one of the following, does not exceed unity:

(a) The sum of the fractions of the inhalation ALI for each radionuclide; or

(b) The total number of derived air concentration-hours (DAC-hours) for all radionuclides divided by (~~two thousand~~) 2,000; or

(c) The sum of the calculated committed effective dose equivalents to all significantly irradiated organs or tissues (T) calculated

from bioassay data using appropriate biological models and expressed as a fraction of the annual limit. For purposes of this requirement, an organ or tissue is deemed to be significantly irradiated if, for that organ or tissue, the product of the weighting factors, w_T , and the committed dose equivalent, $H_{T,50}$, per unit intake is greater than (~~(ten)~~) 10 percent of the maximum weighted value of H_{50} , that is, $w_T H_{T,50}$, per unit intake for any organ or tissue.

(3) **Intake by oral ingestion.** If the occupationally exposed individual also receives an intake of radionuclides by oral ingestion greater than (~~(ten)~~) 10 percent of the applicable oral ALI, the licensee shall account for this intake and include it in demonstrating compliance with the limits.

(4) **Intake through wounds or absorption through skin.** The licensee shall evaluate and, to the extent practical, account for intakes through wounds or skin absorption. The intake through intact skin has been included in the calculation of DAC for hydrogen-3 and does not need to be evaluated or accounted for pursuant to this section.

(5) **External dose from airborne radioactive material.** Licensees shall, when determining the dose from airborne radioactive material, include the contribution to the deep dose equivalent, lens dose equivalent, and shallow dose equivalent from external exposure to the radioactive cloud. Airborne radioactivity measurements and DAC values shall not be used as the primary means to assess the deep dose equivalent when the airborne radioactive material includes radionuclides other than noble gases or if the cloud of airborne radioactive material is not relatively uniform. The determination of the deep dose equivalent to an individual shall be based upon measurements using instruments or individual monitoring devices.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-030 Requirements for planned special exposures. A licensee or registrant may authorize an adult worker to receive doses in addition to and accounted for separately from the doses received under the limits specified in WAC 246-221-010 provided that each of the following conditions is satisfied:

(1) The licensee or registrant authorizes a planned special exposure only in an exceptional situation when alternatives that might avoid the dose estimated to result from the planned special exposure are unavailable or impractical.

(2) The licensee or registrant, and employer if the employer is not the licensee or registrant, specifically authorizes the planned special exposure, in writing, before the exposure occurs.

(3) Before a planned special exposure, the licensee or registrant ensures that each individual involved is:

(a) Informed of the purpose of the planned operation; and

(b) Informed of the estimated doses and associated potential risks and specific radiation levels or other conditions that might be involved in performing the task; and

(c) Instructed in the measures to be taken to keep the dose ALARA considering other risks that may be present.

(4) Prior to permitting an individual to participate in a planned special exposure, the licensee or registrant ascertains prior doses as required by WAC 246-221-020(2) during the lifetime of the individual for each individual involved.

(5) Subject to WAC 246-221-010(2), the licensee or registrant shall not authorize a planned special exposure that would cause an individual to receive a dose from all planned special exposures and all doses in excess of the limits to exceed:

(a) The numerical values of any of the dose limits in WAC 246-221-010(1) in any year; and

(b) Five times the annual dose limits in WAC 246-221-010(1) during the individual's lifetime.

(6) The licensee or registrant maintains records that describe:

(a) The exceptional circumstances requiring the use of a planned special exposure;

(b) The name of the management official who authorized the planned special exposure and a copy of the signed authorization;

(c) What actions were necessary;

(d) Why the actions were necessary;

(e) What precautions were taken to assure that doses were maintained ALARA; and

(f) What individual and collective doses were expected to result.

(7) The licensee or registrant records the best estimate of the dose resulting from the planned special exposure in the individual's record and informs the individual, in writing, of the dose within (~~thirty~~) 30 days from the date of the planned special exposure. The dose from planned special exposures shall not be considered in controlling future occupational dose of the individual under WAC 246-221-010(1) but shall be included in evaluations required by subsections (4) and (5) of this section.

(8) The licensee or registrant submits a written report in accordance with WAC 246-221-265.

AMENDATORY SECTION (Amending WSR 14-01-077, filed 12/16/13, effective 1/16/14)

WAC 246-221-040 Determination of internal exposure of individuals to concentrations of radioactive materials in restricted areas.

(1) For purposes of assessing dose used to determine compliance with occupational dose equivalent limits, the licensee shall, when required under WAC 246-221-100, take suitable and timely measurements of:

(a) Concentrations of radioactive materials in air in work areas;
or

(b) Quantities of radionuclides in the body; or

(c) Quantities of radionuclides excreted from the body; or

(d) Combinations of these measurements.

(2) Unless respiratory protective equipment is used, as provided in WAC 246-221-117, or the assessment of intake is based on bioassays, the licensee shall assume that an individual inhales radioactive material at the airborne concentration in which the individual is present.

(3) When specific information on the physical and biochemical properties of the radionuclides taken into the body or the behavior or the material in an individual is known, the licensee may:

(a) Use that information to calculate the committed effective dose equivalent, and, if used, the licensee shall document that information in the individual's record; and

(b) Upon prior approval of the department, adjust the DAC or ALI values to reflect the actual physical and chemical characteristics of airborne radioactive material, for example, aerosol size distribution or density; and

(c) Separately assess the contribution of fractional intakes of Class D, W, or Y compounds of a given radionuclide to the committed effective dose equivalent. See WAC 246-221-290.

(4) If the licensee chooses to assess intakes of Class Y material using the measurements given in subsection (1)(b) or (c) of this section, the licensee may delay the recording and reporting of the assessments for periods up to seven months, unless otherwise required by WAC 246-221-250 or 246-221-260. This delay permits the licensee to make additional measurements basic to the assessments.

(5) If the identity and concentration of each radionuclide in a mixture are known, the fraction of the DAC applicable to the mixture for use in calculating DAC-hours shall be either:

(a) The sum of the ratios of the concentration to the appropriate DAC value, that is, D, W, or Y, from WAC 246-221-290 for each radionuclide in the mixture; or

(b) The ratio of the total concentration for all radionuclides in the mixture to the most restrictive DAC value for any radionuclide in the mixture.

(6) If the identity of each radionuclide in a mixture is known, but the concentration of one or more of the radionuclides in the mixture is not known, the DAC for the mixture shall be the most restrictive DAC of any radionuclide in the mixture.

(7) When a mixture of radionuclides in air exists, a licensee may disregard certain radionuclides in the mixture if:

(a) The licensee uses the total activity of the mixture in demonstrating compliance with the dose limits in WAC 246-221-010 and in complying with the monitoring requirements in WAC 246-221-100; and

(b) The concentration of any radionuclide disregarded is less than ~~((ten))~~ 10 percent of its DAC; and

(c) The sum of these percentages for all of the radionuclides disregarded in the mixture does not exceed ~~((thirty))~~ 30 percent.

(8) When determining the committed effective dose equivalent, the following information may be considered:

(a) In order to calculate the committed effective dose equivalent, the licensee may assume that the inhalation of one ALI, or an exposure of 2,000 DAC-hours, results in a committed effective dose equivalent of 0.05 Sv ~~((5))~~ five rem) for radionuclides that have their ALIs or DACs based on the committed effective dose equivalent.

(b) For an ALI and the associated DAC determined by the nonstochastic organ dose limit of 0.50 Sv (50 rem), the intake of radionuclides that would result in a committed effective dose equivalent of 0.05 Sv ~~((5))~~ five rem), that is, the stochastic ALI, is listed in parentheses in Table I of WAC 246-221-290. The licensee may, as a simplifying assumption, use the stochastic ALIs to determine committed effective dose equivalent. However, if the licensee uses the stochastic ALIs, the licensee shall also demonstrate that the limit in WAC 246-221-010 (1)(a)(ii) is met.

AMENDATORY SECTION (Amending WSR 17-12-046, filed 6/1/17, effective 7/2/17)

WAC 246-221-055 Dose equivalent to an embryo/fetus. (1) The licensee or registrant shall ensure that the dose equivalent to an embryo/fetus during the entire pregnancy, due to occupational exposure of a declared pregnant woman, does not exceed ((5)) five mSv (0.5 rem).

(2) Once pregnancy has been declared, the licensee or registrant shall make every effort to avoid substantial variation above a uniform monthly exposure rate to a declared pregnant woman in order to satisfy the limit in subsection (1) of this section.

(3) If by the time the woman declares pregnancy to the licensee or registrant, the dose equivalent to the embryo/fetus has exceeded ((5)) five mSv (0.5 rem), or is within 0.50 mSv (0.05 rem) of this dose, the licensee or registrant shall be deemed to be in compliance with subsection (1) of this section if the additional dose equivalent to the embryo/fetus does not exceed 0.50 mSv (0.05 rem) during the remainder of the pregnancy.

(4) The dose equivalent to an embryo/fetus shall be taken as the sum of:

(a) The deep dose equivalent to the declared pregnant woman; and

(b) The dose equivalent to the embryo/fetus from radionuclides in the embryo/fetus and radionuclides in the declared pregnant woman.

(5) The licensee or registrant shall maintain the records of dose equivalent to an embryo/fetus with the records of dose equivalent to the declared pregnant woman. The declaration of pregnancy, including the estimated date of conception, shall also be kept on file, but may be maintained separately from the dose records.

AMENDATORY SECTION (Amending WSR 14-01-077, filed 12/16/13, effective 1/16/14)

WAC 246-221-060 Dose limits for individual members of the public. (1) Each licensee or registrant shall conduct operations so that:

(a) The total effective dose equivalent to individual members of the public from the licensed or registered operation does not exceed ((4)) one mSv (0.1 rem) in a year, exclusive of the dose contributions from background radiation, from any medical administration the individual has received, from exposure to individuals administered radioactive material and released under chapter 246-240 WAC, from voluntary participation in medical research programs, and from the licensee's or registrant's disposal of radioactive material into sanitary sewerage in accordance with WAC 246-221-190; and

(b) The dose in any unrestricted area from external sources, exclusive of the dose contributions from patients administered radioactive material and released under chapter 246-240 WAC, does not exceed 0.02 mSv (0.002 rem) in any one hour.

(2) If the licensee or registrant permits members of the public to have access to restricted areas, they shall be escorted and the limits for members of the public continue to apply to those individuals.

(3) Notwithstanding subsection (1) of this section, a licensee or registrant may continue to operate a facility constructed and put into operation prior to January 1, 1994, where the annual dose limit for an individual member of the public is more than ((±)) one mSv (0.1 rem) and less than ((5)) five mSv (0.5 rem) total effective dose equivalent, if:

(a) The facility's approved operating conditions for each radiation source remain the same. Any increase in the following operating conditions shall require reevaluation by the department and modification of the facility shielding applicable to the source of radiation to meet the ((±)) one mSv (0.1 rem) total effective dose equivalent limit for individual members of the public: Size of the radiation source, workload, or occupancy factors associated with the source of radiation; and

(b) Any change in the permanent shielding of the facility due to remodeling, repair or replacement requires the facility to meet the ((±)) one mSv (0.1 rem) total effective dose equivalent limit for individual members of the public for areas affected by that portion of the shielding.

(4) Each licensee or registrant shall maintain records sufficient to demonstrate compliance with the dose limit for individual members of the public.

AMENDATORY SECTION (Amending WSR 14-01-077, filed 12/16/13, effective 1/16/14)

WAC 246-221-080 Leak tests. (1) Each sealed radioactive source possessed under the provisions of a specific license, other than hydrogen-3 (tritium), with a half-life greater than ((~~thirty~~)) 30 days and in any form other than gas, shall be tested and results obtained for leakage or contamination prior to initial use and at six-month intervals or as specified by the license, except that each source designed for the purpose of emitting alpha particles shall be tested at intervals not to exceed three months. If at any other time there is reason to suspect that a sealed source might have been damaged, it shall be tested for leakage and results obtained before further use. In the absence of a certificate from a transferor indicating that a test for leakage has been made within six months prior to the transfer (three months for a source designed to emit alpha particles), the sealed source shall not be put into use until tested and the results received.

(2) Leak tests shall be capable of detecting the presence of 185 Bq (0.005 microcurie) of removable contamination. The results of leak tests made pursuant to subsection (1) of this section shall be recorded in units of becquerel or microcuries and shall be maintained for inspection by the department. Any test conducted pursuant to subsection (1) of this section which reveals the presence of 185 Bq (0.005 microcurie) or more of removable contamination shall be considered evidence that the sealed source is leaking. The licensee shall immediately withdraw the source from use shall take action to prevent the spread of contamination and shall cause it to be decontaminated and repaired or to be disposed in accordance with WAC 246-232-080. If a sealed source shows evidence of leaking, a report shall be filed with

the department within five days of the test, describing the equipment involved, the test results, and the corrective action taken.

(3) Test samples shall be taken from the sealed source or from the internal surfaces or the opening of the container in which the sealed source is stored or from surfaces of devices or equipment in which the sealed source is permanently mounted. Tests for contamination and leakage may be made by wiping appropriate accessible surfaces on which one might expect contamination to accumulate and measuring these wipes for transferred contamination. Test samples shall also be taken from the interior surfaces of the container in which a sealed source of radium is stored.

(4) Leak tests are required for sealed radioactive sources that are greater than 3.7 MBq (100 microcuries) for beta and gamma emitting sources and greater than 370 KBq (10 microcuries) for sources designed to emit alpha particles.

(5) Tests for leakage or contamination shall be performed by persons specifically authorized by the department, an agreement state, or the NRC to perform such services.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-090 Personnel monitoring for external dose. Each licensee or registrant shall monitor occupational exposure from sources of radiation at levels sufficient to demonstrate compliance with the occupational dose limits of WAC 246-221-010, 246-221-030, 246-221-050 and 246-221-055.

(1) Each licensee or registrant shall monitor occupational exposure to radiation from licensed (or registered) and unlicensed (or unregistered) radiation sources under the control of the licensee or registrant and shall supply and shall require the use of individual monitoring devices by:

(a) Each adult likely to receive, in one year from sources external to the body, a dose in excess of (~~ten~~) 10 percent of the applicable limits specified in WAC 246-221-010(1).

(b) Each minor likely to receive, in one year from sources external to the body, a deep dose equivalent in excess of (~~±~~) one mSv (0.1 rem), a lens dose equivalent in excess of 1.5 mSv (0.15 rem), or a shallow dose equivalent to the skin or to the extremities in excess of (~~5~~) five mSv (0.5 rem).

(c) Each declared pregnant woman likely to receive during the entire pregnancy, from radiation sources external to the body, a deep dose equivalent in excess of (~~±~~) one mSv (0.1 rem). All of the occupational dose limits specified in WAC 246-221-010 continue to be applicable to the declared pregnant worker as long as the embryo/fetus dose limit is not exceeded.

(d) Each individual who enters a high or very high radiation area.

(2) Personnel monitoring devices assigned to an individual:

(a) Shall not intentionally be exposed to give a false or erroneous reading;

(b) Shall be assigned to one individual per exposure interval (i.e., weekly, monthly) and used to determine exposure for that individual only;

(c) Shall not be worn by any individual other than that individual originally assigned to the device;

(d) Personnel monitoring devices that are exposed while not being worn by the assigned individual shall be processed and recorded as soon as possible. A replacement monitoring device shall be assigned to the individual immediately. A record of the circumstances of the exposure shall be retained.

(3) All personnel dosimeters, except for direct and indirect reading pocket ionization chambers and those dosimeters used to measure the dose to any extremities, that require processing to determine the radiation dose and that are utilized by licensees or registrants to comply with subsection (1) of this section, with other applicable provisions of chapters 246-220 through 246-255 WAC, or with conditions specified in a licensee's license must be processed and evaluated by a dosimetry processor:

(a) Holding current personnel dosimetry accreditation from either the National Voluntary Laboratory Accreditation Program (NVLAP) of the National Institute of Standards and Technology (formerly known as the National Bureau of Standards) or the United States Department of Energy Laboratory Accreditation Program for Personnel Dosimetry Systems (DOELAP); and

(b) Approved in this accreditation process for the type of radiation or radiations included in the NVLAP or DOELAP program that most closely approximate the type of radiation or radiations for which the individual wearing the dosimeter is monitored.

(4) For the purposes of this section "dosimetry processor" means an individual or an organization that processes and evaluates personnel monitoring devices in order to determine the radiation dose delivered to the device.

(5) Each licensee or registrant shall maintain records of doses received by all individuals for whom monitoring was required under subsection (1) of this section, and records of doses received during planned special exposures, accidents, and emergency conditions. Assessments of dose equivalent and records made using units in effect before January 1, 1994, need not be changed. These records shall include, when applicable:

(a) The deep dose equivalent to the whole body, lens dose equivalent, shallow dose equivalent to the skin, and shallow dose equivalent to the extremities; and

(b) The total effective dose equivalent when required by WAC 246-221-015; and

(c) The total of the deep dose equivalent and the committed dose to the organ receiving the highest total dose (total organ dose equivalent).

(6) The licensee or registrant shall maintain the records specified in subsection (5) of this section on department Form RHF-5A, in accordance with the instructions provided on the form, or in clear and legible records containing all the information required by Form RHF-5A; and shall update the information at least annually.

(7) Each licensee or registrant shall ensure that individuals, for whom they are required to monitor occupational doses in accordance with subsection (1) of this section, wear individual monitoring devices as follows:

(a) An individual monitoring device used for monitoring the dose to the whole body shall be worn at the unshielded or least shielded location of the whole body likely to receive the highest exposure.

When a protective apron is worn, the location of the individual monitoring device is typically at the neck (collar).

(b) Any additional individual monitoring device used for monitoring the dose to an embryo/fetus of a declared pregnant woman, pursuant to WAC 246-221-055(1), shall be located at the waist under any protective apron being worn by the woman.

(c) An individual monitoring device used for monitoring the lens dose equivalent, to demonstrate compliance with WAC 246-221-010 (1)(b)(i), shall be located at the neck (collar), outside any protective apron being worn by the monitored individual, or at an unshielded location closer to the eye.

(d) An individual monitoring device used for monitoring the dose to the extremities, to demonstrate compliance with WAC 246-221-010 (1)(b)(ii), shall be worn on the extremity likely to receive the highest exposure. Each individual monitoring device shall be oriented to measure the highest dose to the extremity being monitored.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-100 Personnel monitoring for internal dose. (1) Each licensee shall monitor, to determine compliance with WAC 246-221-040, the occupational intake of radioactive material by and assess the committed effective dose equivalent to:

(a) Adults likely to receive, in ((±)) one year, an intake in excess of ((~~ten~~)) 10 percent of the applicable ALI in Table I, Columns 1 and 2, of WAC 246-221-290;

(b) Minors likely to receive, in one year, a committed effective dose equivalent in excess of ((±)) one mSv (0.1 rem); and

(c) Declared pregnant women likely to receive, during the entire pregnancy, a committed effective dose equivalent in excess of ((±)) one mSv (0.1 rem).

(2) Where necessary or desirable in order to aid in determining the extent of an individual's exposure to concentrations of radioactive material, the department may incorporate license provisions or issue an order requiring a licensee or registrant to make available to the individual appropriate bioassay services and to furnish a copy of the reports of such services to the department.

(3) Each licensee shall maintain records of doses received by all individuals for whom monitoring was required pursuant to subsections (1) and (2) of this section, and records of doses received during planned special exposures, accidents, and emergency conditions. Assessments of dose equivalent and records made using units in effect before January 1, 1994, need not be changed. These records shall include, when applicable:

(a) The estimated intake or body burden of radionuclides;

(b) The committed effective dose equivalent assigned to the intake or body burden of radionuclides;

(c) The specific information used to calculate the committed effective dose equivalent pursuant to WAC 246-221-040;

(d) The total effective dose equivalent when required by WAC 246-221-015; and

(e) The total of the deep dose equivalent and the committed dose to the organ receiving the highest total dose (total organ dose equivalent).

(4) The licensee or registrant shall maintain the records specified in subsection (3) of this section on department Form RHF-5A, in accordance with the instructions provided on the form, or in clear and legible records containing all the information required by Form RHF-5A; and shall update the information at least annually.

AMENDATORY SECTION (Amending WSR 94-01-073, filed 12/9/93, effective 1/9/94)

WAC 246-221-102 Control of access to high radiation areas. (1)

The licensee or registrant shall ensure that each entrance or access point to a high radiation area has one or more of the following features:

(a) A control device that, upon entry into the area, causes the level of radiation to be reduced below that level at which an individual might receive a deep dose equivalent of ((±)) one mSv (0.1 rem) in one hour at ((~~thirty~~)) 30 centimeters from the source of radiation or from any surface that the radiation penetrates; or

(b) A control device that energizes a conspicuous visible or audible alarm signal so that the individual entering the high radiation area and the supervisor of the activity are made aware of the entry; or

(c) Entryways that are locked, except during periods when access to the areas is required, with positive control over each individual entry.

(2) In place of the controls required by subsection (1) of this section for a high radiation area, the licensee or registrant may substitute continuous direct or electronic surveillance that is capable of preventing unauthorized entry.

(3) The licensee or registrant may apply to the department for approval of alternative methods for controlling access to high radiation areas.

(4) The licensee or registrant shall establish the controls required by subsections (1) and (3) of this section in a way that does not prevent individuals from leaving a high radiation area.

(5) The licensee is not required to control each entrance or access point to a room or other area that is a high radiation area solely because of the presence of radioactive materials prepared for transport and packaged and labeled in accordance with the regulations of the United States Department of Transportation provided that:

(a) The packages do not remain in the area longer than three days; and

(b) The dose rate at one meter from the external surface of any package does not exceed 0.1 mSv (0.01 rem) per hour.

(6) The licensee is not required to control entrance or access to rooms or other areas in hospitals solely because of the presence of patients containing radioactive material, provided that there are personnel in attendance who are taking the necessary precautions to prevent the exposure of individuals to radiation or radioactive material in excess of the established limits and to operate within the ALARA provisions of the licensee's radiation protection program.

(7) The licensee or registrant is not required to control entrance or access to rooms or other areas as described in this section if the licensee or registrant has met all the specific requirements for access and control specified in other applicable chapters of these regulations, such as, chapter 246-243 WAC for industrial radiography, chapter 246-225 WAC for X-rays in the healing arts, and chapter 246-229 WAC for particle accelerators.

AMENDATORY SECTION (Amending WSR 14-01-077, filed 12/16/13, effective 1/16/14)

WAC 246-221-110 Surveys. (1) Each licensee or registrant shall make or cause to be made such surveys, as defined in WAC 246-220-010, as may be necessary for the licensee or registrant to establish compliance with these regulations and are reasonable under the circumstances to evaluate the magnitude and extent of radiation levels, concentrations or quantities of radioactive material, and potential radiation hazards. Records of such surveys shall be preserved as specified in WAC 246-221-230. Information on performing surveys may be found in the NRC's Regulatory Guide 8.23 "Radiation Safety Surveys at Medical Institutions."

(2) The licensee shall ensure that instruments and equipment used for quantitative radiation measurements, for example, dose rate and effluent monitoring, are calibrated annually at intervals not to exceed (~~thirteen~~) 13 months for the radiation measured.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-117 Use of individual respiratory protection equipment. If the licensee assigns or permits the use of respiratory protection equipment to limit the intake of radioactive material:

(1) The licensee shall use only respiratory protection equipment that is:

(a) Tested and certified by the National Institute for Occupational Safety and Health (NIOSH); or

(b) Approved by the department on the basis of the licensee's submittal of an application for authorized use of other respiratory protection equipment, including a demonstration by testing, or a demonstration on the basis of reliable test information, that the material and performance characteristics of the equipment are capable of providing the proposed degree of protection under anticipated conditions of use.

(2) The licensee shall implement and maintain a respiratory protection program that includes:

(a) Air sampling sufficient to identify the potential hazard, permit proper equipment selection, and estimate exposures;

(b) Surveys and bioassays, as appropriate, to evaluate actual intakes;

(c) Testing of respirators for operability (user seal check for face sealing devices and functional check for others) immediately prior to each use;

(d) Written procedures regarding:

(i) Monitoring, including air sampling and bioassays;

(ii) Supervision and training of respirator users;

(iii) Fit testing;

(iv) Respirator selection;

(v) Breathing air quality;

(vi) Inventory and control;

(vii) Storage, issuance, maintenance, repair, testing, and quality assurance of respiratory protection equipment;

(viii) Recordkeeping; and

(ix) Limitations on periods of respirator use and relief from respirator use;

(e) Determination by a physician that the individual user is medically fit to use respiratory protection equipment:

(i) Before the initial fitting of a face sealing respirator;

(ii) Before the first field use of nonface sealing respirators; and

(iii) Either every (~~twelve~~) 12 months thereafter, or periodically at a frequency determined by a physician; and

(f) Fit testing, with a fit factor greater than or equal to (~~ten~~) 10 times the APF for negative pressure devices, and a fit factor greater than or equal to (~~five hundred~~) 500 for any positive pressure, continuous flow, and pressure-demand devices, before the first field use of tight fitting, face sealing respirators, and periodically thereafter at a frequency not to exceed one year. Fit testing must be performed with the facepiece operating in the negative pressure mode.

(3) The licensee shall advise each respirator user that the user may leave the area at any time for relief from respirator use in the event of equipment malfunction, physical or psychological distress, procedural or communication failure, significant deterioration of operating conditions, or any other conditions that might require relief.

(4) The licensee shall also consider limitations appropriate to the type and mode of use. When selecting respiratory devices the licensee shall provide for vision correction, adequate communication, low temperature work environments, and the concurrent use of other safety or radiological protection equipment. The licensee shall use equipment in such a way as not to interfere with the proper operation of the respirator.

(5) Standby rescue persons are required whenever one-piece atmosphere-supplying suits, or any combination of supplied air respiratory protection device and personnel protective equipment are used from which an unaided individual would have difficulty extricating himself or herself. The standby persons must be equipped with respiratory protection devices or other apparatus appropriate for the potential hazards. The standby rescue persons shall observe or otherwise maintain continuous communication with the workers (visual, voice, signal line, telephone, radio, or other suitable means), and be immediately available to assist them in case of a failure of the air supply or for any other reason that requires relief from distress. A sufficient number of standby rescue persons must be immediately available to assist all users of this type of equipment and to provide effective emergency rescue if needed.

(6) Atmosphere-supplying respirators must be supplied with respirable air of grade D quality or better as defined by the Compressed Gas Association in publication G-7.1, "Commodity Specification for Air," 1997 and included in the regulations of the Occupational Safety and Health Administration (29 C.F.R. 1910.134 (i)(1)(ii)(A) through (E)). Grade D quality air criteria include:

(a) Oxygen content (v/v) of 19.5-23.5%;

(b) Hydrocarbon (condensed) content of ((5)) five milligrams per cubic meter of air or less;

(c) Carbon monoxide (CO) content of 10 ppm or less;

(d) Carbon dioxide content of 1,000 ppm or less; and

(e) Lack of noticeable odor.

(7) The licensee shall ensure that no objects, materials or substances, such as facial hair, or any conditions that interfere with the face-to-facepiece seal or valve function, and that are under the control of the respirator wearer, are present between the skin of the wearer's face and the sealing surface of a tight-fitting respirator facepiece.

(8) In estimating the dose to individuals from intake of airborne radioactive materials, the concentration of radioactive material in the air that is inhaled when respirators are worn is initially assumed to be the ambient concentration in air without respiratory protection, divided by the assigned protection factor. If the dose is later found to be greater than the estimated dose, the corrected value must be used. If the dose is later found to be less than the estimated dose, the corrected value may be used.

(9) The department may impose restrictions in addition to the provisions of this section, WAC 246-221-113 and 246-221-285, in order to:

(a) Ensure that the respiratory protection program of the licensee is adequate to limit doses to individuals from intakes of airborne radioactive materials consistent with maintaining total effective dose equivalent ALARA; and

(b) Limit the extent to which a licensee may use respiratory protection equipment instead of process or other engineering controls.

(10) The licensee shall obtain authorization from the department before using assigned protection factors in excess of those specified in WAC 246-221-285. The department may authorize a licensee to use higher assigned protection factors on receipt of an application that:

(a) Describes the situation for which a need exists for higher protection factors; and

(b) Demonstrates that the respiratory protection equipment provides these higher protection factors under the proposed conditions of use.

AMENDATORY SECTION (Amending WSR 16-13-054, filed 6/10/16, effective 7/11/16)

WAC 246-221-160 Procedures for picking up, receiving, and opening packages. (1) Each licensee who expects to receive a package containing quantities of radioactive material in excess of the Type A₁ or A₂ quantities specified in WAC 246-231-200 shall make arrangements to receive:

- (a) The package when it is offered for delivery by the carrier;
- or
- (b) Immediate notification from the carrier of the arrival of the package at the carrier's terminal.
- (2) Each licensee who picks up a package of radioactive material from a carrier's terminal shall pick up the package expeditiously upon receipt of notification from the carrier of its arrival.
- (3) Each licensee shall:
- (a) Monitor for radioactive contamination the external surfaces of any package labeled with a Radioactive White I, Yellow II or Yellow III label unless the package contains only radioactive material in the form of gas or in special form as defined in WAC 246-231-010; and
- (b) Monitor the radiation levels of the external surfaces of any package labeled with a Radioactive White I, Yellow II or Yellow III label unless the package contains quantities of radioactive material that are less than or equal to the Type A quantity, as defined in WAC 246-231-200; and
- (c) Monitor all packages known to contain radioactive material for radioactive contamination and radiation levels if the package has evidence of potential contamination, such as packages that are crushed, wet, or damaged.
- (4) Monitoring shall be performed:
- (a) Immediately upon receipt if there is evidence of package degradation or any other evidence of potential contamination or excessive radiation levels; or
- (b) As soon as practicable after receipt, but no later than three hours after the package is received at the licensee's facility if received during the licensee's normal working hours, or no later than three hours from the beginning of the next working day if received after normal working hours.
- (5) The licensee shall immediately notify the final delivery carrier and, by telephone, facsimile, or email, (~~or letter,~~) the department when:
- (a) For normal shipments, removable radioactive surface contamination exceeds either 22 dpm/cm² for beta-gamma emitting radionuclides, all radionuclides with half-lives less than (~~ten~~) 10 days, natural uranium, natural thorium, uranium-235, uranium-238, thorium-232, and thorium-228 and thorium 230 when contained in ores or concentrates; or 2.2 dpm/cm² for all other alpha emitting radionuclides;
- or
- (b) For exclusive use shipments, removable radioactive surface contamination exceeds either 220 dpm/cm² for beta-gamma emitting radionuclides, all radionuclides with half-lives less than (~~ten~~) 10 days, natural uranium, natural thorium, uranium-235, uranium-238, thorium-232, and thorium-228 and thorium 230 when contained in ores or concentrates; or 22 dpm/cm² for all other alpha emitting radionuclides; or
- (c) For normal or exclusive use shipments, external radiation levels exceed two mSv/hour (200 millirem per hour) at any point on the external surface of the package; or
- (d) For exclusive use shipments where the shipment is made in a closed transport vehicle, packages are secured in a fixed position, and no loading or unloading occurs between the beginning and end of transportation, external radiation levels exceed (~~ten~~) 10 mSv/hour (1000 millirem per hour) at any point on the external surface of the package.

(6) Each licensee shall establish and maintain procedures for safely opening packages in which radioactive material is received, and shall assure that such procedures are followed and that due consideration is given to instructions for the type of package being opened and the monitoring of potentially contaminated packaging material (including packages containing radioactive material in gaseous form) to assure that only background levels of radiation are present prior to disposal of such material as nonradioactive waste.

(7) Licensees transferring special form sources to and from a work site in vehicles owned or operated by the licensee are exempt from the contamination monitoring requirements of subsection (3)(a) of this section but are not exempt from the monitoring requirement in subsection (3)(b) of this section for measuring radiation levels to ensure that the source is still properly lodged in its shield.

AMENDATORY SECTION (Amending WSR 94-01-073, filed 12/9/93, effective 1/9/94)

WAC 246-221-190 Disposal by release into sanitary sewerage systems. (1) No licensee shall discharge radioactive material into a sanitary sewerage system unless:

(a) It is readily soluble or it is biological material which is readily dispersible in water;

(b) The quantity of any radioactive material released in any one month, if diluted by the average monthly quantity of water released by the licensee, will not result in an average concentration exceeding the limits specified in WAC 246-221-290, Table III; and

(c) The sum of the fractions for each radionuclide, if more than one radionuclide is released, will not exceed unity; where the fraction for each radionuclide is determined by dividing the actual monthly average concentration of each radionuclide released by the licensee into the sewer by the concentration of that radionuclide listed in Table III of WAC 246-221-290; and

(d) The total quantity of licensed and other radioactive material that the licensee releases into the sanitary sewerage system in a year does not exceed 185 GBq (~~((5))~~ five Ci) of hydrogen-3, 37 GBq (~~((4))~~ one Ci) of carbon-14, and 37 GBq (~~((4))~~ one Ci) of all other radioactive materials combined.

(2) Excreta from individuals undergoing medical diagnosis or therapy with radioactive material shall be exempt from any limitations contained in this section.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-230 Records important to radiation safety. (1) Each licensee or registrant shall make and retain records of activities, program reviews, measurements, and calculations which may be necessary to determine the extent of occupational and public exposure from sources of radiation under the control of the licensee or registrant.

(2) Each record required by this section shall be legible throughout the specified retention period.

(3) Each licensee or registrant shall use the SI units: Becquerel, gray, sievert and coulomb per kilogram, or the special units: Curie, rad, rem, and roentgen, including multiples and subdivisions, and shall clearly indicate the units of all quantities on records required by these regulations.

(4) The licensee or registrant shall make a clear distinction among the quantities entered on the records required by these regulations such as, total effective dose equivalent, total organ dose equivalent, shallow dose equivalent, lens dose equivalent, deep dose equivalent, or committed effective dose equivalent.

(5) Records which must be maintained under this part shall be the original or a reproduced copy or microform if such reproduced copy or microform is duly authenticated by authorized personnel and the microform is capable of producing a clear and legible copy after storage for the period specified by department regulations. The record may also be stored in electronic media with the capability for producing legible, accurate, and complete records during the required retention period. Electronic media data storage systems shall incorporate standard or universally recognized security measures. Records, such as letters, drawings, and specifications, shall include all pertinent information, such as stamps, initials, and signatures.

(6) The licensee shall maintain adequate safeguards against tampering with and loss of records.

(7) The licensee or registrant shall retain the following required records until the department terminates each pertinent license or registration requiring the record, and upon termination of the license or registration, the licensee or registrant shall store for at least (~~thirty~~) 30 years:

(a) Records of prior occupational dose and exposure history as recorded on department Form RHF-4 or RHF-4A, or equivalent;

(b) Records on department Form RHF-5 or RHF-5A, or equivalent, of doses received by all individuals for whom monitoring was required pursuant to WAC 246-221-090 and 246-221-100;

(c) Records of doses received during planned special exposures, accidents, and emergency conditions;

(d) The specific information used to calculate the committed effective dose equivalent pursuant to WAC 246-221-040(3);

(e) Records of the results of surveys to determine the dose from external sources of radiation used, in the absence of or in combination with individual monitoring data, in the assessment of individual dose equivalents;

(f) Records of the results of measurements and calculations used to determine individual intakes of radioactive material and used in the assessment of internal dose;

(g) Records showing the results of air sampling, surveys, and bioassays required pursuant to WAC 246-221-117 (1)(b)(i) and (ii);

(h) Records of the results of measurements and calculations used to evaluate the release of radioactive effluents to the environment.

(8) The licensee or registrant shall retain the following records until the department terminates the pertinent license or registration requiring the record:

(a) Records of waste disposal made under the provisions of WAC 246-221-180, 246-221-190, 246-221-210 and 246-221-220, chapter 246-249 WAC, and any burials in soil as previously authorized;

(b) Records of dose to individual members of the public as required by WAC 246-221-060(4);

(c) Records of the provisions of the radiation protection program as required by WAC 246-221-005.

(9) The licensee or registrant shall retain the following records for three years after the record is made:

(a) Records of testing entry control devices for very high radiation areas as required by WAC 246-221-106(3);

(b) Records used in preparing department Form RHF-4 or RHF-4A;

(c) Records showing the results of general surveys required by WAC 246-221-110 and package surveys required by WAC 246-221-160;

(d) Records of calibrations required by WAC 246-221-110;

(e) Records of program audits and other reviews of the content and implementation of the radiation protection program required by WAC 246-221-005;

(f) Records of waste disposal by decay in storage.

(10) If there is a conflict between the department's regulations in this part, license condition, or other written department approval or authorization pertaining to the retention period for the same type of record, the retention period specified in the regulations in this part for such records shall apply unless the department, under WAC 246-220-050, has granted a specific exemption from the record retention requirements specified in the regulations in this part.

(11) The discontinuance or curtailment of activities does not relieve the licensee or registrant of responsibility for retaining all records required by this section.

AMENDATORY SECTION (Amending WSR 09-06-003, filed 2/18/09, effective 3/21/09)

WAC 246-221-235 Reports of transactions involving nationally tracked sources. Each licensee who manufactures, transfers, receives, disassembles, or disposes of a nationally tracked source shall complete and submit a National Source Tracking Transaction Report as specified in subsections (1) through (5) of this section for each type of transaction.

(1) Each licensee who manufactures a nationally tracked source shall complete and submit a National Source Tracking Transaction Report. The report must include the following information:

(a) The name, address, and license number of the reporting licensee;

(b) The name of the individual preparing the report;

(c) The manufacturer, model, and serial number of the source;

(d) The radioactive material in the source;

(e) The initial source strength in becquerels (curies) at the time of manufacture; and

(f) The manufacture date of the source.

(2) Each licensee that transfers a nationally tracked source to another person shall complete and submit a National Source Tracking Transaction Report. The report must include the following information:

(a) The name, address, and license number of the reporting licensee;

(b) The name of the individual preparing the report;

- (c) The name and license number of the recipient facility and the shipping address;
 - (d) The manufacturer, model, and serial number of the source or, if not available, other information to uniquely identify the source;
 - (e) The radioactive material in the source;
 - (f) The initial or current source strength in becquerels (curies);
 - (g) The date for which the source strength is reported;
 - (h) The shipping date;
 - (i) The estimated arrival date; and
 - (j) For nationally tracked sources transferred as waste under a Uniform Low-Level Radioactive Waste Manifest, the waste manifest number and the container identification of the container with the nationally tracked source.
- (3) Each licensee that receives a nationally tracked source shall complete and submit a National Source Tracking Transaction Report. The report must include the following information:
- (a) The name, address, and license number of the reporting licensee;
 - (b) The name of the individual preparing the report;
 - (c) The name, address, and license number of the person that provided the source;
 - (d) The manufacturer, model, and serial number of the source or, if not available, other information to uniquely identify the source;
 - (e) The radioactive material in the source;
 - (f) The initial or current source strength in becquerels (curies);
 - (g) The date for which the source strength is reported;
 - (h) The date of receipt; and
 - (i) For material received under a Uniform Low-Level Radioactive Waste Manifest, the waste manifest number and the container identification with the nationally tracked source.
- (4) Each licensee that disassembles a nationally tracked source shall complete and submit a National Source Tracking Transaction Report. The report must include the following information:
- (a) The name, address, and license number of the reporting licensee;
 - (b) The name of the individual preparing the report;
 - (c) The manufacturer, model, and serial number of the source or, if not available, other information to uniquely identify the source;
 - (d) The radioactive material in the source;
 - (e) The initial or current source strength in becquerels (curies);
 - (f) The date for which the source strength is reported;
 - (g) The disassemble date of the source.
- (5) Each licensee who disposes of a nationally tracked source shall complete and submit a National Source Tracking Transaction Report. The report must include the following information:
- (a) The name, address, and license number of the reporting licensee;
 - (b) The name of the individual preparing the report;
 - (c) The waste manifest number;
 - (d) The container identification with the nationally tracked source;
 - (e) The date of disposal; and
 - (f) The method of disposal.

(6) The reports discussed in subsections (1) through (5) of this section must be submitted by the close of the next business day after the transaction. A single report may be submitted for multiple sources and transactions. The reports must be submitted to the National Source Tracking System by using:

- (a) The online National Source Tracking System;
- (b) Electronically using a computer-readable format;
- (c) By facsimile;
- (d) By mail to the address on the National Source Tracking Transaction Report Form (NRC Form 748); or
- (e) By telephone with follow-up by facsimile or mail.

(7) Each licensee shall correct any error in previously filed reports or file a new report for any missed transaction within five business days of the discovery of the error or missed transaction. Such errors may be detected by a variety of methods such as administrative reviews or by physical inventories required by regulation. In addition, each licensee shall reconcile the inventory of nationally tracked sources possessed by the licensee against that licensee's data in the National Source Tracking System. The reconciliation must be conducted during the month of January in each year. The reconciliation process must include resolving any discrepancies between the National Source Tracking System and the actual inventory by filing the reports identified by subsections (1) through (5) of this section. By January 31, of each year, each licensee must submit to the National Source Tracking System confirmation that the data in the National Source Tracking System is correct.

~~((8) Each licensee that possesses Category 1 or 2 nationally tracked sources shall report its initial inventory of Category 1 or 2 nationally tracked sources to the National Source Tracking System by January 31, 2009. The information may be submitted by using any of the methods identified in subsection (6)(a) through (d) of this section. The initial inventory report shall include the following information:~~

- ~~(a) The name, address, and license number of the reporting licensee;~~
- ~~(b) The name of the individual preparing the report;~~
- ~~(c) The manufacturer, model, and serial number of each nationally tracked source or, if not available, other information to uniquely identify the source;~~
- ~~(d) The radioactive material in the sealed source;~~
- ~~(e) The initial or current source strength in becquerels (curies); and~~
- ~~(f) The date for which the source strength is reported.)~~

AMENDATORY SECTION (Amending WSR 16-13-054, filed 6/10/16, effective 7/11/16)

WAC 246-221-240 Reports of stolen, lost or missing radiation sources. (1) Each licensee and registrant shall report by telephone (206-682-5327) and confirm promptly by letter, facsimile, or email to the State Department of Health, Office of Radiation Protection, P.O. Box 47827, Olympia, Washington 98504-7827.

(a) Immediately after its occurrence becomes known to the licensee, stolen, lost, or missing radioactive material in an aggregate

quantity equal to or greater than (~~one thousand~~) 1,000 times the quantity specified in WAC 246-221-300, Appendix B; or

(b) Within (~~thirty~~) 30 days after its occurrence becomes known to the licensee, lost, stolen, or missing radioactive material in an aggregate quantity greater than (~~ten~~) 10 times the quantity specified in WAC 246-221-300, Appendix B that is still missing or any item not exempted in chapter 246-232 WAC; or

(c) Immediately after its occurrence becomes known to the registrant, a stolen, lost, or missing radiation machine.

(2) Each licensee or registrant required to make a report pursuant to subsection (1) of this section shall, within (~~thirty~~) 30 days after making the telephone report, make a written report to the department setting forth the following information:

(a) A description of the licensed or registered source of radiation involved, including, for radioactive material, the kind, quantity, and chemical and physical form; and, for radiation machines, the manufacturer, model and serial number, type and maximum energy of radiation emitted; and

(b) A description of the circumstances under which the loss or theft occurred; and

(c) A statement of disposition, or probable disposition, of the licensed or registered source of radiation involved; and

(d) Exposures of individuals to radiation, circumstances under which the exposures occurred, and the possible total effective dose equivalent to persons in unrestricted areas; and

(e) Actions that have been taken, or will be taken, to recover the source of radiation; and

(f) Procedures or measures that have been, or will be, adopted to ensure against a recurrence of the loss or theft of licensed or registered sources of radiation.

(3) Subsequent to filing the written report, the licensee or registrant shall also report additional substantive information on the loss or theft within (~~thirty~~) 30 days after the licensee or registrant learns of such information.

(4) The licensee or registrant shall prepare any report filed with the department pursuant to this section so that names of individuals who may have received exposure to radiation are stated in a separate and detachable portion of the report.

AMENDATORY SECTION (Amending WSR 16-13-054, filed 6/10/16, effective 7/11/16)

WAC 246-221-250 Notification of incidents. (1) **Immediate notification.** Notwithstanding other requirements for notification, each licensee and registrant shall immediately (as soon as possible but no later than four hours after discovery of an incident) notify the State Department of Health, Office of Radiation Protection, P.O. Box 47827, Olympia, Washington 98504-7827, by telephone (206-682-5327) and confirming letter, facsimile, or email with a follow-up written report within (~~thirty~~) 30 days of any incident involving any radiation source which may have caused or threatens to cause:

(a) An individual to receive:

(i) A total effective dose equivalent of 0.25 Sv (25 rem) or more;

(ii) A lens dose equivalent of 0.75 Sv (75 rem) or more; or
(iii) A shallow dose equivalent to the skin or extremities or a total organ dose equivalent of 2.5 Sv (250 rem) or more;

(b) The release of radioactive material, inside or outside of a restricted area, so that, had an individual been present for (~~twenty-four~~) 24 hours, the individual could have received an intake five times the occupational ALI. This provision does not apply to locations where personnel are not normally stationed during routine operations, such as hot-cells or process enclosures; or

(c) The loss of ability to take immediate protective actions necessary to avoid exposure to sources of radiation or releases of radioactive material that could exceed regulatory limits. Events which could cause such a loss of ability include fires, explosions, toxic gas releases, etc.

(2) **Twenty-four hour notification.** Each licensee and registrant shall within (~~twenty-four~~) 24 hours of discovery of the event, notify the State Department of Health, Office of Radiation Protection, P.O. Box 47827, Olympia, Washington 98504-7827, by telephone (206-682-5327) and confirming letter, facsimile, or email with a follow-up written report within (~~thirty~~) 30 days of any incident involving any radiation source possessed which may have caused or threatens to cause:

(a) An individual to receive, in a period of (~~twenty-four~~) 24 hours:

(i) A total effective dose equivalent exceeding 0.05 Sv (~~(5)~~) five rem);

(ii) A lens dose equivalent exceeding 0.15 Sv (15 rem); or

(iii) A shallow dose equivalent to the skin or extremities or a total organ dose equivalent exceeding 0.5 Sv (50 rem);

(b) The release of radioactive material, inside or outside of a restricted area, so that, had an individual been present for (~~twenty-four~~) 24 hours, the individual could have received an intake in excess of one occupational ALI. This provision does not apply to locations where personnel are not normally stationed during routine operations, such as hot-cells or process enclosures;

(c) An unplanned contamination incident that:

(i) Requires access to the contaminated area, by workers or the general public, to be restricted for more than (~~twenty-four~~) 24 hours by imposing additional radiological controls or by prohibiting entry into the area;

(ii) Involves a quantity of material greater than five times the lowest annual limit on intake specified in WAC 246-221-290; and

(iii) Has access to the area restricted for a reason other than to allow radionuclides with a half-life of less than (~~twenty-four~~) 24 hours to decay prior to decontamination;

(d) Equipment failure or inability to function as designed when:

(i) The equipment is required by regulation or license condition to prevent releases exceeding regulatory limits, to prevent exposures to radiation and radioactive material exceeding regulatory limits or to mitigate the consequences of an accident;

(ii) The equipment is required to be available and operable at the time it becomes disabled or fails to function; and

(iii) No redundant equipment is available and operable to perform the required safety functions;

(e) An unplanned medical treatment at a medical facility of an individual with removable radioactive contamination on the individual's clothing or body; or

(f) An unplanned fire or explosion damaging any radioactive material or any device, container or equipment containing radioactive material when:

(i) The quantity of radioactive material involved is greater than five times the lowest annual limit on intake specified in WAC 246-221-290; and

(ii) The damage affects the integrity of the radioactive material or its container.

(3) For each occurrence requiring notification pursuant to this section, a prompt investigation of the situation shall be initiated by the licensee/registrant. A written report of the findings of the investigation shall be sent to the department within (~~(thirty)~~) 30 days.

(4) The licensee or registrant shall prepare each report filed with the department under this section so that names of individuals who have received exposure to sources of radiation are stated in a separate and detachable portion of the report.

Any report filed with the department under this section shall contain the information described in WAC 246-221-260 (2) and (3).

(5) The provisions of this section do not apply to doses that result from planned special exposures, provided such doses are within the limits for planned special exposures and are reported pursuant to WAC 246-221-265.

(6) Telephone notifications that do not involve immediate or (~~(twenty-four)~~) 24 hour notification should be made to the Tumwater office (360-236-3300).

(7) Telephone notification required under this section shall include, to the extent that the information is available at the time of notification:

(a) The caller's name and call-back telephone number;

(b) A description of the incident including date and time;

(c) The exact location of the incident;

(d) The radionuclides, quantities, and chemical and physical forms of the radioactive materials involved; and

(e) Any personnel radiation exposure data available.

AMENDATORY SECTION (Amending WSR 99-15-105, filed 7/21/99, effective 8/21/99)

WAC 246-221-260 Reports of overexposures and excessive levels and concentrations. (1) In addition to any notification required by WAC 246-221-250, each licensee or registrant shall submit a written report to the department within (~~(thirty)~~) 30 days after learning of any of the following occurrences:

(a) Incidents for which notification is required by WAC 246-221-250; or

(b) Doses in excess of any of the following:

(i) The occupational dose limits for adults in WAC 246-221-010;
or

(ii) The occupational dose limits for a minor in WAC 246-221-050;
or

(iii) The limits for an embryo/fetus of a declared pregnant woman in WAC 246-221-055; or

(iv) The limits for an individual member of the public in WAC 246-221-060; or

(v) Any applicable limit in the license; or
(vi) The ALARA constraints for air emissions established under WAC 246-221-005; or

(c) Levels of radiation or concentrations of radioactive material in:

(i) A restricted area in excess of applicable limits in the license; or

(ii) An unrestricted area in excess of (~~ten~~) 10 times the applicable limit set forth in this chapter or in the license or registration, whether or not involving exposure of any individual in excess of the limits in WAC 246-221-060; or

(d) For source materials milling licensees and nuclear power plants subject to the provisions of United States Environmental Protection Agency's generally applicable environmental radiation standards in 40 C.F.R. 190, levels of radiation or releases of radioactive material in excess of those standards, or of license conditions related to those standards.

(2) Each report required by subsection (1) of this section shall describe:

(a) The incident and its exact location, time and date;

(b) The extent of exposure of individuals to radiation or to radioactive material, including estimates of each individual's dose as required by subsection (3) of this section;

(c) Levels of radiation and concentrations of radioactive material involved, including the radionuclides, quantities, and chemical and physical form;

(d) The cause or probable cause of the exposure, levels of radiation or concentrations;

(e) The manufacturer and model number (if applicable) of any equipment that failed or malfunctioned;

(f) The results of any evaluations or assessments; and

(g) Corrective steps taken or planned to assure against a recurrence, including the schedule for achieving conformance with applicable limits, ALARA constraints, generally applicable environmental standards, and associated license conditions.

(3) Each report filed with the department pursuant to this section shall include for each individual exposed the name, Social Security number, and date of birth, and an estimate of the individual's dose. With respect to the limit for the embryo/fetus in WAC 246-221-055, the identifiers should be those of the declared pregnant woman. The report shall be prepared so that this information is stated in a separate and detachable part of the report.

(4) Individuals shall be notified of reports in accordance with the requirements of WAC 246-222-040.

AMENDATORY SECTION (Amending WSR 99-05-013, filed 2/5/99, effective 3/8/99)

WAC 246-221-265 Special reports to the department—Planned special exposures and leaking sources. (1) The licensee or registrant shall submit a written report to the department within (~~thirty~~) 30 days following any planned special exposure conducted in accordance with WAC 246-221-030. The written report shall:

- (a) Inform the department that a planned special exposure was conducted;
 - (b) Indicate the date the planned special exposure occurred; and
 - (c) Provide the information required by WAC 246-221-030.
- (2) The licensee shall file a written report with the department within five days after learning that a sealed source is leaking or contaminated. The report shall describe:
- (a) The source;
 - (b) The source holder;
 - (c) The equipment in which the source is installed;
 - (d) The test results; and
 - (e) The corrective action taken.

AMENDATORY SECTION (Amending WSR 00-07-085, filed 3/15/00, effective 4/15/00)

WAC 246-221-270 Vacating premises and release of equipment. (1)

Each specific licensee shall notify the department in writing of intent to vacate, at least (~~thirty~~) 30 days before vacating or relinquishing possession or control of premises which may have been contaminated with radioactive material as a result of licensed activities.

(2) Each licensee shall permanently decontaminate the premise, before vacating any premise or transferring the premise, in accordance with the standards specified in chapter 246-246 WAC. A survey by the licensee shall be made after the decontamination and the department and the landlord or subsequent tenant or transferee shall be provided with a copy of the survey no later than the date of vacating or relinquishing possession or control of the premise.

(3) No machinery, instruments, laboratory equipment or any other property used in contact with, or close proximity to radioactive material at a licensed premise shall be assigned, sold, leased, or transferred to an unlicensed person unless the property has been decontaminated and meets the standards specified in WAC 246-232-140. A survey shall be made after the decontamination and the department and subsequent owner or transferee shall be provided with a copy of the survey report.

AMENDATORY SECTION (Amending WSR 01-05-110, filed 2/21/01, effective 3/24/01)

WAC 246-221-285 Assigned protection factors for respirators^a.

	Operating mode	Assigned Protection Factors
I. Air-Purifying Respirators (Particulate ^b only) ^c :		
Filtering facepiece disposable ^d	Negative Pressure	(^d)
Facepiece, half ^e	Negative Pressure	10
Facepiece, full	Negative Pressure	100
Facepiece, half	Powered air-purifying respirators	50

	Operating mode	Assigned Protection Factors
Facepiece, full	Powered air-purifying respirators	1000
Helmet/hood	Powered air-purifying respirators	1000
Facepiece, loose-fitting	Powered air-purifying respirators	25
II. Atmosphere-Supplying Respirators (Particulate, gases and vapors ^f):		
1. Air-line respirator:		
Facepiece, half	Demand	10
Facepiece, half	Continuous Flow	50
Facepiece, half	Pressure Demand	50
Facepiece, full	Demand	100
Facepiece, full	Continuous Flow	1000
Facepiece, full	Pressure Demand	1000
Helmet/hood	Continuous Flow	1000
Facepiece, loose-fitting	Continuous Flow	25
Suit	Continuous Flow	(g)
2. Self-contained breathing apparatus (SCBA):		
Facepiece, full	Demand	h100
Facepiece, full	Pressure Demand	i10,000
Facepiece, full	Demand, Recirculating	h100
Facepiece, full	Positive Pressure Recirculating	i10,000

III. Combination Respirators:

Any combination of air-purifying and atmosphere-supplying respirators. Assigned protection factor for type and mode of operation as listed above.

- a These assigned protection factors apply only in a respiratory protection program that meets the requirements of this chapter. They are applicable only to airborne radiological hazards and may not be appropriate to circumstances when chemical or other respiratory hazards exist instead of, or in addition to, radioactive hazards. Selection and use of respirators for these circumstances must also comply with Department of Labor regulations.
Radioactive contaminants for which the concentration values in Table 1, Column 3 of WAC 246-221-290, Appendix A, are based on internal dose due to inhalation may, in addition, present external exposure hazards at higher concentrations. Under these circumstances, limitations on occupancy may have to be governed by external dose limits.
- b Air-purifying respirators with APF <100 must be equipped with particulate filters that are at least 95 percent efficient. Air-purifying respirators with APF = 100 must be equipped with particulate filters that are at least 99 percent efficient. Air-purifying respirators with APFs >100 must be equipped with particulate filters that are at least 99.97 percent efficient.
- c The licensee may apply to the department for the use of an APF greater than ((+)) one for sorbent cartridges as protection against airborne radioactive gases and vapors (e.g., radioiodine).
- d Licensees may permit individuals to use this type of respirator who have not been medically screened or fit tested on the device provided that no credit be taken for their use in estimating intake or dose. It is also recognized that it is difficult to perform an effective positive or negative pressure preuse user seal check on this type of device. All other respiratory protection program requirements listed in WAC 246-221-117 apply. An assigned protection factor has not been assigned for these devices. However, an APF equal to 10 may be used if the licensee can demonstrate a fit factor of at least 100 by use of a validated or evaluated, qualitative or quantitative fit test.
- e Under-chin type only. No distinction is made in this section between elastomeric half-masks with replaceable cartridges and those designed with the filter medium as an integral part of the facepiece (e.g., disposable or reusable disposable). Both types are acceptable so long as the seal area of the latter contains some substantial type of seal-enhancing material such as rubber or plastic, the two or more suspension straps are adjustable, the filter medium is at least 95 percent efficient and all other requirements of this part are met.
- f The assigned protection factors for gases and vapors are not applicable to radioactive contaminants that present an absorption or submersion hazard. For tritium oxide vapor, approximately ((~~one-third~~)) 1/3 of the intake occurs by absorption through the skin so that an overall protection factor of ((3)) three is appropriate when atmosphere-supplying respirators are used to protect against tritium oxide. Exposure to radioactive noble gases is not considered a significant respiratory hazard, and protective actions for these contaminants should be based on external (submersion) dose considerations.
- g No NIOSH approval schedule is currently available for atmosphere-supplying suits. This equipment may be used in an acceptable respiratory protection program as long as all the other minimum program requirements, with the exception of fit testing, are met (i.e., WAC 246-221-117).
- h The licensee should implement institutional controls to assure that these devices are not used in areas immediately dangerous to life or health (IDLH).
- i This type of respirator may be used as an emergency device in unknown concentrations for protection against inhalation hazards. External radiation hazards and other limitations to permitted exposure such as skin absorption shall be taken into account in these circumstances. This device may not be used by any individual who experiences perceptible outward leakage of breathing gas while wearing the device.

WAC 246-221-290 Appendix A—Annual limits on intake (ALI) and derived air concentrations (DAC) of radionuclides for occupational exposure; effluent concentrations; concentrations for release to sanitary sewerage. For each radionuclide, Table I indicates the chemical form which is to be used for selecting the appropriate ALI or DAC value. The ALIs and DACs for inhalation are given for an aerosol with an activity median aerodynamic diameter (AMAD) of ((±)) one µm (micron) and for three classes (D,W,Y) of radioactive material, which refer to their retention (approximately days, weeks or years) in the pulmonary region of the lung. This classification applies to a range of clearance half-times for D if less than ((~~ten~~)) 10 days, for W from ((~~ten to one hundred~~)) 10 to 100 days, and for Y greater than ((~~one hundred~~)) 100 days. Table II provides concentration limits for airborne and liquid effluents released to the general environment. Table III provides concentration limits for discharges to sanitary sewerage.

Note: The values in Tables I, II, and III are presented in the computer "E" notation. In this notation a value of 6E-02 represents a value of 6×10^{-2} or 0.06, 6E+2 represents 6×10^2 or 600, and 6E+0 represents 6×10^0 or 6.

Table I "Occupational Values"

Note that the columns in Table I of this appendix captioned "Oral Ingestion ALI," "Inhalation ALI," and "DAC," are applicable to occupational exposure to radioactive material.

The ALIs in this appendix are the annual intakes of given radionuclide by "Reference Man" which would result in either: A committed effective dose equivalent of 0.05 Sv ((~~5~~)) five rem), stochastic ALI; or a committed dose equivalent of 0.5 Sv (50 rem) to an organ or tissue, nonstochastic ALI. The stochastic ALIs were derived to result in a risk, due to irradiation of organs and tissues, comparable to the risk associated with deep dose equivalent to the whole body of 0.05 Sv ((~~5~~)) five rem). The derivation includes multiplying the committed dose equivalent to an organ or tissue by a weighting factor, w_T . This weighting factor is the proportion of the risk of stochastic effects resulting from irradiation of the organ or tissue, T, to the total risk of stochastic effects when the whole body is irradiated uniformly. The values of w_T are listed under the definition of weighting factor in WAC 246-221-005. The nonstochastic ALIs were derived to avoid nonstochastic effects, such as prompt damage to tissue or reduction in organ function.

A value of $w_T = 0.06$ is applicable to each of the five organs or tissues in the "remainder" category receiving the highest dose equivalents, and the dose equivalents of all other remaining tissues may be disregarded. The following portions of the GI tract — stomach, small intestine, upper large intestine, and lower large intestine — are to be treated as four separate organs.

Note that the dose equivalents for an extremity, elbows, arms below the elbows, feet and lower legs, knees, and legs below the knees, skin, and lens of the eye are not considered in computing the committed effective dose equivalent, but are subject to limits that must be met separately.

When an ALI is defined by the stochastic dose limit, this value alone is given. When an ALI is determined by the non-stochastic dose limit to an organ, the organ or tissue to which the limit applies is shown, and the ALI for the stochastic limit is shown in parentheses. Abbreviated organ or tissue designations are used:

LLI wall = lower large intestine wall;
St. wall = stomach wall;
Blad wall = bladder wall; and
Bone surf = bone surface.

The use of the ALIs listed first, the more limiting of the stochastic and nonstochastic ALIs, will ensure that nonstochastic effects are avoided and that the risk of stochastic effects is limited to an acceptably low value. If, in a particular situation involving a radionuclide for which the nonstochastic ALI is limiting, use of that nonstochastic ALI is considered unduly conservative, the licensee may use the stochastic ALI to determine the committed effective dose equivalent. However, the licensee shall also ensure that the 0.5 Sv (50 rem) dose equivalent limit for any organ or tissue is not exceeded by the sum of the external deep dose equivalent plus the internal committed dose equivalent to that organ, not the effective dose. For the case where there is no external dose contribution, this would be demonstrated if the sum of the fractions of the nonstochastic ALIs (ALI_{ns}) that contribute to the committed dose equivalent to the organ receiving the highest dose does not exceed unity, that is, $\sum (\text{intake (in } \mu\text{Ci) of each radionuclide} / ALI_{ns}) \leq 1.0$. If there is an external deep dose equivalent contribution of H_d , then this sum must be less than $(\pm) \text{one} - (H_d/50)$, instead of ≤ 1.0 .

The derived air concentration (DAC) values are derived limits intended to control chronic occupational exposures. The relationship between the DAC and the ALI is given by:

$$DAC = ALI (\text{in } \mu\text{Ci}) / (2000 \text{ hours per working year} \times 60 \text{ minutes/hour} \times 2 \times 10^4 \text{ ml per minute}) = [ALI / 2.4 \times 10^9] \mu\text{Ci/ml}$$

where 2×10^4 ml per minute is the volume of air breathed per minute at work by Reference Man under working conditions of light work.

The DAC values relate to one of two modes of exposure: Either external submersion or the internal committed dose equivalents resulting from inhalation of radioactive materials. DACs based upon submersion are for immersion in a semi-infinite cloud of uniform concentration and apply to each radionuclide separately.

The ALI and DAC values include contributions to exposure by the single radionuclide named and any in-growth of daughter radionuclides produced in the body by decay of the parent. However, intakes that include both the parent and daughter radionuclides should be treated by the general method appropriate for mixtures.

The values of ALI and DAC do not apply directly when the individual both ingests and inhales a radionuclide, when the individual is exposed to a mixture of radionuclides by either inhalation or ingestion or both, or when the individual is exposed to both internal and external irradiation. See WAC 246-221-015. When an individual is exposed to radioactive materials which fall under several of the translocation classifications of the same radionuclide, such as, Class D,

Class W, or Class Y, the exposure may be evaluated as if it were a mixture of different radionuclides.

It should be noted that the classification of a compound as Class D, W, or Y is based on the chemical form of the compound and does not take into account the radiological half-life of different radionuclides. For this reason, values are given for Class D, W, and Y compounds, even for very short-lived radionuclides.

Table II "Effluent Concentrations"

The columns in Table II of this appendix captioned "Effluents," "Air" and "Water" are applicable to the assessment and control of dose to the public, particularly in the implementation of the provisions of WAC 246-221-070. The concentration values given in Columns 1 and 2 of Table II are equivalent to the radionuclide concentrations which, if inhaled or ingested continuously over the course of a year, would produce a total effective dose equivalent of 0.50 mSv (0.05 rem).

Consideration of nonstochastic limits has not been included in deriving the air and water effluent concentration limits because non-stochastic effects are presumed not to occur at or below the dose levels established for individual members of the public. For radionuclides, where the nonstochastic limit was governing in deriving the occupational DAC, the stochastic ALI was used in deriving the corresponding airborne effluent limit in Table II. For this reason, the DAC and airborne effluent limits are not always proportional as was the case in the previous Appendix A of this chapter.

The air concentration values listed in Table II, Column 1 were derived by one of two methods. For those radionuclides for which the stochastic limit is governing, the occupational stochastic inhalation ALI was divided by 2.4×10^9 , relating the inhalation ALI to the DAC, as explained above, and then divided by a factor of ~~((three hundred))~~ 300. The factor of ~~((three hundred))~~ 300 includes the following components: A factor of ~~((fifty))~~ 50 to relate the 0.05 Sv ~~((5))~~ five rem annual occupational dose limit to the ~~((1))~~ one mSv (0.1 rem) limit for members of the public, a factor of three to adjust for the difference in exposure time and the inhalation rate for a worker and that for members of the public; and a factor of two to adjust the occupational values, derived for adults, so that they are applicable to other age groups.

For those radionuclides for which submersion, that is external dose, is limiting, the occupational DAC in Table I, Column 3 was divided by ~~((two hundred nineteen))~~ 219. The factor of ~~((two hundred nineteen))~~ 219 is composed of a factor of ~~((fifty))~~ 50, as described above, and a factor of 4.38 relating occupational exposure for ~~((two thousand))~~ 2,000 hours per year to full-time exposure ~~((eight thousand seven hundred sixty))~~ 8,760 hours per year). Note that an additional factor of two for age considerations is not warranted in the submersion case.

The water concentrations were derived by taking the most restrictive occupational stochastic oral ingestion ALI and dividing by 7.3×10^7 . The factor of 7.3×10^7 (ml) includes the following components: The factors of ~~((fifty))~~ 50 and two described above and a factor of 7.3×10^5 (ml) which is the annual water intake of Reference Man.

Note 2 of this appendix provides groupings of radionuclides which are applicable to unknown mixtures of radionuclides. These groupings, including occupational inhalation ALIs and DACs, air and water effluent concentrations and releases to sewer, require demonstrating that the most limiting radionuclides in successive classes are absent. The limit for the unknown mixture is defined when the presence of one of the listed radionuclides cannot be definitely excluded as being present either from knowledge of the radionuclide composition of the source or from actual measurements.

Table III "Releases to Sewers"

The monthly average concentrations for release to sanitary sewerage are applicable to the provisions in WAC 246-221-190. The concentration values were derived by taking the most restrictive occupational stochastic oral ingestion ALI and dividing by 7.3×10^6 (ml). The factor of 7.3×10^6 (ml) is composed of a factor of 7.3×10^5 (ml), the annual water intake by Reference Man, and a factor of ~~((ten))~~ 10, such that the concentrations, if the sewage released by the licensee were the only source of water ingested by a Reference Man during a year, would result in a committed effective dose equivalent of ~~((5))~~ five mSv (0.5 rem).

LIST OF ELEMENTS

Name	Symbol	Atomic Number	Name	Symbol	Atomic Number
Actinium	Ac	89	Molybdenum	Mo	42
Aluminum	Al	13	Neodymium	Nd	60
Americium	Am	95	Neptunium	Np	93
Antimony	Sb	51	Nickel	Ni	28
Argon	Ar	18	Nitrogen	N	7
Arsenic	As	33	Niobium	Nb	41
Astatine	At	85	Osmium	Os	76
Barium	Ba	56	Oxygen	O	8
Berkelium	Bk	97	Palladium	Pd	46
Beryllium	Be	4	Phosphorus	P	15
Bismuth	Bi	83	Platinum	Pt	78
Bromine	Br	35	Plutonium	Pu	94
Cadmium	Cd	48	Polonium	Po	84
Calcium	Ca	20	Potassium	K	19
Californium	Cf	98	Praseodymium	Pr	59
Carbon	C	6	Promethium	Pm	61
Cerium	Ce	58	Protactinium	Pa	91
Cesium	Cs	55	Radium	Ra	88
Chlorine	Cl	17	Radon	Rn	86
Chromium	Cr	24	Rhenium	Re	75
Cobalt	Co	27	Rhodium	Rh	45
Copper	Cu	29	Rubidium	Rb	37
Curium	Cm	96	Ruthenium	Ru	44
Dysprosium	Dy	66	Samarium	Sm	62
Einsteinium	Es	99	Scandium	Sc	21
Erbium	Er	68	Selenium	Se	34
Europium	Eu	63	Silicon	Si	14

LIST OF ELEMENTS

Name	Symbol	Atomic Number	Name	Symbol	Atomic Number
Fermium	Fm	100	Silver	Ag	47
Fluorine	F	9	Sodium	Na	11
Francium	Fr	87	Strontium	Sr	38
Gadolinium	Gd	64	Sulfur	S	16
Gallium	Ga	31	Tantalum	Ta	73
Germanium	Ge	32	Technetium	Tc	43
Gold	Au	79	Tellurium	Te	52
Hafnium	Hf	72	Terbium	Tb	65
Holmium	Ho	67	Thallium	Tl	81
Hydrogen	H	1	Thorium	Th	90
Indium	In	49	Thulium	Tm	69
Iodine	I	53	Tin	Sn	50
Iridium	Ir	77	Titanium	Ti	22
Iron	Fe	26	Tungsten	W	74
Krypton	Kr	36	Uranium	U	92
Lanthanum	La	57	Vanadium	V	23
Lead	Pb	82	Xenon	Xe	54
Lutetium	Lu	71	Ytterbium	Yb	70
Magnesium	Mg	12	Yttrium	Y	39
Manganese	Mn	25	Zinc	Zn	30
Mendelevium	Md	101	Zirconium	Zr	40
Mercury	Hg	80			

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion ALI μCi	Inhalation ALI μCi DAC μCi/ml		Air μCi/ml	Water μCi/ml	
1	Hydrogen-3	Water, DAC includes skin absorption	8E+4	8E+4	2E-5	1E-7	1E-3	1E-2
	Gas (HT or T ₂) Submersion ¹ : Use above values as HT and T ₂ oxidize in air and in the body to HTO.							
4	Beryllium-7	W, all compounds except those given for Y	4E+4	2E+4	9E-6	3E-8	6E-4	6E-3
		Y, oxides, halides, and nitrates	-	2E+4	8E-6	3E-8	-	-
4	Beryllium-10	W, see ⁷ Be	1E+3	2E+2	6E-8	2E-10	-	-
		LLI wall (1E+3)	-	-	-	-	2E-5	2E-4
		Y, see ⁷ Be	-	1E+1	6E-9	2E-11	-	-
6	Carbon-11 ²	Monoxide	-	1E+6	5E-4	2E-6	-	-
		Dioxide	-	6E+5	3E-4	9E-7	-	-
		Compounds	4E+5	4E+5	2E-4	6E-7	6E-3	6E-2
6	Carbon-14	Monoxide	-	2E+6	7E-4	2E-6	-	-
		Dioxide	-	2E+5	9E-5	3E-7	-	-
		Compounds	2E+3	2E+3	1E-6	3E-9	3E-5	3E-4
7	Nitrogen-13 ²	Submersion ¹	-	-	4E-6	2E-8	-	-
8	Oxygen-15 ²	Submersion ¹	-	-	4E-6	2E-8	-	-
9	Fluorine-18 ²	D, fluorides of H, Li, Na, K, Rb, Cs, and Fr	5E+4	7E+4	3E-5	1E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
			St wall (5E+4)	-	-	-	7E-4	7E-3
		W, fluorides of Be, Mg, Ca, Sr, Ba, Ra, Al, Ga, In, Tl, As, Sb, Bi, Fe, Ru, Os, Co, Ni, Pd, Pt, Cu, Ag, Au, Zn, Cd, Hg, Sc, Y, Ti, Zr, V, Nb, Ta, Mn, Tc, and Re	-	9E+4	4E-5	1E-7	-	-
		Y, lanthanum fluoride	-	8E+4	3E-5	1E-7	-	-
11	Sodium-22	D, all compounds	4E+2	6E+2	3E-7	9E-10	6E-6	6E-5
11	Sodium-24	D, all compounds	4E+3	5E+3	2E-6	7E-9	5E-5	5E-4
12	Magnesium-28	D, all compounds except those given for W	7E+2	2E+3	7E-7	2E-9	9E-6	9E-5
		W, oxides, hydroxides, carbides, halides, and nitrates	-	1E+3	5E-7	2E-9	-	-
13	Aluminum-26	D, all compounds except those given for W	4E+2	6E+1	3E-8	9E-11	6E-6	6E-5
		W, oxides, hydroxides, carbides, halides, and nitrates	-	9E+1	4E-8	1E-10	-	-
14	Silicon-31	D, all compounds except those given for W and Y	9E+3	3E+4	1E-5	4E-8	1E-4	1E-3
		W, oxides, hydroxides, carbides, and nitrates	-	3E+4	1E-5	5E-8	-	-
		Y, aluminosilicate glass	-	3E+4	1E-5	4E-8	-	-
14	Silicon-32	D, see ³¹ Si	2E+3	2E+2	1E-7	3E-10	-	-
			LLI wall (3E+3)	-	-	-	4E-5	4E-4
		W, see ³¹ Si	-	1E+2	5E-8	2E-10	-	-
		Y, see ³¹ Si	-	5E+0	2E-9	7E-12	-	-
15	Phosphorus-32	D, all compounds except phosphates given for W	6E+2	9E+2	4E-7	1E-9	9E-6	9E-5
		W, phosphates of Zn ²⁺ , S ³⁺ , Mg ²⁺ , Fe ³⁺ , Bi ³⁺ , and lanthanides	-	4E+2	2E-7	5E-10	-	-
15	Phosphorus-33	D, see ³² P	6E+3	8E+3	4E-6	1E-8	8E-5	8E-4
		W, see ³² P	-	3E+3	1E-6	4E-9	-	-
16	Sulfur-35	Vapor	-	1E+4	6E-6	2E-8	-	-
		D, sulfides and sulfates except those given for W	1E+4	2E+4	7E-6	2E-8	-	-
			LLI wall (8E+3)	-	-	-	1E-4	1E-3
		W, elemental sulfur, sulfides of Sr, Ba, Ge, Sn, Pb, As, Sb, Bi, Cu, Ag, Au, Zn, Cd, Hg, W, and Mo. Sulfates of Ca, Sr, Ba, Ra, As, Sb, and Bi	6E+3	-	-	-	-	-
			-	2E+3	9E-7	3E-9	-	-
17	Chlorine-36	D, chlorides of H, Li, Na, K, Rb, Cs, and Fr	2E+3	2E+3	1E-6	3E-9	2E-5	2E-4
		W, chlorides of lanthanides, Be, Mg, Ca, Sr, Ba, Ra, Al, Ga, In, Tl, Ge, Sn, Pb, As, Sb, Bi, Fe, Ru, Os, Co, Rh, Ir, Ni, Pd, Pt, Cu, Ag, Au, Zn, Cd, Hg, Sc, Y, Ti, Zr, Hf, V, Nb, Ta, Cr, Mo, W, Mn, Tc, and Re	-	2E+2	1E-7	3E-10	-	-
17	Chlorine-38 ²	D, see ³⁶ Cl	2E+4	4E+4	2E-5	6E-8	-	-
			St wall (3E+4)	-	-	-	3E-4	3E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
17	Chlorine-39 ²	W, see ³⁶ Cl	-	5E+4	2E-5	6E-8	-	-
		D, see ³⁶ Cl	2E+4	5E+4	2E-5	7E-8	-	-
18	Argon-37	St wall (4E+4)	-	-	-	-	5E-4	5E-3
		W, see ³⁶ Cl	-	6E+4	2E-5	8E-8	-	-
18	Argon-39	Submersion ¹	-	-	1E+0	6E-3	-	-
18	Argon-41	Submersion ¹	-	-	2E-4	8E-7	-	-
19	Potassium-40	Submersion ¹	-	-	3E-6	1E-8	-	-
19	Potassium-42	D, all compounds	3E+2	4E+2	2E-7	6E-10	4E-6	4E-5
19	Potassium-43	D, all compounds	5E+3	5E+3	2E-6	7E-9	6E-5	6E-4
19	Potassium-44 ²	D, all compounds	6E+3	9E+3	4E-6	1E-8	9E-5	9E-4
19	Potassium-45 ²	D, all compounds	2E+4	7E+4	3E-5	9E-8	-	-
		St wall (4E+4)	-	-	-	-	5E-4	5E-3
20	Calcium-41	D, all compounds	3E+4	1E+5	5E-5	2E-7	-	-
		St wall (5E+4)	-	-	-	-	7E-4	7E-3
20	Calcium-45	W, all compounds	3E+3	4E+3	2E-6	-	-	-
20	Calcium-47	W, all compounds	Bone surf (4E+3)	Bone surf (4E+3)	-	5E-9	6E-5	6E-4
21	Scandium-43	Y, all compounds	2E+3	8E+2	4E-7	1E-9	2E-5	2E-4
21	Scandium-44m	Y, all compounds	8E+2	9E+2	4E-7	1E-9	1E-5	1E-4
21	Scandium-44	Y, all compounds	7E+3	2E+4	9E-6	3E-8	1E-4	1E-3
21	Scandium-46	Y, all compounds	5E+2	7E+2	3E-7	1E-9	7E-6	7E-5
21	Scandium-47	Y, all compounds	4E+3	1E+4	5E-6	2E-8	5E-5	5E-4
21	Scandium-48	Y, all compounds	9E+2	2E+2	1E-7	3E-10	1E-5	1E-4
		LLI wall (3E+3)	2E+3	3E+3	1E-6	4E-9	-	-
21	Scandium-49 ²	Y, all compounds	8E+2	1E+3	6E-7	2E-9	1E-5	1E-4
22	Titanium-44	D, all compounds except those given for W and Y	2E+4	5E+4	2E-5	8E-8	3E-4	3E-3
		W, oxides, hydroxides, carbides, halides, and nitrates	3E+2	1E+1	5E-9	2E-11	4E-6	4E-5
22	Titanium-45	Y, SrTiO	-	3E+1	1E-8	4E-11	-	-
		D, see ⁴⁴ Ti	-	6E+0	2E-9	8E-12	-	-
23	Vanadium-472	W, see ⁴⁴ Ti	9E+3	3E+4	1E-5	3E-8	1E-4	1E-3
		Y, see ⁴⁴ Ti	-	3E+4	1E-5	4E-8	-	-
23	Vanadium-48	D, all compounds except those given for W	3E+4	8E+4	3E-5	1E-7	-	-
		St wall (3E+4)	-	-	-	-	4E-4	4E-3
23	Vanadium-49	W, oxides, hydroxides, carbides, and halides	-	1E+5	4E-5	1E-7	-	-
		D, see ⁴⁷ V	6E+2	1E+3	5E-7	2E-9	9E-6	9E-5
23	Vanadium-49	W, see ⁴⁷ V	-	6E+2	3E-7	9E-10	-	-
		D, see ⁴⁷ V	7E+4	3E+4	1E-5	-	-	-
		LLI wall (9E+4)	-	Bone surf (3E+4)	-	5E-8	1E-3	1E-2
		W, see ⁴⁷ V	-	2E+4	8E-6	2E-8	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
24	Chromium-48	D, all compounds except those given for W and Y	6E+3	1E+4	5E-6	2E-8	8E-5	8E-4
		W, halides and nitrates	-	7E+3	3E-6	1E-8	-	-
		Y, oxides and hydroxides	-	7E+3	3E-6	1E-8	-	-
24	Chromium-49 ²	D, see ⁴⁸ Cr	3E+4	8E+4	4E-5	1E-7	4E-4	4E-3
		W, see ⁴⁸ Cr	-	1E+5	4E-5	1E-7	-	-
		Y, see ⁴⁸ Cr	-	9E+4	4E-5	1E-7	-	-
24	Chromium-51	D, see ⁴⁸ Cr	4E+4	5E+4	2E-5	6E-8	5E-4	5E-3
		W, see ⁴⁸ Cr	-	2E+4	1E-5	3E-8	-	-
		Y, see ⁴⁸ Cr	-	2E+4	8E-6	3E-8	-	-
25	Manganese-51 ²	D, all compounds except those given for W	2E+4	5E+4	2E-5	7E-8	3E-4	3E-3
		W, oxides, hydroxides, halides, and nitrates	-	6E+4	3E-5	8E-8	-	-
25	Manganese-52m ²	D, see ⁵¹ Mn	3E+4	9E+4	4E-5	1E-7	-	-
		St wall (4E+4)	-	-	-	-	5E-4	5E-3
		W, see ⁵¹ Mn	-	1E+5	4E-5	1E-7	-	-
25	Manganese-52	D, see ⁵¹ Mn	7E+2	1E+3	5E-7	2E-9	1E-5	1E-4
		W, see ⁵¹ Mn	-	9E+2	4E-7	1E-9	-	-
25	Manganese-53	D, see ⁵¹ Mn	5E+4	1E+4	5E-6	-	7E-4	7E-3
		Bone surf (2E+4)	-	-	-	3E-8	-	-
		W, see ⁵¹ Mn	-	1E+4	5E-6	2E-8	-	-
25	Manganese-54	D, see ⁵¹ Mn	2E+3	9E+2	4E-7	1E-9	3E-5	3E-4
		W, see ⁵¹ Mn	-	8E+2	3E-7	1E-9	-	-
25	Manganese-56	D, see ⁵¹ Mn	5E+3	2E+4	6E-6	2E-8	7E-5	7E-4
		W, see ⁵¹ Mn	-	2E+4	9E-6	3E-8	-	-
26	Iron-52	D, all compounds except those given for W	9E+2	3E+3	1E-6	4E-9	1E-5	1E-4
		W, oxides, hydroxides, and halides	-	2E+3	1E-6	3E-9	-	-
26	Iron-55	D, see ⁵² Fe	9E+3	2E+3	8E-7	3E-9	1E-4	1E-3
		W, see ⁵² Fe	-	4E+3	2E-6	6E-9	-	-
26	Iron-59	D, see ⁵² Fe	8E+2	3E+2	1E-7	5E-10	1E-5	1E-4
		W, see ⁵² Fe	-	5E+2	2E-7	7E-10	-	-
26	Iron-60	D, see ⁵² Fe	3E+1	6E+0	3E-9	9E-12	4E-7	4E-6
		W, see ⁵² Fe	-	2E+1	8E-9	3E-11	-	-
27	Cobalt-55	W, all compounds except those given for Y	1E+3	3E+3	1E-6	4E-9	2E-5	2E-4
		Y, oxides, hydroxides, halides, and nitrates	-	3E+3	1E-6	4E-9	-	-
27	Cobalt-56	W, see ⁵⁵ Co	5E+2	3E+2	1E-7	4E-10	6E-6	6E-5
		Y, see ⁵⁵ Co	4E+2	2E+2	8E-8	3E-10	-	-
27	Cobalt-57	W, see ⁵⁵ Co	8E+3	3E+3	1E-6	4E-9	6E-5	6E-4
		Y, see ⁵⁵ Co	4E+3	7E+2	3E-7	9E-10	-	-
27	Cobalt-58m	W, see ⁵⁵ Co	6E+4	9E+4	4E-5	1E-7	8E-4	8E-3
		Y, see ⁵⁵ Co	-	6E+4	3E-5	9E-8	-	-
27	Cobalt-58	W, see ⁵⁵ Co	2E+3	1E+3	5E-7	2E-9	2E-5	2E-4

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
27	Cobalt-60m ²	Y, see ⁵⁵ Co	1E+3	7E+2	3E-7	1E-9	-	-
		W, see ⁵⁵ Co	1E+6	4E+6	2E-3	6E-6	-	-
		St wall (1E+6)	-	-	-	2E-2	2E-1	
27	Cobalt-60	Y, see ⁵⁵ Co	-	3E+6	1E-3	4E-6	-	-
		W, see ⁵⁵ Co	5E+2	2E+2	7E-8	2E-10	3E-6	3E-5
		Y, see ⁵⁵ Co	2E+2	3E+1	1E-8	5E-11	-	-
27	Cobalt-61 ²	W, see ⁵⁵ Co	2E+4	6E+4	3E-5	9E-8	3E-4	3E-3
		Y, see ⁵⁵ Co	2E+4	6E+4	2E-5	8E-8	-	-
		W, see ⁵⁵ Co	4E+4	2E+5	7E-5	2E-7	-	-
27	Cobalt-62m ²	St wall (5E+4)	-	-	-	7E-4	7E-3	
		Y, see ⁵⁵ Co	-	2E+5	6E-5	2E-7	-	-
		D, all compounds except those given for W	1E+3	2E+3	8E-7	3E-9	2E-5	2E-4
28	Nickel-56	W, oxides, hydroxides, and carbides	-	1E+3	5E-7	2E-9	-	-
		Vapor	-	1E+3	5E-7	2E-9	-	-
		D, see ⁵⁶ Ni	2E+3	5E+3	2E-6	7E-9	2E-5	2E-4
28	Nickel-57	W, see ⁵⁶ Ni	-	3E+3	1E-6	4E-9	-	-
		Vapor	-	6E+3	3E-6	9E-9	-	-
		D, see ⁵⁶ Ni	2E+4	4E+3	2E-6	5E-9	3E-4	3E-3
28	Nickel-59	W, see ⁵⁶ Ni	-	7E+3	3E-6	1E-8	-	-
		Vapor	-	2E+3	8E-7	3E-9	-	-
		D, see ⁵⁶ Ni	9E+3	2E+3	7E-7	2E-9	1E-4	1E-3
28	Nickel-63	W, see ⁵⁶ Ni	-	3E+3	1E-6	4E-9	-	-
		Vapor	-	8E+2	3E-7	1E-9	-	-
		D, see ⁵⁶ Ni	8E+3	2E+4	1E-5	3E-8	1E-4	1E-3
28	Nickel-65	W, see ⁵⁶ Ni	-	3E+4	1E-5	4E-8	-	-
		Vapor	-	2E+4	7E-6	2E-8	-	-
		D, see ⁵⁶ Ni	4E+2	2E+3	7E-7	2E-9	-	-
28	Nickel-66	LLI wall (5E+2)	-	-	-	6E-6	6E-5	
		W, see ⁵⁶ Ni	-	6E+2	3E-7	9E-10	-	-
		Vapor	-	3E+3	1E-6	4E-9	-	-
29	Copper-60 ²	D, all compounds except those given for W and Y	3E+4	9E+4	4E-5	1E-7	-	-
		St wall (3E+4)	-	-	-	4E-4	4E-3	
		W, sulfides, halides, and nitrates	-	1E+5	5E-5	2E-7	-	-
29	Copper-61	Y, oxides and hydroxides	-	1E+5	4E-5	1E-7	-	-
		D, see ⁶⁰ Cu	1E+4	3E+4	1E-5	4E-8	2E-4	2E-3
		W, see ⁶⁰ Cu	-	4E+4	2E-5	6E-8	-	-
29	Copper-64	Y, see ⁶⁰ Cu	-	4E+4	1E-5	5E-8	-	-
		D, see ⁶⁰ Cu	1E+4	3E+4	1E-5	4E-8	2E-4	2E-3
		W, see ⁶⁰ Cu	-	2E+4	1E-5	3E-8	-	-
29	Copper-67	Y, see ⁶⁰ Cu	-	2E+4	9E-6	3E-8	-	-
		D, see ⁶⁰ Cu	5E+3	8E+3	3E-6	1E-8	6E-5	6E-4

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
		W, see ^{60}Cu	-	5E+3	2E-6	7E-9	-	-
		Y, see ^{60}Cu	-	5E+3	2E-6	6E-9	-	-
30	Zinc-62	Y, all compounds	1E+3	3E+3	1E-6	4E-9	2E-5	2E-4
30	Zinc-63 ²	Y, all compounds	2E+4	7E+4	3E-5	9E-8	-	-
			St wall (3E+4)	-	-	-	3E-4	3E-3
30	Zinc-65	Y, all compounds	4E+2	3E+2	1E-7	4E-10	5E-6	5E-5
30	Zinc-69m	Y, all compounds	4E+3	7E+3	3E-6	1E-8	6E-5	6E-4
30	Zinc-69 ²	Y, all compounds	6E+4	1E+5	6E-5	2E-7	8E-4	8E-3
30	Zinc-71m	Y, all compounds	6E+3	2E+4	7E-6	2E-8	8E-5	8E-4
30	Zinc-72	Y, all compounds	1E+3	1E+3	5E-7	2E-9	1E-5	1E-4
31	Gallium-65 ²	D, all compounds ((except except)) except those given for W	5E+4	2E+5	7E-5	2E-7	-	-
			St wall (6E+4)	-	-	-	9E-4	9E-3
		W, oxides, hydroxides, carbides, halides, and nitrates	-	2E+5	8E-5	3E-7	-	-
31	Gallium-66	D, see ^{65}Ga	1E+3	4E+3	1E-6	5E-9	1E-5	1E-4
		W, see ^{65}Ga	-	3E+3	1E-6	4E-9	-	-
31	Gallium-67	D, see ^{65}Ga	7E+3	1E+4	6E-6	2E-8	1E-4	1E-3
		W, see ^{65}Ga	-	1E+4	4E-6	1E-8	-	-
31	Gallium-68 ²	D, see ^{65}Ga	2E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, see ^{65}Ga	-	5E+4	2E-5	7E-8	-	-
31	Gallium-70 ²	D, see ^{65}Ga	5E+4	2E+5	7E-5	2E-7	-	-
			St wall (7E+4)	-	-	-	1E-3	1E-2
		W, see ^{65}Ga	-	2E+5	8E-5	3E-7	-	-
31	Gallium-72	D, see ^{65}Ga	1E+3	4E+3	1E-6	5E-9	2E-5	2E-4
		W, see ^{65}Ga	-	3E+3	1E-6	4E-9	-	-
31	Gallium-73	D, see ^{65}Ga	5E+3	2E+4	6E-6	2E-8	7E-5	7E-4
		W, see ^{65}Ga	-	2E+4	6E-6	2E-8	-	-
32	Germanium-66	D, all compounds except those given for W	2E+4	3E+4	1E-5	4E-8	3E-4	3E-3
		W, oxides, sulfides, and halides	-	2E+4	8E-6	3E-8	-	-
32	Germanium-67 ²	D, see ^{66}Ge	3E+4	9E+4	4E-5	1E-7	-	-
			St wall (4E+4)	-	-	-	6E-4	6E-3
		W, see ^{66}Ge	-	1E+5	4E-5	1E-7	-	-
32	Germanium-68	D, see ^{66}Ge	5E+3	4E+3	2E-6	5E-9	6E-5	6E-4
		W, see ^{66}Ge	-	1E+2	4E-8	1E-10	-	-
32	Germanium-69	D, see ^{66}Ge	1E+4	2E+4	6E-6	2E-8	2E-4	2E-3
		W, see ^{66}Ge	-	8E+3	3E-6	1E-8	-	-
32	Germanium-71	D, see ^{66}Ge	5E+5	4E+5	2E-4	6E-7	7E-3	7E-2
		W, see ^{66}Ge	-	4E+4	2E-5	6E-8	-	-
32	Germanium-75 ²	D, see ^{66}Ge	4E+4	8E+4	3E-5	1E-7	-	-
			St wall (7E+4)	-	-	-	9E-4	9E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
32	Germanium-77	W, see ⁶⁶ Ge	-	8E+4	4E-5	1E-7	-	-
		D, see ⁶⁶ Ge	9E+3	1E+4	4E-6	1E-8	1E-4	1E-3
32	Germanium-78 ²	W, see ⁶⁶ Ge	-	6E+3	2E-6	8E-9	-	-
		D, see ⁶⁶ Ge	2E+4	2E+4	9E-6	3E-8	-	-
33	Arsenic-69 ²	St wall (2E+4)	-	-	-	-	3E-4	3E-3
		W, see ⁶⁶ Ge	-	2E+4	9E-6	3E-8	-	-
33	Arsenic-70 ²	W, all compounds	3E+4	1E+5	5E-5	2E-7	-	-
		St wall (4E+4)	-	-	-	-	6E-4	6E-3
33	Arsenic-70 ²	W, all compounds	1E+4	5E+4	2E-5	7E-8	2E-4	2E-3
33	Arsenic-71	W, all compounds	4E+3	5E+3	2E-6	6E-9	5E-5	5E-4
33	Arsenic-72	W, all compounds	9E+2	1E+3	6E-7	2E-9	1E-5	1E-4
33	Arsenic-73	W, all compounds	8E+3	2E+3	7E-7	2E-9	1E-4	1E-3
33	Arsenic-74	W, all compounds	1E+3	8E+2	3E-7	1E-9	2E-5	2E-4
33	Arsenic-76	W, all compounds	1E+3	1E+3	6E-7	2E-9	1E-5	1E-4
33	Arsenic-77	W, all compounds	4E+3	5E+3	2E-6	7E-9	-	-
33	Arsenic-78 ²	LLI wall (5E+3)	-	-	-	-	6E-5	6E-4
		W, all compounds	8E+3	2E+4	9E-6	3E-8	1E-4	1E-3
34	Selenium-70 ²	D, all compounds except those given for W	2E+4	4E+4	2E-5	5E-8	1E-4	1E-3
34	Selenium-73m ²	W, oxides, hydroxides, carbides, and elemental Se	1E+4	4E+4	2E-5	6E-8	-	-
		D, see ⁷⁰ Se	6E+4	2E+5	6E-5	2E-7	4E-4	4E-3
34	Selenium-73	W, see ⁷⁰ Se	3E+4	1E+5	6E-5	2E-7	-	-
		D, see ⁷⁰ Se	3E+3	1E+4	5E-6	2E-8	4E-5	4E-4
34	Selenium-75	W, see ⁷⁰ Se	-	2E+4	7E-6	2E-8	-	-
		D, see ⁷⁰ Se	5E+2	7E+2	3E-7	1E-9	7E-6	7E-5
34	Selenium-79	W, see ⁷⁰ Se	-	6E+2	3E-7	8E-10	-	-
		D, see ⁷⁰ Se	6E+2	8E+2	3E-7	1E-9	8E-6	8E-5
34	Selenium-81m ²	W, see ⁷⁰ Se	-	6E+2	2E-7	8E-10	-	-
		D, see ⁷⁰ Se	4E+4	7E+4	3E-5	9E-8	3E-4	3E-3
34	Selenium-81 ²	W, see ⁷⁰ Se	2E+4	7E+4	3E-5	1E-7	-	-
		D, see ⁷⁰ Se	6E+4	2E+5	9E-5	3E-7	-	-
34	Selenium-83 ²	St wall (8E+4)	-	-	-	-	1E-3	1E-2
		W, see ⁷⁰ Se	-	2E+5	1E-4	3E-7	-	-
34	Selenium-83 ²	D, see ⁷⁰ Se	4E+4	1E+5	5E-5	2E-7	4E-4	4E-3
		W, see ⁷⁰ Se	3E+4	1E+5	5E-5	2E-7	-	-
35	Bromine-74m ²	D, bromides of H, Li, Na, K, Rb, Cs, and Fr	1E+4	4E+4	2E-5	5E-8	-	-
		St wall (2E+4)	-	-	-	-	3E-4	3E-3
		W, bromides of lanthanides, Be, Mg, Ca, Sr, Ba, Ra, Al, Ga, In, Tl, Ge, Sn, Pb, As, Sb, Bi, Fe, Ru, Os, Co, Rh, Ir, Ni, Pd, Pt, Cu, Ag, Au, Zn, Cd, Hg, Sc, Y, Ti, Zr, Hf, V, Nb, Ta, Mn, Tc, and Re	-	4E+4	2E-5	6E-8	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
35	Bromine-74 ²	D, see ^{74m} Br	2E+4	7E+4	3E-5	1E-7	-	-
			St wall (4E+4)	-	-	-	5E-45E-3	-
35	Bromine-75 ²	W, see ^{74m} Br	-	8E+4	4E-5	1E-7	-	-
		D, see ^{74m} Br	3E+4	5E+4	2E-5	7E-8	-	-
35	Bromine-76	St wall (4E+4)	-	-	-	-	5E-4	5E-3
		W, see ^{74m} Br	-	5E+4	2E-5	7E-8	-	-
35	Bromine-76	D, see ^{74m} Br	4E+3	5E+3	2E-6	7E-9	5E-5	5E-4
		W, see ^{74m} Br	-	4E+3	2E-6	6E-9	-	-
35	Bromine-77	D, see ^{74m} Br	2E+4	2E+4	1E-5	3E-8	2E-4	2E-3
		W, see ^{74m} Br	-	2E+4	8E-6	3E-8	-	-
35	Bromine-80m	D, see ^{74m} Br	2E+4	2E+4	7E-6	2E-8	3E-4	3E-3
		W, see ^{74m} Br	-	1E+4	6E-6	2E-8	-	-
35	Bromine-80 ²	D, see ^{74m} Br	5E+4	2E+5	8E-5	3E-7	-	-
			St wall (9E+4)	-	-	-	1E-3	1E-2
35	Bromine-82	W, see ^{74m} Br	-	2E+5	9E-5	3E-7	-	-
		D, see ^{74m} Br	3E+3	4E+3	2E-6	6E-9	4E-5	4E-4
35	Bromine-83	W, see ^{74m} Br	-	4E+3	2E-6	5E-9	-	-
		D, see ^{74m} Br	5E+4	6E+4	3E-5	9E-8	-	-
35	Bromine-84 ²		St wall (7E+4)	-	-	-	9E-4	9E-3
		W, see ^{74m} Br	-	6E+4	3E-5	9E-8	-	-
35	Bromine-84 ²	D, see ^{74m} Br	2E+4	6E+4	2E-5	8E-8	-	-
			St wall (3E+4)	-	-	-	4E-4	4E-3
		W, see ^{74m} Br	-	6E+4	3E-5	9E-8	-	-
36	Krypton-74 ²	Submersion ¹	-	-	3E-6	1E-8	-	-
36	Krypton-76	Submersion ¹	-	-	9E-6	4E-8	-	-
36	Krypton-77 ²	Submersion ¹	-	-	4E-6	2E-8	-	-
36	Krypton-79	Submersion ¹	-	-	2E-5	7E-8	-	-
36	Krypton-81	Submersion ¹	-	-	7E-4	3E-6	-	-
36	Krypton-83m ²	Submersion ¹	-	-	1E-2	5E-5	-	-
36	Krypton-85m	Submersion ¹	-	-	2E-5	1E-7	-	-
36	Krypton-85	Submersion ¹	-	-	1E-4	7E-7	-	-
36	Krypton-87 ²	Submersion ¹	-	-	5E-6	2E-8	-	-
36	Krypton-88	Submersion ¹	-	-	2E-6	9E-9	-	-
37	Rubidium-79 ²	D, all compounds	4E+4	1E+5	5E-5	2E-7	-	-
			St wall (6E+4)	-	-	-	8E-4	8E-3
37	Rubidium-81m ²	D, all compounds	2E+5	3E+5	1E-4	5E-7	-	-
			St wall (3E+5)	-	-	-	4E-3	4E-2
37	Rubidium-81	D, all compounds	4E+4	5E+4	2E-5	7E-8	5E-4	5E-3
37	Rubidium-82m	D, all compounds	1E+4	2E+4	7E-6	2E-8	2E-4	2E-3
37	Rubidium-83	D, all compounds	6E+2	1E+3	4E-7	1E-9	9E-6	9E-5
37	Rubidium-84	D, all compounds	5E+2	8E+2	3E-7	1E-9	7E-6	7E-5

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
37	Rubidium-86	D, all compounds	5E+2	8E+2	3E-7	1E-9	7E-6	7E-5
37	Rubidium-87	D, all compounds	1E+3	2E+3	6E-7	2E-9	1E-5	1E-4
37	Rubidium-88 ²	D, all compounds	2E+4	6E+4	3E-5	9E-8	-	-
37	Rubidium-89 ²	D, all compounds	St wall (3E+4)	-	-	-	4E-4	4E-3
			4E+4	1E+5	6E-5	2E-7	-	-
38	Strontium-80 ²	D, all soluble compound except SrTiO Y, all insoluble compounds and SrTiO	St wall (6E+4)	-	-	-	9E-4	9E-3
			4E+3	1E+4	5E-6	2E-8	6E-5	6E-4
38	Strontium-81 ²	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	-	1E+4	5E-6	2E-8	-	-
			3E+4	8E+4	3E-5	1E-7	3E-4	3E-3
38	Strontium-82	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	2E+4	8E+4	3E-5	1E-7	-	-
			3E+2	4E+2	2E-7	6E-10	-	-
38	Strontium-83	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	LLI wall (2E+2)	-	-	-	3E-6	3E-5
			2E+2	9E+1	4E-8	1E-10	-	-
38	Strontium-85m ²	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	3E+3	7E+3	3E-6	1E-8	3E-5	3E-4
			2E+3	4E+3	1E-6	5E-9	-	-
38	Strontium-85	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	2E+5	6E+5	3E-4	9E-7	3E-3	3E-2
			-	8E+5	4E-4	1E-6	-	-
38	Strontium-87m	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	3E+3	3E+3	1E-6	4E-9	4E-5	4E-4
			-	2E+3	6E-7	2E-9	-	-
38	Strontium-89	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	5E+4	1E+5	5E-5	2E-7	6E-4	6E-3
			4E+4	2E+5	6E-5	2E-7	-	-
38	Strontium-90	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	6E+2	8E+2	4E-7	1E-9	-	-
			LLI wall (6E+2)	-	-	-	8E-6	8E-5
38	Strontium-91	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	5E+2	1E+2	6E-8	2E-10	-	-
			3E+1	2E+1	8E-9	-	-	-
38	Strontium-92	D, see ⁸⁰ Sr Y, see ⁸⁰ Sr	Bone surf (4E+1)	Bone surf (2E+1)	-	3E-11	5E-7	5E-6
			-	4E+0	2E-9	6E-12	-	-
39	Yttrium-86m ²	W, all compounds except those given for Y Y, oxides and hydroxides	2E+3	6E+3	2E-6	8E-9	2E-5	2E-4
			-	4E+3	1E-6	5E-9	-	-
39	Yttrium-86	W, see ^{86m} Y Y, see ^{86m} Y	3E+3	9E+3	4E-6	1E-8	4E-5	4E-4
			-	7E+3	3E-6	9E-9	-	-
39	Yttrium-87	W, see ^{86m} Y Y, see ^{86m} Y	2E+4	6E+4	2E-5	8E-8	3E-4	3E-3
			-	5E+4	2E-5	8E-8	-	-
39	Yttrium-88	W, see ^{86m} Y Y, see ^{86m} Y	1E+3	3E+3	1E-6	5E-9	2E-5	2E-4
			-	3E+3	1E-6	5E-9	-	-
39	Yttrium-90m	W, see ^{86m} Y	2E+3	3E+3	1E-6	5E-9	3E-5	3E-4
			-	3E+3	1E-6	5E-9	-	-
39	Yttrium-90m	W, see ^{86m} Y	1E+3	3E+2	1E-7	3E-10	1E-5	1E-4
			-	2E+2	1E-7	3E-10	-	-
39	Yttrium-90m	W, see ^{86m} Y	8E+3	1E+4	5E-6	2E-8	1E-4	1E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
39	Yttrium-90	Y, see ^{86}mY	-	1E+4	5E-6	2E-8	-	-
		W, see ^{86}mY	4E+2	7E+2	3E-7	9E-10	-	-
39	Yttrium-91m ²	Y, see ^{86}mY	-	6E+2	3E-7	9E-10	-	-
		W, see ^{86}mY	1E+5	2E+5	1E-4	3E-7	2E-3	2E-2
39	Yttrium-91	Y, see ^{86}mY	-	2E+5	7E-5	2E-7	-	-
		W, see ^{86}mY	5E+2	2E+2	7E-8	2E-10	-	-
39	Yttrium-92	Y, see ^{86}mY	-	1E+2	5E-8	2E-10	-	-
		W, see ^{86}mY	3E+3	9E+3	4E-6	1E-8	4E-5	4E-4
39	Yttrium-93	Y, see ^{86}mY	-	8E+3	3E-6	1E-8	-	-
		W, see ^{86}mY	1E+3	3E+3	1E-6	4E-9	2E-5	2E-4
39	Yttrium-94 ²	Y, see ^{86}mY	-	2E+3	1E-6	3E-9	-	-
		W, see ^{86}mY	2E+4	8E+4	3E-5	1E-7	-	-
39	Yttrium-95 ²	Y, see ^{86}mY	-	8E+4	3E-5	1E-7	-	-
		W, see ^{86}mY	4E+4	2E+5	6E-5	2E-7	-	-
40	Zirconium-86	Y, see ^{86}mY	-	1E+5	6E-5	2E-7	-	-
		D, all compounds except those given for W and Y	1E+3	4E+3	2E-6	6E-9	2E-5	2E-4
40	Zirconium-88	W, oxides, hydroxides, halides, and nitrates	-	3E+3	1E-6	4E-9	-	-
		Y, carbide	-	2E+3	1E-6	3E-9	-	-
40	Zirconium-89	D, see ^{86}Zr	4E+3	2E+2	9E-8	3E-10	5E-5	5E-4
		W, see ^{86}Zr	-	5E+2	2E-7	7E-10	-	-
40	Zirconium-93	Y, see ^{86}Zr	-	3E+2	1E-7	4E-10	-	-
		D, see ^{86}Zr	2E+3	4E+3	1E-6	5E-9	2E-5	2E-4
40	Zirconium-95	W, see ^{86}Zr	-	2E+3	1E-6	3E-9	-	-
		Y, see ^{86}Zr	-	2E+3	1E-6	3E-9	-	-
40	Zirconium-97	D, see ^{86}Zr	1E+3	6E+0	3E-9	-	-	-
		W, see ^{86}Zr	-	2E+1	1E-8	-	-	-
40	Zirconium-99	Y, see ^{86}Zr	-	6E+1	2E-8	-	-	-
		D, see ^{86}Zr	1E+3	1E+2	5E-8	-	2E-5	2E-4
40	Zirconium-101	W, see ^{86}Zr	-	4E+2	2E-7	5E-10	-	-
		Y, see ^{86}Zr	-	3E+2	1E-7	4E-10	-	-
40	Zirconium-102	D, see ^{86}Zr	6E+2	2E+3	8E-7	3E-9	9E-6	9E-5
		W, see ^{86}Zr	-	4E+2	2E-7	5E-10	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
41	Niobium-88 ²	W, see ⁸⁶ Zr	-	1E+3	6E-7	2E-9	-	-
		Y, see ⁸⁶ Zr	-	1E+3	5E-7	2E-9	-	-
		W, all compounds except those given for Y	5E+4	2E+5	9E-5	3E-7	-	-
41	Niobium-89 ² (66 min)	St wall (7E+4)	-	-	-	-	1E-3	1E-2
		Y, oxides and hydroxides	-	2E+5	9E-5	3E-7	-	-
		W, see ⁸⁸ Nb	1E+4	4E+4	2E-5	6E-8	1E-4	1E-3
41	Niobium-89 (122 min)	Y, see ⁸⁸ Nb	-	4E+4	2E-5	5E-8	-	-
		W, see ⁸⁸ Nb	5E+3	2E+4	8E-6	3E-8	7E-5	7E-4
41	Niobium-90	Y, see ⁸⁸ Nb	-	2E+4	6E-6	2E-8	-	-
		W, see ⁸⁸ Nb	1E+3	3E+3	1E-6	4E-9	1E-5	1E-4
41	Niobium-93m	Y, see ⁸⁸ Nb	-	2E+3	1E-6	3E-9	-	-
		W, see ⁸⁸ Nb	9E+3	2E+3	8E-7	3E-9	-	-
		LLI wall (1E+4)	-	-	-	-	2E-4	2E-3
41	Niobium-94	Y, see ⁸⁸ Nb	-	2E+2	7E-8	2E-10	-	-
		W, see ⁸⁸ Nb	9E+2	2E+2	8E-8	3E-10	1E-5	1E-4
41	Niobium-95m	Y, see ⁸⁸ Nb	-	2E+1	6E-9	2E-11	-	-
		W, see ⁸⁸ Nb	2E+3	3E+3	1E-6	4E-9	-	-
		LLI wall (2E+3)	-	-	-	-	3E-5	3E-4
41	Niobium-95	Y, see ⁸⁸ Nb	-	2E+3	9E-7	3E-9	-	-
		W, see ⁸⁸ Nb	2E+3	1E+3	5E-7	2E-9	3E-5	3E-4
41	Niobium-96	Y, see ⁸⁸ Nb	-	1E+3	5E-7	2E-9	-	-
		W, see ⁸⁸ Nb	1E+3	3E+3	1E-6	4E-9	2E-5	2E-4
41	Niobium-97 ²	Y, see ⁸⁸ Nb	-	2E+3	1E-6	3E-9	-	-
		W, see ⁸⁸ Nb	2E+4	8E+4	3E-5	1E-7	3E-4	3E-3
41	Niobium-98 ²	Y, see ⁸⁸ Nb	-	7E+4	3E-5	1E-7	-	-
		W, see ⁸⁸ Nb	1E+4	5E+4	2E-5	8E-8	2E-4	2E-3
42	Molybdenum-90	Y, see ⁸⁸ Nb	-	5E+4	2E-5	7E-8	-	-
		D, all compounds except those given for Y	4E+3	7E+3	3E-6	1E-8	3E-5	3E-4
42	Molybdenum-93m	Y, oxides, hydroxides, and MoS	2E+3	5E+3	2E-6	6E-9	-	-
		D, see ⁹⁰ Mo	9E+3	2E+4	7E-6	2E-8	6E-5	6E-4
42	Molybdenum-93	Y, see ⁹⁰ Mo	4E+3	1E+4	6E-6	2E-8	-	-
		D, see ⁹⁰ Mo	4E+3	5E+3	2E-6	8E-9	5E-5	5E-4
42	Molybdenum-99	Y, see ⁹⁰ Mo	2E+4	2E+2	8E-8	2E-10	-	-
		D, see ⁹⁰ Mo	2E+3	3E+3	1E-6	4E-9	-	-
42	Molybdenum-101 ²	LLI wall (1E+3)	-	-	-	-	2E-5	2E-4
		Y, see ⁹⁰ Mo	1E+3	1E+3	6E-7	2E-9	-	-
		D, see ⁹⁰ Mo	4E+4	1E+5	6E-5	2E-7	-	-
		St wall (5E+4)	-	-	-	-	7E-4	7E-3
		Y, see ⁹⁰ Mo	-	1E+5	6E-5	2E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
43	Technetium-93m ²	D, all compounds except those given for W	7E+4	2E+5	6E-5	2E-7	1E-3	1E-2
		W, oxides, hydroxides, halides, and nitrates	-	3E+5	1E-4	4E-7	-	-
43	Technetium-93	D, see ^{93m} Tc	3E+4	7E+4	3E-5	1E-7	4E-4	4E-3
		W, see ^{93m} Tc	-	1E+5	4E-5	1E-7	-	-
43	Technetium-94m ²	D, see ^{93m} Tc	2E+4	4E+4	2E-5	6E-8	3E-4	3E-3
		W, see ^{93m} Tc	-	6E+4	2E-5	8E-8	-	-
43	Technetium-94	D, see ^{93m} Tc	9E+3	2E+4	8E-6	3E-8	1E-4	1E-3
		W, see ^{93m} Tc	-	2E+4	1E-5	3E-8	-	-
43	Technetium-95m	D, see ^{93m} Tc	4E+3	5E+3	2E-6	8E-9	5E-5	5E-4
		W, see ^{93m} Tc	-	2E+3	8E-7	3E-9	-	-
43	Technetium-95	D, see ^{93m} Tc	1E+4	2E+4	9E-6	3E-8	1E-4	1E-3
		W, see ^{93m} Tc	-	2E+4	8E-6	3E-8	-	-
43	Technetium-96m ²	D, see ^{93m} Tc	2E+5	3E+5	1E-4	4E-7	2E-3	2E-2
		W, see ^{93m} Tc	-	2E+5	1E-4	3E-7	-	-
43	Technetium-96	D, see ^{93m} Tc	2E+3	3E+3	1E-6	5E-9	3E-5	3E-4
		W, see ^{93m} Tc	-	2E+3	9E-7	3E-9	-	-
43	Technetium-97m	D, see ^{93m} Tc	5E+3	7E+3	3E-6	-	6E-5	6E-4
		W, see ^{93m} Tc	-	St wall (7E+3)	-	1E-8	-	-
		W, see ^{93m} Tc	-	1E+3	5E-7	2E-9	-	-
43	Technetium-97	D, see ^{93m} Tc	4E+4	5E+4	2E-5	7E-8	5E-4	5E-3
		W, see ^{93m} Tc	-	6E+3	2E-6	8E-9	-	-
43	Technetium-98	D, see ^{93m} Tc	1E+3	2E+3	7E-7	2E-9	1E-5	1E-4
		W, see ^{93m} Tc	-	3E+2	1E-7	4E-10	-	-
43	Technetium-99m	D, see ^{93m} Tc	8E+4	2E+5	6E-5	2E-7	1E-3	1E-2
		W, see ^{93m} Tc	-	2E+5	1E-4	3E-7	-	-
43	Technetium-99	D, see ^{93m} Tc	4E+3	5E+3	2E-6	-	6E-5	6E-4
		W, see ^{93m} Tc	-	St wall (6E+3)	-	8E-9	-	-
		W, see ^{93m} Tc	-	7E+2	3E-7	9E-10	-	-
43	Technetium-101 ²	D, see ^{93m} Tc	9E+4	3E+5	1E-4	5E-7	-	-
		W, see ^{93m} Tc	-	St wall (1E+5)	-	-	2E-3	2E-2
		W, see ^{93m} Tc	-	4E+5	2E-4	5E-7	-	-
43	Technetium-104 ²	D, see ^{93m} Tc	2E+4	7E+4	3E-5	1E-7	-	-
		W, see ^{93m} Tc	-	St wall (3E+4)	-	-	4E-4	4E-3
		W, see ^{93m} Tc	-	9E+4	4E-5	1E-7	-	-
44	Ruthenium-94 ²	D, all compounds except those given for W and Y	2E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, halides	-	6E+4	3E-5	9E-8	-	-
		Y, oxides and hydroxides	-	6E+4	2E-5	8E-8	-	-
44	Ruthenium-97	D, see ⁹⁴ Ru	8E+3	2E+4	8E-6	3E-8	1E-4	1E-3
		W, see ⁹⁴ Ru	-	1E+4	5E-6	2E-8	-	-
		Y, see ⁹⁴ Ru	-	1E+4	5E-6	2E-8	-	-
44	Ruthenium-103	D, see ⁹⁴ Ru	2E+3	2E+3	7E-7	2E-9	3E-5	3E-4

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
44	Ruthenium-105	W, see ^{94}Ru	-	1E+3	4E-7	1E-9	-	-
		Y, see ^{94}Ru	-	6E+2	3E-7	9E-10	-	-
		D, see ^{94}Ru	5E+3	1E+4	6E-6	2E-8	7E-5	7E-4
44	Ruthenium-106	W, see ^{94}Ru	-	1E+4	6E-6	2E-8	-	-
		Y, see ^{94}Ru	-	1E+4	5E-6	2E-8	-	-
		D, see ^{94}Ru	2E+2	9E+1	4E-8	1E-10	-	-
45	Rhodium-99m	LLI wall (2E+2)	-	-	-	-	3E-6	3E-5
		W, see ^{94}Ru	-	5E+1	2E-8	8E-11	-	-
		Y, see ^{94}Ru	-	1E+1	5E-9	2E-11	-	-
		D, all compounds except those given for W and Y	2E+4	6E+4	2E-5	8E-8	2E-4	2E-3
		W, halides	-	8E+4	3E-5	1E-7	-	-
45	Rhodium-99	Y, oxides and hydroxides	-	7E+4	3E-5	9E-8	-	-
		D, see ^{99m}Rh	2E+3	3E+3	1E-6	4E-9	3E-5	3E-4
		W, see ^{99m}Rh	-	2E+3	9E-7	3E-9	-	-
45	Rhodium-100	Y, see ^{99m}Rh	-	2E+3	8E-7	3E-9	-	-
		D, see ^{99m}Rh	2E+3	5E+3	2E-6	7E-9	2E-5	2E-4
		W, see ^{99m}Rh	-	4E+3	2E-6	6E-9	-	-
45	Rhodium-101m	Y, see ^{99m}Rh	-	4E+3	2E-6	5E-9	-	-
		D, see ^{99m}Rh	6E+3	1E+4	5E-6	2E-8	8E-5	8E-4
		W, see ^{99m}Rh	-	8E+3	4E-6	1E-8	-	-
45	Rhodium-101	Y, see ^{99m}Rh	-	8E+3	3E-6	1E-8	-	-
		D, see ^{99m}Rh	2E+3	5E+2	2E-7	7E-10	3E-5	3E-4
		W, see ^{99m}Rh	-	8E+2	3E-7	1E-9	-	-
45	Rhodium-102m	Y, see ^{99m}Rh	-	2E+2	6E-8	2E-10	-	-
		D, see ^{99m}Rh	1E+3	5E+2	2E-7	7E-10	-	-
		LLI wall (1E+3)	-	-	-	-	2E-5	2E-4
		W, see ^{99m}Rh	-	4E+2	2E-7	5E-10	-	-
		Y, see ^{99m}Rh	-	1E+2	5E-8	2E-10	-	-
45	Rhodium-102	D, see ^{99m}Rh	6E+2	9E+1	4E-8	1E-10	8E-6	8E-5
		W, see ^{99m}Rh	-	2E+2	7E-8	2E-10	-	-
		Y, see ^{99m}Rh	-	6E+1	2E-8	8E-11	-	-
45	Rhodium-103m ²	D, see ^{99m}Rh	4E+5	1E+6	5E-4	2E-6	6E-3	6E-2
		W, see ^{99m}Rh	-	1E+6	5E-4	2E-6	-	-
		Y, see ^{99m}Rh	-	1E+6	5E-4	2E-6	-	-
45	Rhodium-105	D, see ^{99m}Rh	4E+3	1E+4	5E-6	2E-8	-	-
		LLI wall (4E+3)	-	-	-	-	5E-5	5E-4
		W, see ^{99m}Rh	-	6E+3	3E-6	9E-9	-	-
		Y, see ^{99m}Rh	-	6E+3	2E-6	8E-9	-	-
		D, see ^{99m}Rh	8E+3	3E+4	1E-5	4E-8	1E-4	1E-3
45	Rhodium-106m	W, see ^{99m}Rh	-	4E+4	2E-5	5E-8	-	-
		Y, see ^{99m}Rh	-	4E+4	1E-5	5E-8	-	-
		D, see ^{99m}Rh	7E+4	2E+5	1E-4	3E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
			St wall (9E+4)	-	-	-	1E-3	1E-2
		W, see ^{99m} Rh	-	3E+5	1E-4	4E-7	-	-
		Y, see ^{99m} Rh	-	3E+5	1E-4	3E-7	-	-
46	Palladium-100	D, all compounds except those given for W and Y	1E+3	1E+3	6E-7	2E-9	2E-5	2E-4
		W, nitrates	-	1E+3	5E-7	2E-9	-	-
		Y, oxides and hydroxides	-	1E+3	6E-7	2E-9	-	-
46	Palladium-101	D, see ¹⁰⁰ Pd	1E+4	3E+4	1E-5	5E-8	2E-4	2E-3
		W, see ¹⁰⁰ Pd	-	3E+4	1E-5	5E-8	-	-
		Y, see ¹⁰⁰ Pd	-	3E+4	1E-5	4E-8	-	-
46	Palladium-103	D, see ¹⁰⁰ Pd	6E+3	6E+3	3E-6	9E-9	-	-
			LLI wall (7E+3)	-	-	-	1E-4	1E-3
		W, see ¹⁰⁰ Pd	-	4E+3	2E-6	6E-9	-	-
		Y, see ¹⁰⁰ Pd	-	4E+3	1E-6	5E-9	-	-
46	Palladium-107	D, see ¹⁰⁰ Pd	3E+4	2E+4	9E-6	-	-	-
			LLI wall (4E+4)	Kidneys (2E+4)	-	3E-8	5E-4	5E-3
		W, see ¹⁰⁰ Pd	-	7E+3	3E-6	1E-8	-	-
		Y, see ¹⁰⁰ Pd	-	4E+2	2E-7	6E-10	-	-
46	Palladium-109	D, see ¹⁰⁰ Pd	2E+3	6E+3	3E-6	9E-9	3E-5	3E-4
		W, see ¹⁰⁰ Pd	-	5E+3	2E-6	8E-9	-	-
		Y, see ¹⁰⁰ Pd	-	5E+3	2E-6	6E-9	-	-
47	Silver-102 ²	D, all compounds except those given for W and Y	5E+4	2E+5	8E-5	2E-7	-	-
			St wall (6E+4)	-	-	-	9E-4	9E-3
		W, nitrates and sulfides	-	2E+5	9E-5	3E-7	-	-
		Y, oxides and hydroxides	-	2E+5	8E-5	3E-7	-	-
47	Silver-103 ²	D, see ¹⁰² Ag	4E+4	1E+5	4E-5	1E-7	5E-4	5E-3
		W, see ¹⁰² Ag	-	1E+5	5E-5	2E-7	-	-
		Y, see ¹⁰² Ag	-	1E+5	5E-5	2E-7	-	-
47	Silver-104m ²	D, see ¹⁰² Ag	3E+4	9E+4	4E-5	1E-7	4E-4	4E-3
		W, see ¹⁰² Ag	-	1E+5	5E-5	2E-7	-	-
		Y, see ¹⁰² Ag	-	1E+5	5E-5	2E-7	-	-
47	Silver-104 ²	D, see ¹⁰² Ag	2E+4	7E+4	3E-5	1E-7	3E-4	3E-3
		W, see ¹⁰² Ag	-	1E+5	6E-5	2E-7	-	-
		Y, see ¹⁰² Ag	-	1E+5	6E-5	2E-7	-	-
47	Silver-105	D, see ¹⁰² Ag	3E+3	1E+3	4E-7	1E-9	4E-5	4E-4
		W, see ¹⁰² Ag	-	2E+3	7E-7	2E-9	-	-
		Y, see ¹⁰² Ag	-	2E+3	7E-7	2E-9	-	-
47	Silver-106m	D, see ¹⁰² Ag	8E+2	7E+2	3E-7	1E-9	1E-5	1E-4
		W, see ¹⁰² Ag	-	9E+2	4E-7	1E-9	-	-
		Y, see ¹⁰² Ag	-	9E+2	4E-7	1E-9	-	-
47	Silver-106 ²	D, see ¹⁰² Ag	6E+4	2E+5	8E-5	3E-7	-	-
			St. wall (6E+4)	-	-	-	9E-4	9E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
47	Silver-108m	W, see ¹⁰² Ag	-	2E+5	9E-5	3E-7	-	-
		Y, see ¹⁰² Ag	-	2E+5	8E-5	3E-7	-	-
		D, see ¹⁰² Ag	6E+2	2E+2	8E-8	3E-10	9E-6	9E-5
		W, see ¹⁰² Ag	-	3E+2	1E-7	4E-10	-	-
		Y, see ¹⁰² Ag	-	2E+1	1E-8	3E-11	-	-
47	Silver-110m	D, see ¹⁰² Ag	5E+2	1E+2	5E-8	2E-10	6E-6	6E-5
		W, see ¹⁰² Ag	-	2E+2	8E-8	3E-10	-	-
		Y, see ¹⁰² Ag	-	9E+1	4E-8	1E-10	-	-
47	Silver-111	D, see ¹⁰² Ag	9E+2	2E+3	6E-7	-	-	-
			LLI wall (1E+3)	Liver (2E+3)	-	2E-9	2E-5	2E-4
47	Silver-112	W, see ¹⁰² Ag	-	9E+2	4E-7	1E-9	-	-
		Y, see ¹⁰² Ag	-	9E+2	4E-7	1E-9	-	-
		D, see ¹⁰² Ag	3E+3	8E+3	3E-6	1E-8	4E-5	4E-4
47	Silver-115 ²	W, see ¹⁰² Ag	-	1E+4	4E-6	1E-8	-	-
		Y, see ¹⁰² Ag	-	9E+3	4E-6	1E-8	-	-
		D, see ¹⁰² Ag	3E+4	9E+4	4E-5	1E-7	-	-
			St wall (3E+4)	-	-	-	4E-4	4E-3
48	Cadmium-104 ²	W, see ¹⁰² Ag	-	9E+4	4E-5	1E-7	-	-
		Y, see ¹⁰² Ag	-	8E+4	3E-5	1E-7	-	-
		D, all compounds except those given for W and Y	2E+4	7E+4	3E-5	9E-8	3E-4	3E-3
		W, sulfides, halides, and nitrates	-	1E+5	5E-5	2E-7	-	-
48	Cadmium-107	Y, oxides and hydroxides	-	1E+5	5E-5	2E-7	-	-
		D, see ¹⁰⁴ Cd	2E+4	5E+4	2E-5	8E-8	3E-4	3E-3
		W, see ¹⁰⁴ Cd	-	6E+4	2E-5	8E-8	-	-
48	Cadmium-109	Y, see ¹⁰⁴ Cd	-	5E+4	2E-5	7E-8	-	-
		D, see ¹⁰⁴ Cd	3E+2	4E+1	1E-8	-	-	-
			Kidneys (4E+2)	Kidneys (5E+1)	-	7E-11	6E-6	6E-5
48	Cadmium-113m	W, see ¹⁰⁴ Cd	-	1E+2	5E-8	-	-	-
			-	Kidneys (1E+2)	-	2E-10	-	-
		Y, see ¹⁰⁴ Cd	-	1E+2	5E-8	2E-10	-	-
		D, see ¹⁰⁴ Cd	2E+1	2E+0	1E-9	-	-	-
			Kidneys (4E+1)	Kidneys (4E+0)	-	5E-12	5E-7	5E-6
48	Cadmium-113	W, see ¹⁰⁴ Cd	-	8E+0	4E-9	-	-	-
			-	Kidneys (1E+1)	-	2E-11	-	-
		Y, see ¹⁰⁴ Cd	-	1E+1	5E-9	2E-11	-	-
		D, see ¹⁰⁴ Cd	2E+1	2E+0	9E-10	-	-	-
	Kidneys (3E+1)	Kidneys (3E+0)	-	5E-12	4E-7	4E-6		
	W, see ¹⁰⁴ Cd	-	8E+0	3E-9	-	-	-	
	-	Kidneys (1E+1)	-	2E-11	-	-		

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
48	Cadmium-115m	Y, see ^{104}Cd	-	1E+1	6E-9	2E-11	-	-
		D, see ^{104}Cd	3E+2	5E+1	2E-8	-	4E-6	4E-5
48	Cadmium-115		-	Kidneys (8E+1)	-	1E-10	-	-
		W, see ^{104}Cd	-	1E+2	5E-8	2E-10	-	-
		Y, see ^{104}Cd	-	1E+2	6E-8	2E-10	-	-
		D, see ^{104}Cd	9E+2	1E+3	6E-7	2E-9	-	-
48	Cadmium-117m		LLI wall (1E+3)	-	-	-	1E-5	1E-4
		W, see ^{104}Cd	-	1E+3	5E-7	2E-9	-	-
		Y, see ^{104}Cd	-	1E+3	6E-7	2E-9	-	-
		D, see ^{104}Cd	5E+3	1E+4	5E-6	2E-8	6E-5	6E-4
48	Cadmium-117	W, see ^{104}Cd	-	2E+4	7E-6	2E-8	-	-
		Y, see ^{104}Cd	-	1E+4	6E-6	2E-8	-	-
		D, see ^{104}Cd	5E+3	1E+4	5E-6	2E-8	6E-5	6E-4
49	Indium-109	W, see ^{104}Cd	-	2E+4	7E-6	2E-8	-	-
		Y, see ^{104}Cd	-	1E+4	6E-6	2E-8	-	-
		D, all compounds except those given for W	2E+4	4E+4	2E-5	6E-8	3E-4	3E-3
49	Indium-110 ² (69.1 min)	W, oxides, hydroxides, halides, and nitrates	-	6E+4	3E-5	9E-8	-	-
		D, see ^{109}In	2E+4	4E+4	2E-5	6E-8	2E-4	2E-3
49	Indium-110 (4.9 h)	W, see ^{109}In	-	6E+4	2E-5	8E-8	-	-
		D, see ^{109}In	5E+3	2E+4	7E-6	2E-8	7E-5	7E-4
49	Indium-111	W, see ^{109}In	-	2E+4	8E-6	3E-8	-	-
		D, see ^{109}In	4E+3	6E+3	3E-6	9E-9	6E-5	6E-4
49	Indium-112 ²	W, see ^{109}In	-	6E+3	3E-6	9E-9	-	-
		D, see ^{109}In	2E+5	6E+5	3E-4	9E-7	2E-3	2E-2
49	Indium-113m ²	W, see ^{109}In	-	7E+5	3E-4	1E-6	-	-
		D, see ^{109}In	5E+4	1E+5	6E-5	2E-7	7E-4	7E-3
49	Indium-114m	W, see ^{109}In	-	2E+5	8E-5	3E-7	-	-
		D, see ^{109}In	3E+2	6E+1	3E-8	9E-11	-	-
49	Indium-115m		LLI wall (4E+2)	-	-	-	5E-6	5E-5
		W, see ^{109}In	-	1E+2	4E-8	1E-10	-	-
49	Indium-115	D, see ^{109}In	1E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, see ^{109}In	-	5E+4	2E-5	7E-8	-	-
49	Indium-116m ²	D, see ^{109}In	4E+1	1E+0	6E-10	2E-12	5E-7	5E-6
		W, see ^{109}In	-	5E+0	2E-9	8E-12	-	-
49	Indium-117m ²	D, see ^{109}In	2E+4	8E+4	3E-5	1E-7	3E-4	3E-3
		W, see ^{109}In	-	1E+5	5E-5	2E-7	-	-
49	Indium-117 ²	D, see ^{109}In	1E+4	3E+4	1E-5	5E-8	2E-4	2E-3
		W, see ^{109}In	-	4E+4	2E-5	6E-8	-	-
49	Indium-119m ²	D, see ^{109}In	6E+4	2E+5	7E-5	2E-7	8E-4	8E-3
		W, see ^{109}In	-	2E+5	9E-5	3E-7	-	-
49	Indium-119m ²	D, see ^{109}In	4E+4	1E+5	5E-5	2E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
			St wall (5E+4)	-	-	-	7E-4	7E-3
50	Tin-110	W, see ¹⁰⁹ In D, all compounds except those given for W	-	1E+5	6E-5	2E-7	-	-
			4E+3	1E+4	5E-6	2E-8	5E-5	5E-4
50	Tin-111 ²	W, sulfides, oxides, hydroxides, halides, nitrates, and stannic phosphate D, see ¹¹⁰ Sn	-	1E+4	5E-6	2E-8	-	-
			7E+4	2E+5	9E-5	3E-7	1E-3	1E-2
50	Tin-113	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	3E+5	1E-4	4E-7	-	-
			2E+3	1E+3	5E-7	2E-9	-	-
			LLI wall (2E+3)	-	-	-	3E-5	3E-4
50	Tin-117m	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	5E+2	2E-7	8E-10	-	-
			2E+3	1E+3	5E-7	-	-	-
			LLI wall (2E+3)	Bone surf (2E+3)	-	3E-9	3E-5	3E-4
50	Tin-119m	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	1E+3	6E-7	2E-9	-	-
			3E+3	2E+3	1E-6	3E-9	-	-
			LLI wall (4E+3)	-	-	-	6E-5	6E-4
50	Tin-121m	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	1E+3	4E-7	1E-9	-	-
			3E+3	9E+2	4E-7	1E-9	-	-
			LLI wall (4E+3)	-	-	-	5E-5	5E-4
50	Tin-121	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	5E+2	2E-7	8E-10	-	-
			6E+3	2E+4	6E-6	2E-8	-	-
			LLI wall (6E+3)	-	-	-	8E-5	8E-4
50	Tin-123m ²	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	1E+4	5E-6	2E-8	-	-
			5E+4	1E+5	5E-5	2E-7	7E-4	7E-3
50	Tin-123	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	1E+5	6E-5	2E-7	-	-
			5E+2	6E+2	3E-7	9E-10	-	-
			LLI wall (6E+2)	-	-	-	9E-6	9E-5
50	Tin-125	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	2E+2	7E-8	2E-10	-	-
			4E+2	9E+2	4E-7	1E-9	-	-
			LLI wall (5E+2)	-	-	-	6E-6	6E-5
50	Tin-126	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	4E+2	1E-7	5E-10	-	-
			3E+2	6E+1	2E-8	8E-11	4E-6	4E-5
50	Tin-127	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	7E+1	3E-8	9E-11	-	-
			7E+3	2E+4	8E-6	3E-8	9E-5	9E-4
50	Tin-128 ²	W, see ¹¹⁰ Sn D, see ¹¹⁰ Sn	-	2E+4	8E-6	3E-8	-	-
			9E+3	3E+4	1E-5	4E-8	1E-4	1E-3
51	Antimony-115 ²	W, see ¹¹⁰ Sn D, all compounds except those given for W	-	4E+4	1E-5	5E-8	-	-
			8E+4	2E+5	1E-4	3E-7	1E-3	1E-2
		W, oxides, hydroxides, halides, sulfides, sulfates, and nitrates	-	3E+5	1E-4	4E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
51	Antimony-116m ²	D, see ¹¹⁵ Sb	2E+4	7E+4	3E-5	1E-7	3E-4	3E-3
		W, see ¹¹⁵ Sb	-	1E+5	6E-5	2E-7	-	-
51	Antimony-116 ²	D, see ¹¹⁵ Sb	7E+4	3E+5	1E-4	4E-7	-	-
		St wall (9E+4)	-	-	-	-	1E-3	1E-2
51	Antimony-117	W, see ¹¹⁵ Sb	-	3E+5	1E-4	5E-7	-	-
		D, see ¹¹⁵ Sb	7E+4	2E+5	9E-5	3E-7	9E-4	9E-3
51	Antimony-118m	W, see ¹¹⁵ Sb	-	3E+5	1E-4	4E-7	-	-
		D, see ¹¹⁵ Sb	6E+3	2E+4	8E-6	3E-8	7E-5	7E-4
51	Antimony-119	W, see ¹¹⁵ Sb	5E+3	2E+4	9E-6	3E-8	-	-
		D, see ¹¹⁵ Sb	2E+4	5E+4	2E-5	6E-8	2E-4	2E-3
51	Antimony-120 ² (16 min)	W, see ¹¹⁵ Sb	2E+4	3E+4	1E-5	4E-8	-	-
		D, see ¹¹⁵ Sb	1E+5	4E+5	2E-4	6E-7	-	-
51	Antimony-120 (5.76 d)	St wall (2E+5)	-	-	-	-	2E-3	2E-2
		W, see ¹¹⁵ Sb	-	5E+5	2E-4	7E-7	-	-
51	Antimony-122	D, see ¹¹⁵ Sb	1E+3	2E+3	9E-7	3E-9	1E-5	1E-4
		W, see ¹¹⁵ Sb	9E+2	1E+3	5E-7	2E-9	-	-
51	Antimony-124m ²	D, see ¹¹⁵ Sb	8E+2	2E+3	1E-6	3E-9	-	-
		LLI wall (8E+2)	-	-	-	-	1E-5	1E-4
51	Antimony-124	W, see ¹¹⁵ Sb	7E+2	1E+3	4E-7	2E-9	-	-
		D, see ¹¹⁵ Sb	3E+5	8E+5	4E-4	1E-6	3E-3	3E-2
51	Antimony-125	W, see ¹¹⁵ Sb	2E+5	6E+5	2E-4	8E-7	-	-
		D, see ¹¹⁵ Sb	6E+2	9E+2	4E-7	1E-9	7E-6	7E-5
51	Antimony-126m ²	W, see ¹¹⁵ Sb	5E+2	2E+2	1E-7	3E-10	-	-
		D, see ¹¹⁵ Sb	2E+3	2E+3	1E-6	3E-9	3E-5	3E-4
51	Antimony-126	W, see ¹¹⁵ Sb	-	5E+2	2E-7	7E-10	-	-
		D, see ¹¹⁵ Sb	5E+4	2E+5	8E-5	3E-7	-	-
51	Antimony-127	St wall (7E+4)	-	-	-	-	9E-4	9E-3
		W, see ¹¹⁵ Sb	-	2E+5	8E-5	3E-7	-	-
51	Antimony-128 ² (10.4 min)	D, see ¹¹⁵ Sb	6E+2	1E+3	5E-7	2E-9	7E-6	7E-5
		W, see ¹¹⁵ Sb	5E+2	5E+2	2E-7	7E-10	-	-
51	Antimony-128 (9.01 h)	D, see ¹¹⁵ Sb	8E+2	2E+3	9E-7	3E-9	-	-
		LLI wall (8E+2)	-	-	-	-	1E-5	1E-4
51	Antimony-129	W, see ¹¹⁵ Sb	7E+2	9E+2	4E-7	1E-9	-	-
		D, see ¹¹⁵ Sb	8E+4	4E+5	2E-4	5E-7	-	-
51	Antimony-129	St wall (1E+5)	-	-	-	-	1E-3	1E-2
		W, see ¹¹⁵ Sb	-	4E+5	2E-4	6E-7	-	-
51	Antimony-129	D, see ¹¹⁵ Sb	1E+3	4E+3	2E-6	6E-9	2E-5	2E-4
		W, see ¹¹⁵ Sb	-	3E+3	1E-6	5E-9	-	-
51	Antimony-129	D, see ¹¹⁵ Sb	3E+3	9E+3	4E-6	1E-8	4E-5	4E-4
		W, see ¹¹⁵ Sb	-	9E+3	4E-6	1E-8	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
51	Antimony-130 ²	D, see ¹¹⁵ Sb	2E+4	6E+4	3E-5	9E-8	3E-4	3E-3
		W, see ¹¹⁵ Sb	-	8E+4	3E-5	1E-7	-	-
51	Antimony-131 ²	D, see ¹¹⁵ Sb	1E+4	2E+4	1E-5	-	-	-
			Thyroid (2E+4)	Thyroid (4E+4)	-	6E-8	2E-4	2E-3
		W, see ¹¹⁵ Sb	-	2E+4	1E-5	-	-	-
			-	Thyroid (4E+4)	-	6E-8	-	-
52	Tellurium-116	D, all compounds except those given for W	8E+3	2E+4	9E-6	3E-8	1E-4	1E-3
		W, oxides, hydroxides, and nitrates	-	3E+4	1E-5	4E-8	-	-
52	Tellurium-121m	D, see ¹¹⁶ Te	5E+2	2E+2	8E-8	-	-	-
			Bone surf (7E+2)	Bone surf (4E+2)	-	5E-10	1E-5	1E-4
		W, see ¹¹⁶ Te	-	4E+2	2E-7	6E-10	-	-
52	Tellurium-121	D, see ¹¹⁶ Te	3E+3	4E+3	2E-6	6E-9	4E-5	4E-4
		W, see ¹¹⁶ Te	-	3E+3	1E-6	4E-9	-	-
52	Tellurium-123m	D, see ¹¹⁶ Te	6E+2	2E+2	9E-8	-	-	-
			Bone surf (1E+3)	Bone surf (5E+2)	-	8E-10	1E-5	1E-4
		W, see ¹¹⁶ Te	-	5E+2	2E-7	8E-10	-	-
52	Tellurium-123	D, see ¹¹⁶ Te	5E+2	2E+2	8E-8	-	-	-
			Bone surf (1E+3)	Bone surf (5E+2)	-	7E-10	2E-5	2E-4
		W, see ¹¹⁶ Te	-	4E+2	2E-7	-	-	-
			-	Bone surf (1E+3)	-	2E-9	-	-
52	Tellurium-125m	D, see ¹¹⁶ Te	1E+3	4E+2	2E-7	-	-	-
			Bone surf (1E+3)	Bone surf (1E+3)	-	1E-9	2E-5	2E-4
		W, see ¹¹⁶ Te	-	7E+2	3E-7	1E-9	-	-
52	Tellurium-127m	D, see ¹¹⁶ Te	6E+2	3E+2	1E-7	-	9E-6	9E-5
			-	Bone surf (4E+2)	-	6E-10	-	-
		W, see ¹¹⁶ Te	-	3E+2	1E-7	4E-10	-	-
52	Tellurium-127	D, see ¹¹⁶ Te	7E+3	2E+4	9E-6	3E-8	1E-4	1E-3
		W, see ¹¹⁶ Te	-	2E+4	7E-6	2E-8	-	-
52	Tellurium-129m	D, see ¹¹⁶ Te	5E+2	6E+2	3E-7	9E-10	7E-6	7E-5
		W, see ¹¹⁶ Te	-	2E+2	1E-7	3E-10	-	-
52	Tellurium-129 ²	D, see ¹¹⁶ Te	3E+4	6E+4	3E-5	9E-8	4E-4	4E-3
		W, see ¹¹⁶ Te	-	7E+4	3E-5	1E-7	-	-
52	Tellurium-131m	D, see ¹¹⁶ Te	3E+2	4E+2	2E-7	-	-	-
			Thyroid (6E+2)	Thyroid (1E+3)	-	2E-9	8E-6	8E-5
		W, see ¹¹⁶ Te	-	4E+2	2E-7	-	-	-
			-	Thyroid (9E+2)	-	1E-9	-	-
52	Tellurium-131 ²	D, see ¹¹⁶ Te	3E+3	5E+3	2E-6	-	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
52	Tellurium-132	W, see ¹¹⁶ Te	Thyroid (6E+3)	Thyroid (1E+4)	-	2E-8	8E-5	8E-4
			-	5E+3	2E-6	-	-	-
		D, see ¹¹⁶ Te	-	Thyroid (1E+4)	-	2E-8	-	-
			2E+2	2E+2	9E-8	-	-	-
52	Tellurium-133m ²	W, see ¹¹⁶ Te	Thyroid (7E+2)	Thyroid (8E+2)	-	1E-9	9E-6	9E-5
			-	2E+2	9E-8	-	-	-
		D, see ¹¹⁶ Te	-	Thyroid (6E+2)	-	9E-10	-	-
			3E+3	5E+3	2E-6	-	-	-
52	Tellurium-133 ²	W, see ¹¹⁶ Te	Thyroid (6E+3)	Thyroid (1E+4)	-	2E-8	9E-5	9E-4
			-	5E+3	2E-6	-	-	-
		D, see ¹¹⁶ Te	-	Thyroid (1E+4)	-	2E-8	-	-
			1E+4	2E+4	9E-6	-	-	-
52	Tellurium-134 ²	W, see ¹¹⁶ Te	Thyroid (3E+4)	Thyroid (6E+4)	-	8E-8	4E-4	4E-3
			-	2E+4	9E-6	-	-	-
		D, see ¹¹⁶ Te	-	Thyroid (6E+4)	-	8E-8	-	-
			2E+4	2E+4	1E-5	-	-	-
53	Iodine-120m ²	W, see ¹¹⁶ Te	Thyroid (2E+4)	Thyroid (5E+4)	-	7E-8	3E-4	3E-3
			-	2E+4	1E-5	-	-	-
		D, all compounds	-	Thyroid (5E+4)	-	7E-8	-	-
			1E+4	2E+4	9E-6	3E-8	-	-
53	Iodine-120 ²	D, all compounds	Thyroid (1E+4)	-	-	-	2E-4	2E-3
			4E+3	9E+3	4E-6	-	-	-
53	Iodine-121	D, all compounds	Thyroid (8E+3)	Thyroid (1E+4)	-	2E-8	1E-4	1E-3
			1E+4	2E+4	8E-6	-	-	-
53	Iodine-123	D, all compounds	Thyroid (3E+4)	Thyroid (5E+4)	-	7E-8	4E-4	4E-3
			3E+3	6E+3	3E-6	-	-	-
53	Iodine-124	D, all compounds	Thyroid (1E+4)	Thyroid (2E+4)	-	2E-8	1E-4	1E-3
			5E+1	8E+1	3E-8	-	-	-
53	Iodine-125	D, all compounds	Thyroid (2E+2)	Thyroid (3E+2)	-	4E-10	2E-6	2E-5
			4E+1	6E+1	3E-8	-	-	-
53	Iodine-126	D, all compounds	Thyroid (1E+2)	Thyroid (2E+2)	-	3E-10	2E-6	2E-5
			2E+1	4E+1	1E-8	-	-	-
53	Iodine-128 ²	D, all compounds	Thyroid (7E+1)	Thyroid (1E+2)	-	2E-10	1E-6	1E-5
			4E+4	1E+5	5E-5	2E-7	-	-
			St wall (6E+4)	-	-	-	8E-4	8E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
53	Iodine-129	D, all compounds	5E+0	9E+0	4E-9	-	-	-
			Thyroid (2E+1)	Thyroid (3E+1)	-	4E-11	2E-7	2E-6
53	Iodine-130	D, all compounds	4E+2	7E+2	3E-7	-	-	-
			Thyroid (1E+3)	Thyroid (2E+3)	-	3E-9	2E-5	2E-4
53	Iodine-131	D, all compounds	3E+1	5E+1	2E-8	-	-	-
			Thyroid (9E+1)	Thyroid (2E+2)	-	2E-10	1E-6	1E-5
53	Iodine-132m ²	D, all compounds	4E+3	8E+3	4E-6	-	-	-
			Thyroid (1E+4)	Thyroid (2E+4)	-	3E-8	1E-4	1E-3
53	Iodine-132	D, all compounds	4E+3	8E+3	3E-6	-	-	-
			Thyroid (9E+3)	Thyroid (1E+4)	-	2E-8	1E-4	1E-3
53	Iodine-133	D, all compounds	1E+2	3E+2	1E-7	-	-	-
			Thyroid (5E+2)	Thyroid (9E+2)	-	1E-9	7E-6	7E-5
53	Iodine-134 ²	D, all compounds	2E+4	5E+4	2E-5	6E-8	-	-
			Thyroid (3E+4)	-	-	-	4E-4	4E-3
53	Iodine-135	D, all compounds	8E+2	2E+3	7E-7	-	-	-
			Thyroid (3E+3)	Thyroid (4E+3)	-	6E-9	3E-5	3E-4
54	Xenon-120 ²	Submersion ¹	-	-	1E-5	4E-8	-	-
54	Xenon-121 ²	Submersion ¹	-	-	2E-6	1E-8	-	-
54	Xenon-122	Submersion ¹	-	-	7E-5	3E-7	-	-
54	Xenon-123	Submersion ¹	-	-	6E-6	3E-8	-	-
54	Xenon-125	Submersion ¹	-	-	2E-5	7E-8	-	-
54	Xenon-127	Submersion ¹	-	-	1E-5	6E-8	-	-
54	Xenon-129m	Submersion ¹	-	-	2E-4	9E-7	-	-
54	Xenon-131m	Submersion ¹	-	-	4E-4	2E-6	-	-
54	Xenon-133m	Submersion ¹	-	-	1E-4	6E-7	-	-
54	Xenon-133	Submersion ¹	-	-	1E-4	5E-7	-	-
54	Xenon-135m ²	Submersion ¹	-	-	9E-6	4E-8	-	-
54	Xenon-135	Submersion ¹	-	-	1E-5	7E-8	-	-
54	Xenon-138 ²	Submersion ¹	-	-	4E-6	2E-8	-	-
55	Cesium-125 ²	D, all compounds	5E+4	1E+5	6E-5	2E-7	-	-
			St wall (9E+4)	-	-	-	1E-3	1E-2
55	Cesium-127	D, all compounds	6E+4	9E+4	4E-5	1E-7	9E-4	9E-3
55	Cesium-129	D, all compounds	2E+4	3E+4	1E-5	5E-8	3E-4	3E-3
55	Cesium-130 ²	D, all compounds	6E+4	2E+5	8E-5	3E-7	-	-
			St wall (1E+5)	-	-	-	1E-3	1E-2
55	Cesium-131	D, all compounds	2E+4	3E+4	1E-5	4E-8	3E-4	3E-3
55	Cesium-132	D, all compounds	3E+3	4E+3	2E-6	6E-9	4E-5	4E-4
55	Cesium-134m	D, all compounds	1E+5	1E+5	6E-5	2E-7	-	-
			St wall (1E+5)	-	-	-	2E-3	2E-2
55	Cesium-134	D, all compounds	7E+1	1E+2	4E-8	2E-10	9E-7	9E-6

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
55	Cesium-135m ²	D, all compounds	1E+5	2E+5	8E-5	3E-7	1E-3	1E-2
55	Cesium-135	D, all compounds	7E+2	1E+3	5E-7	2E-9	1E-5	1E-4
55	Cesium-136	D, all compounds	4E+2	7E+2	3E-7	9E-10	6E-6	6E-5
55	Cesium-137	D, all compounds	1E+2	2E+2	6E-8	2E-10	1E-6	1E-5
55	Cesium-138 ²	D, all compounds	2E+4	6E+4	2E-5	8E-8	-	-
			St wall (3E+4)	-	-	-	4E-4	4E-3
56	Barium-126 ²	D, all compounds	6E+3	2E+4	6E-6	2E-8	8E-5	8E-4
56	Barium-128	D, all compounds	5E+2	2E+3	7E-7	2E-9	7E-6	7E-5
56	Barium-131m ²	D, all compounds	4E+5	1E+6	6E-4	2E-6	-	-
			St wall (5E+5)	-	-	-	7E-3	7E-2
56	Barium-131	D, all compounds	3E+3	8E+3	3E-6	1E-8	4E-5	4E-4
56	Barium-133m	D, all compounds	2E+3	9E+3	4E-6	1E-8	-	-
			LLI wall (3E+3)	-	-	-	4E-5	4E-4
56	Barium-133	D, all compounds	2E+3	7E+2	3E-7	9E-10	2E-5	2E-4
56	Barium-135m	D, all compounds	3E+3	1E+4	5E-6	2E-8	4E-5	4E-4
56	Barium-139 ²	D, all compounds	1E+4	3E+4	1E-5	4E-8	2E-4	2E-3
56	Barium-140	D, all compounds	5E+2	1E+3	6E-7	2E-9	-	-
			LLI wall (6E+2)	-	-	-	8E-6	8E-5
56	Barium-141 ²	D, all compounds	2E+4	7E+4	3E-5	1E-7	3E-4	3E-3
56	Barium-142 ²	D, all compounds	5E+4	1E+5	6E-5	2E-7	7E-4	7E-3
57	Lanthanum-131 ²	D, all compounds except those given for W	5E+4	1E+5	5E-5	2E-7	6E-4	6E-3
		W, oxides and hydroxides	-	2E+5	7E-5	2E-7	-	-
57	Lanthanum-132	D, see ¹³¹ La	3E+3	1E+4	4E-6	1E-8	4E-5	4E-4
		W, see ¹³¹ La	-	1E+4	5E-6	2E-8	-	-
57	Lanthanum-135	D, see ¹³¹ La	4E+4	1E+5	4E-5	1E-7	5E-4	5E-3
		W, see ¹³¹ La	-	9E+4	4E-5	1E-7	-	-
57	Lanthanum-137	D, see ¹³¹ La	1E+4	6E+1	3E-8	-	2E-4	2E-3
			-	Liver (7E+1)	-	1E-10	-	-
		W, see ¹³¹ La	-	3E+2	1E-7	-	-	-
			-	Liver (3E+2)	-	4E-10	-	-
57	Lanthanum-138	D, see ¹³¹ La	9E+2	4E+0	1E-9	5E-12	1E-5	1E-4
		W, see ¹³¹ La	-	1E+1	6E-9	2E-11	-	-
57	Lanthanum-140	D, see ¹³¹ La	6E+2	1E+3	6E-7	2E-9	9E-6	9E-5
		W, see ¹³¹ La	-	1E+3	5E-7	2E-9	-	-
57	Lanthanum-141	D, see ¹³¹ La	4E+3	9E+3	4E-6	1E-8	5E-5	5E-4
		W, see ¹³¹ La	-	1E+4	5E-6	2E-8	-	-
57	Lanthanum-142 ²	D, see ¹³¹ La	8E+3	2E+4	9E-6	3E-8	1E-4	1E-3
		W, see ¹³¹ La	-	3E+4	1E-5	5E-8	-	-
57	Lanthanum-143 ²	D, see ¹³¹ La	4E+4	1E+5	4E-5	1E-7	-	-
			St wall (4E+4)	-	-	-	5E-4	5E-3
		W, see ¹³¹ La	-	9E+4	4E-5	1E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
58	Cerium-134	W, all compounds except those given for Y	5E+2	7E+2	3E-7	1E-9	-	-
			LLI wall (6E+2)	-	-	-	8E-6	8E-5
58	Cerium-135	Y, oxides, hydroxides, and fluorides	-	7E+2	3E-7	9E-10	-	-
			W, see ¹³⁴ Ce	2E+3	4E+3	2E-6	5E-9	2E-5
58	Cerium-137m	Y, see ¹³⁴ Ce	-	4E+3	1E-6	5E-9	-	-
			W, see ¹³⁴ Ce	2E+3	4E+3	2E-6	6E-9	-
58	Cerium-137	Y, see ¹³⁴ Ce	LLI wall (2E+3)	-	-	-	3E-5	3E-4
			W, see ¹³⁴ Ce	5E+4	1E+5	6E-5	2E-7	7E-4
58	Cerium-139	Y, see ¹³⁴ Ce	-	4E+3	2E-6	5E-9	-	-
			W, see ¹³⁴ Ce	5E+3	8E+2	3E-7	1E-9	7E-5
58	Cerium-141	Y, see ¹³⁴ Ce	-	7E+2	3E-7	9E-10	-	-
			W, see ¹³⁴ Ce	2E+3	7E+2	3E-7	1E-9	-
58	Cerium-143	Y, see ¹³⁴ Ce	LLI wall (2E+3)	-	-	-	3E-5	3E-4
			W, see ¹³⁴ Ce	1E+3	2E+3	8E-7	3E-9	-
58	Cerium-144	Y, see ¹³⁴ Ce	-	6E+2	2E-7	8E-10	-	-
			W, see ¹³⁴ Ce	2E+2	3E+1	1E-8	4E-11	-
59	Praseodymium-136 ²	Y, see ¹³⁴ Ce	LLI wall (1E+3)	-	-	-	2E-5	2E-4
			W, see ¹³⁴ Ce	2E+2	3E+1	1E-8	4E-11	-
59	Praseodymium-137 ²	Y, see ¹³⁴ Ce	-	1E+1	6E-9	2E-11	-	-
			W, all compounds except those given for Y	5E+4	2E+5	1E-4	3E-7	-
59	Praseodymium-138m	Y, oxides, hydroxides, carbides, and fluorides	St wall (7E+4)	-	-	-	1E-3	1E-2
			W, see ¹³⁶ Pr	4E+4	2E+5	6E-5	2E-7	5E-4
59	Praseodymium-139	Y, see ¹³⁶ Pr	-	2E+5	9E-5	3E-7	-	-
			W, see ¹³⁶ Pr	1E+4	5E+4	2E-5	8E-8	1E-4
59	Praseodymium-142m ²	Y, see ¹³⁶ Pr	-	1E+5	6E-5	2E-7	-	-
			W, see ¹³⁶ Pr	4E+4	1E+5	5E-5	2E-7	6E-4
59	Praseodymium-142	Y, see ¹³⁶ Pr	-	1E+5	5E-5	2E-7	-	-
			W, see ¹³⁶ Pr	8E+4	2E+5	7E-5	2E-7	1E-3
59	Praseodymium-143	Y, see ¹³⁶ Pr	-	1E+5	6E-5	2E-7	-	-
			W, see ¹³⁶ Pr	1E+3	2E+3	9E-7	3E-9	1E-5
59	Praseodymium-144 ²	Y, see ¹³⁶ Pr	-	2E+3	8E-7	3E-9	-	-
			W, see ¹³⁶ Pr	9E+2	8E+2	3E-7	1E-9	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
			St wall (4E+4)	-	-	-	6E-4	6E-3
59	Praseodymium-145	Y, see ¹³⁶ Pr	-	1E+5	5E-5	2E-7	-	-
		W, see ¹³⁶ Pr	3E+3	9E+3	4E-6	1E-8	4E-5	4E-4
59	Praseodymium-147 ²	Y, see ¹³⁶ Pr	-	8E+3	3E-6	1E-8	-	-
		W, see ¹³⁶ Pr	5E+4	2E+5	8E-5	3E-7	-	-
			St wall (8E+4)	-	-	-	1E-3	1E-2
60	Neodymium-136 ²	Y, see ¹³⁶ Pr	-	2E+5	8E-5	3E-7	-	-
		W, all compounds except those given for Y	1E+4	6E+4	2E-5	8E-8	2E-4	2E-3
60	Neodymium-138	Y, oxides, hydroxides, carbides, and fluorides	-	5E+4	2E-5	8E-8	-	-
		W, see ¹³⁶ Nd	2E+3	6E+3	3E-6	9E-9	3E-5	3E-4
60	Neodymium-139m	Y, see ¹³⁶ Nd	-	5E+3	2E-6	7E-9	-	-
		W, see ¹³⁶ Nd	5E+3	2E+4	7E-6	2E-8	7E-5	7E-4
60	Neodymium-139 ²	Y, see ¹³⁶ Nd	-	1E+4	6E-6	2E-8	-	-
		W, see ¹³⁶ Nd	9E+4	3E+5	1E-4	5E-7	1E-3	1E-2
60	Neodymium-141	Y, see ¹³⁶ Nd	-	3E+5	1E-4	4E-7	-	-
		W, see ¹³⁶ Nd	2E+5	7E+5	3E-4	1E-6	2E-3	2E-2
60	Neodymium-147	Y, see ¹³⁶ Nd	-	6E+5	3E-4	9E-7	-	-
		W, see ¹³⁶ Nd	1E+3	9E+2	4E-7	1E-9	-	-
			LLI wall (1E+3)	-	-	-	2E-5	2E-4
60	Neodymium-149 ²	Y, see ¹³⁶ Nd	-	8E+2	4E-7	1E-9	-	-
		W, see ¹³⁶ Nd	1E+4	3E+4	1E-5	4E-8	1E-4	1E-3
60	Neodymium-151 ²	Y, see ¹³⁶ Nd	-	2E+4	1E-5	3E-8	-	-
		W, see ¹³⁶ Nd	7E+4	2E+5	8E-5	3E-7	9E-4	9E-3
61	Promethium-141 ²	Y, see ¹³⁶ Nd	-	2E+5	8E-5	3E-7	-	-
		W, all compounds except those given for Y	5E+4	2E+5	8E-5	3E-7	-	-
			St wall (6E+4)	-	-	-	8E-4	8E-3
61	Promethium-143	Y, oxides, hydroxides, carbides, and fluorides	-	2E+5	7E-5	2E-7	-	-
		W, see ¹⁴¹ Pm	5E+3	6E+2	2E-7	8E-10	7E-5	7E-4
61	Promethium-144	Y, see ¹⁴¹ Pm	-	7E+2	3E-7	1E-9	-	-
		W, see ¹⁴¹ Pm	1E+3	1E+2	5E-8	2E-10	2E-5	2E-4
61	Promethium-145	Y, see ¹⁴¹ Pm	-	1E+2	5E-8	2E-10	-	-
		W, see ¹⁴¹ Pm	1E+4	2E+2	7E-8	-	1E-4	1E-3
			-	Bone surf (2E+2)	-	3E-10	-	-
61	Promethium-146	Y, see ¹⁴¹ Pm	-	2E+2	8E-8	3E-10	-	-
		W, see ¹⁴¹ Pm	2E+3	5E+1	2E-8	7E-11	2E-5	2E-4
61	Promethium-147	Y, see ¹⁴¹ Pm	-	4E+1	2E-8	6E-11	-	-
		W, see ¹⁴¹ Pm	4E+3	1E+2	5E-8	-	-	-
			LLI wall (5E+3)	Bone surf (2E+2)	-	3E-10	7E-5	7E-4
		Y, see ¹⁴¹ Pm	-	1E+2	6E-8	2E-10	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
61	Promethium-148m	W, see ¹⁴¹ Pm	7E+2	3E+2	1E-7	4E-10	1E-5	1E-4
		Y, see ¹⁴¹ Pm	-	3E+2	1E-7	5E-10	-	-
61	Promethium-148	W, see ¹⁴¹ Pm	4E+2	5E+2	2E-7	8E-10	-	-
			LLI wall (5E+2)	-	-	-	7E-6	7E-5
		Y, see ¹⁴¹ Pm	-	5E+2	2E-7	7E-10	-	-
61	Promethium-149	W, see ¹⁴¹ Pm	1E+3	2E+3	8E-7	3E-9	-	-
			LLI wall (1E+3)	-	-	-	2E-5	2E-4
		Y, see ¹⁴¹ Pm	-	2E+3	8E-7	2E-9	-	-
61	Promethium-150	W, see ¹⁴¹ Pm	5E+3	2E+4	8E-6	3E-8	7E-5	7E-4
		Y, see ¹⁴¹ Pm	-	2E+4	7E-6	2E-8	-	-
61	Promethium-151	W, see ¹⁴¹ Pm	2E+3	4E+3	1E-6	5E-9	2E-5	2E-4
		Y, see ¹⁴¹ Pm	-	3E+3	1E-6	4E-9	-	-
62	Samarium-141m ²	W, all compounds	3E+4	1E+5	4E-5	1E-7	4E-4	4E-3
62	Samarium-141 ²	W, all compounds	5E+4	2E+5	8E-5	2E-7	-	-
			St wall (6E+4)	-	-	-	8E-4	8E-3
62	Samarium-142 ²	W, all compounds	8E+3	3E+4	1E-5	4E-8	1E-4	1E-3
62	Samarium-145	W, all compounds	6E+3	5E+2	2E-7	7E-10	8E-5	8E-4
62	Samarium-146	W, all compounds	1E+1	4E-2	1E-11	-	-	-
			Bone surf (3E+1)	Bone surf (6E-2)	-	9E-14	3E-7	3E-6
62	Samarium-147	W, all compounds	2E+1	4E-2	2E-11	-	-	-
			Bone surf (3E+1)	Bone surf (7E-2)	-	1E-13	4E-7	4E-6
62	Samarium-151	W, all compounds	1E+4	1E+2	4E-8	-	-	-
			LLI wall (1E+4)	Bone surf (2E+2)	-	2E-10	2E-4	2E-3
62	Samarium-153	W, all compounds	2E+3	3E+3	1E-6	4E-9	-	-
			LLI wall (2E+3)	-	-	-	3E-5	3E-4
62	Samarium-155 ²	W, all compounds	6E+4	2E+5	9E-5	3E-7	-	-
			St wall (8E+4)	-	-	-	1E-3	1E-2
62	Samarium-156	W, all compounds	5E+3	9E+3	4E-6	1E-8	7E-5	7E-4
63	Europium-145	W, all compounds	2E+3	2E+3	8E-7	3E-9	2E-5	2E-4
63	Europium-146	W, all compounds	1E+3	1E+3	5E-7	2E-9	1E-5	1E-4
63	Europium-147	W, all compounds	3E+3	2E+3	7E-7	2E-9	4E-5	4E-4
63	Europium-148	W, all compounds	1E+3	4E+2	1E-7	5E-10	1E-5	1E-4
63	Europium-149	W, all compounds	1E+4	3E+3	1E-6	4E-9	2E-4	2E-3
63	Europium-150 (12.62h)	W, all compounds	3E+3	8E+3	4E-6	1E-8	4E-5	4E-4
63	Europium-150 (34.2 y)	W, all compounds	8E+2	2E+1	8E-9	3E-11	1E-5	1E-4
63	Europium-152m	W, all compounds	3E+3	6E+3	3E-6	9E-9	4E-5	4E-4
63	Europium-152	W, all compounds	8E+2	2E+1	1E-8	3E-11	1E-5	1E-4
63	Europium-154	W, all compounds	5E+2	2E+1	8E-9	3E-11	7E-6	7E-5
63	Europium-155	W, all compounds	4E+3	9E+1	4E-8	-	5E-5	5E-4
			-	Bone surf (1E+2)	-	2E-10	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
63	Europium-156	W, all compounds	6E+2	5E+2	2E-7	6E-10	8E-6	8E-5
63	Europium-157	W, all compounds	2E+3	5E+3	2E-6	7E-9	3E-5	3E-4
63	Europium-158 ²	W, all compounds	2E+4	6E+4	2E-5	8E-8	3E-4	3E-3
64	Gadolinium-145 ²	D, all compounds except those given for W	5E+4	2E+5	6E-5	2E-7	-	-
		St wall	(5E+4)	-	-	-	6E-4	6E-3
		W, oxides, hydroxides, and fluorides	-	2E+5	7E-5	2E-7	-	-
64	Gadolinium-146	D, see ¹⁴⁵ Gd	1E+3	1E+2	5E-8	2E-10	2E-5	2E-4
		W, see ¹⁴⁵ Gd	-	3E+2	1E-7	4E-10	-	-
64	Gadolinium-147	D, see ¹⁴⁵ Gd	2E+3	4E+3	2E-6	6E-9	3E-5	3E-4
		W, see ¹⁴⁵ Gd	-	4E+3	1E-6	5E-9	-	-
64	Gadolinium-148	D, see ¹⁴⁵ Gd	1E+1	8E+3	3E-12	-	-	-
		Bone surf (2E+1)	-	Bone surf (2E+2)	-	2E-14	3E-7	3E-6
		W, see ¹⁴⁵ Gd	-	3E-2	1E-11	-	-	-
			-	Bone surf (6E-2)	-	8E-14	-	-
64	Gadolinium-149	D, see ¹⁴⁵ Gd	3E+3	2E+3	9E-7	3E-9	4E-5	4E-4
		W, see ¹⁴⁵ Gd	-	2E+3	1E-6	3E-9	-	-
64	Gadolinium-151	D, see ¹⁴⁵ Gd	6E+3	4E+2	2E-7	-	9E-5	9E-4
			-	Bone surf (6E+2)	-	9E-10	-	-
		W, see ¹⁴⁵ Gd	-	1E+3	5E-7	2E-9	-	-
64	Gadolinium-152	D, see ¹⁴⁵ Gd	2E+1	1E-2	4E-12	-	-	-
			Bone surf (3E+1)	Bone surf (2E-2)	-	3E-14	4E-7	4E-6
		W, see ¹⁴⁵ Gd	-	4E-2	2E-11	-	-	-
			-	Bone surf (8E-2)	-	1E-13	-	-
64	Gadolinium-153	D, see ¹⁴⁵ Gd	5E+3	1E+2	6E-8	-	6E-5	6E-4
			-	Bone surf (2E+2)	-	3E-10	-	-
		W, see ¹⁴⁵ Gd	-	6E+2	2E-7	8E-10	-	-
64	Gadolinium-159	D, see ¹⁴⁵ Gd	3E+3	8E+3	3E-6	1E-8	4E-5	4E-4
		W, see ¹⁴⁵ Gd	-	6E+3	2E-6	8E-9	-	-
65	Terbium-147 ²	W, all compounds	9E+3	3E+4	1E-5	5E-8	1E-4	1E-3
65	Terbium-149	W, all compounds	5E+3	7E+2	3E-7	1E-9	7E-5	7E-4
65	Terbium-150	W, all compounds	5E+3	2E+4	9E-6	3E-8	7E-5	7E-4
65	Terbium-151	W, all compounds	4E+3	9E+3	4E-6	1E-8	5E-5	5E-4
65	Terbium-153	W, all compounds	5E+3	7E+3	3E-6	1E-8	7E-5	7E-4
65	Terbium-154	W, all compounds	2E+3	4E+3	2E-6	6E-9	2E-5	2E-4
65	Terbium-155	W, all compounds	6E+3	8E+3	3E-6	1E-8	8E-5	8E-4
65	Terbium-156m (5.0 h)	W, all compounds	2E+4	3E+4	1E-5	4E-8	2E-4	2E-3
65	Terbium-156m (24.4 h)	W, all compounds	7E+3	8E+3	3E-6	1E-8	1E-4	1E-3
65	Terbium-156	W, all compounds	1E+3	1E+3	6E-7	2E-9	1E-5	1E-4
65	Terbium-157	W, all compounds	5E+4	3E+2	1E-7	-	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
			LLI wall (5E+4)	Bone surf (6E+2)	-	8E-10	7E-4	7E-3
65	Terbium-158	W, all compounds	1E+3	2E+1	8E-9	3E-11	2E-5	2E-4
65	Terbium-160	W, all compounds	8E+2	2E+2	9E-8	3E-10	1E-5	1E-4
65	Terbium-161	W, all compounds	2E+3	2E+3	7E-7	2E-9	-	-
			LLI wall (2E+3)	-	-	-	3E-5	3E-4
66	Dysprosium-155	W, all compounds	9E+3	3E+4	1E-5	4E-8	1E-4	1E-3
66	Dysprosium-157	W, all compounds	2E+4	6E+4	3E-5	9E-8	3E-4	3E-3
66	Dysprosium-159	W, all compounds	1E+4	2E+3	1E-6	3E-9	2E-4	2E-3
66	Dysprosium-165	W, all compounds	1E+4	5E+4	2E-5	6E-8	2E-4	2E-3
66	Dysprosium-166	W, all compounds	6E+2	7E+2	3E-7	1E-9	-	-
			LLI wall (8E+2)	-	-	-	1E-5	1E-4
67	Holmium-155 ²	W, all compounds	4E+4	2E+5	6E-5	2E-7	6E-4	6E-3
67	Holmium-157 ²	W, all compounds	3E+5	1E+6	6E-4	2E-6	4E-3	4E-2
67	Holmium-159 ²	W, all compounds	2E+5	1E+6	4E-4	1E-6	3E-3	3E-2
67	Holmium-161	W, all compounds	1E+5	4E+5	2E-4	6E-7	1E-3	1E-2
67	Holmium-162m ²	W, all compounds	5E+4	3E+5	1E-4	4E-7	7E-4	7E-3
67	Holmium-162 ²	W, all compounds	5E+5	2E+6	1E-3	3E-6	-	-
			St wall (8E+5)	-	-	-	1E-2	1E-1
67	Holmium-164m ²	W, all compounds	1E+5	3E+5	1E-4	4E-7	1E-3	1E-2
67	Holmium-164 ²	W, all compounds	2E+5	6E+5	3E-4	9E-7	-	-
			St wall (2E+5)	-	-	-	3E-3	3E-2
67	Holmium-166m	W, all compounds	6E+2	7E+0	3E-9	9E-12	9E-6	9E-5
67	Holmium-166	W, all compounds	9E+2	2E+3	7E-7	2E-9	-	-
			LLI wall (9E+2)	-	-	-	1E-5	1E-4
67	Holmium-167	W, all compounds	2E+4	6E+4	2E-5	8E-8	2E-4	2E-3
68	Erbium-161	W, all compounds	2E+4	6E+4	3E-5	9E-8	2E-4	2E-3
68	Erbium-165	W, all compounds	6E+4	2E+5	8E-5	3E-7	9E-4	9E-3
68	Erbium-169	W, all compounds	3E+3	3E+3	1E-6	4E-9	-	-
			LLI wall (4E+3)	-	-	-	5E-5	5E-4
68	Erbium-171	W, all compounds	4E+3	1E+4	4E-6	1E-8	5E-5	5E-4
68	Erbium-172	W, all compounds	1E+3	1E+3	6E-7	2E-9	-	-
			LLI wall (E+3)	-	-	-	2E-5	2E-4
69	Thulium-162 ²	W, all compounds	7E+4	3E+5	1E-4	4E-7	-	-
			St wall (7E+4)	-	-	-	1E-3	1E-2
69	Thulium-166	W, all compounds	4E+3	1E+4	6E-6	2E-8	6E-5	6E-4
69	Thulium-167	W, all compounds	2E+3	2E+3	8E-7	3E-9	-	-
			LLI wall (2E+3)	-	-	-	3E-5	3E-4
69	Thulium-170	W, all compounds	8E+2	2E+2	9E-8	3E-10	-	-
			LLI wall (1E+3)	-	-	-	1E-5	1E-4
69	Thulium-171	W, all compounds	1E+4	3E+2	1E-7	-	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
69	Thulium-172	W, all compounds	LLI wall (1E+4)	Bone surf (6E+2)	-	8E-10	2E-4	2E-3
			7E+2	1E+3	5E-7	2E-9	-	-
69	Thulium-173	W, all compounds	LLI wall (8E+2)	-	-	-	1E-5	1E-4
			4E+3	1E+4	5E-6	2E-8	6E-5	6E-4
69	Thulium-175 ²	W, all compounds	7E+4	3E+5	1E-4	4E-7	-	-
			St wall (9E+4)	-	-	-	1E-3	1E-2
70	Ytterbium-162 ²	W, all compounds except those given for Y Y, oxides, hydroxides, and fluorides	7E+4	3E+5	1E-4	4E-7	1E-3	1E-2
			-	3E+5	1E-4	4E-7	-	-
70	Ytterbium-166	W, see ¹⁶² Yb Y, see ¹⁶² Yb	1E+3	2E+3	8E-7	3E-9	2E-5	2E-4
			-	2E+3	8E-7	3E-9	-	-
70	Ytterbium-167 ²	W, see ¹⁶² Yb Y, see ¹⁶² Yb	3E+5	8E+5	3E-4	1E-6	4E-3	4E-2
			-	7E+5	3E-4	1E-6	-	-
70	Ytterbium-169	W, see ¹⁶² Yb Y, see ¹⁶² Yb	2E+3	8E+2	4E-7	1E-9	2E-5	2E-4
			-	7E+2	3E-7	1E-9	-	-
70	Ytterbium-175	W, see ¹⁶² Yb Y, see ¹⁶² Yb	3E+3	4E+3	1E-6	5E-9	-	-
			LLI wall (3E+3)	-	-	-	4E-5	4E-4
70	Ytterbium-177 ²	W, see ¹⁶² Yb Y, see ¹⁶² Yb	2E+4	5E+4	2E-5	7E-8	2E-4	2E-3
			-	5E+4	2E-5	6E-8	-	-
70	Ytterbium-178 ²	W, see ¹⁶² Yb Y, see ¹⁶² Yb	1E+4	4E+4	2E-5	6E-8	2E-4	2E-3
			-	4E+4	2E-5	5E-8	-	-
71	Lutetium-169	W, all compounds except those given for Y Y, oxides, hydroxides, and fluorides	3E+3	4E+3	2E-6	6E-9	3E-5	3E-4
			-	4E+3	2E-6	6E-9	-	-
71	Lutetium-170	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	1E+3	2E+3	9E-7	3E-9	2E-5	2E-4
			-	2E+3	8E-7	3E-9	-	-
71	Lutetium-171	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	2E+3	2E+3	8E-7	3E-9	3E-5	3E-4
			-	2E+3	8E-7	3E-9	-	-
71	Lutetium-172	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	1E+3	1E+3	5E-7	2E-9	1E-5	1E-4
			-	1E+3	5E-7	2E-9	-	-
71	Lutetium-173	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	5E+3	3E+2	1E-7	-	7E-5	7E-4
			-	Bone surf (5E+2)	-	6E-10	-	-
71	Lutetium-174m	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	-	3E+2	1E-7	4E-10	-	-
			2E+3	2E+2	1E-7	-	-	-
71	Lutetium-174	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	LLI wall (3E+3)	Bone surf (3E+2)	-	5E-10	4E-5	4E-4
			-	2E+2	9E-8	3E-10	-	-
71	Lutetium-176m	W, see ¹⁶⁹ Lu Y, see ¹⁶⁹ Lu	5E+3	1E+2	5E-8	-	7E-5	7E-4
			-	Bone surf (2E+2)	-	3E-10	-	-
			-	2E+2	6E-8	2E-10	-	-
			8E+3	3E+4	1E-5	3E-8	1E-4	1E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
71	Lutetium-176	Y, see ^{169}Lu	-	2E+4	9E-6	3E-8	-	-
		W, see ^{169}Lu	7E+2	5E+0	2E-9	-	1E-5	1E-4
71	Lutetium-177m	Y, see ^{169}Lu	-	Bone surf (1E+1)	-	2E-11	-	-
		W, see ^{169}Lu	7E+2	8E+0	3E-9	1E-11	-	-
71	Lutetium-177	Y, see ^{169}Lu	-	1E+2	5E-8	-	1E-5	1E-4
		W, see ^{169}Lu	2E+3	8E+1	3E-8	1E-10	-	-
71	Lutetium-178m ²	Y, see ^{169}Lu	-	2E+3	9E-7	3E-9	-	-
		W, see ^{169}Lu	5E+4	2E+5	8E-5	3E-7	-	-
71	Lutetium-178 ²	Y, see ^{169}Lu	-	St. wall (6E+4)	-	-	4E-5	4E-4
		W, see ^{169}Lu	4E+4	2E+3	9E-7	3E-9	-	-
71	Lutetium-179	Y, see ^{169}Lu	-	2E+5	7E-5	2E-7	-	-
		W, see ^{169}Lu	6E+3	1E+5	5E-5	2E-7	-	-
72	Hafnium-170	Y, see ^{169}Lu	-	1E+5	5E-5	2E-7	-	-
		D, all compounds except those given for W	3E+3	2E+4	8E-6	3E-8	9E-5	9E-4
72	Hafnium-172	W, oxides, hydroxides, carbides, and nitrates	-	2E+4	6E-6	3E-8	-	-
		D, see ^{170}Hf	1E+3	6E+3	2E-6	8E-9	4E-5	4E-4
72	Hafnium-173	W, see ^{170}Hf	-	5E+3	2E-6	6E-9	-	-
		D, see ^{170}Hf	5E+3	9E+0	4E-9	-	2E-5	2E-4
72	Hafnium-175	W, see ^{170}Hf	-	Bone surf (2E+1)	-	3E-11	-	-
		D, see ^{170}Hf	3E+3	4E+1	2E-8	-	-	-
72	Hafnium-177m ²	W, see ^{170}Hf	-	Bone surf (6E+1)	-	8E-11	-	-
		D, see ^{170}Hf	2E+4	1E+4	5E-6	2E-8	7E-5	7E-4
72	Hafnium-178m	W, see ^{170}Hf	-	1E+4	5E-6	2E-8	-	-
		D, see ^{170}Hf	3E+2	9E+2	4E-7	-	4E-5	4E-4
72	Hafnium-179m	W, see ^{170}Hf	-	Bone surf (1E+3)	-	1E-9	-	-
		D, see ^{170}Hf	1E+3	1E+3	5E-7	2E-9	-	-
72	Hafnium-179m	W, see ^{170}Hf	-	2E+4	2E-5	8E-8	3E-4	3E-3
		D, see ^{170}Hf	3E+2	9E+4	4E-5	1E-7	-	-
72	Hafnium-179m	W, see ^{170}Hf	-	Bone surf (2E+0)	-	3E-12	-	-
		D, see ^{170}Hf	1E+3	5E+0	2E-9	-	-	-
72	Hafnium-179m	W, see ^{170}Hf	-	Bone surf (9E+0)	-	1E-11	-	-
		D, see ^{170}Hf	1E+3	3E+2	1E-7	-	1E-5	1E-4
72	Hafnium-179m	W, see ^{170}Hf	-	Bone surf (6E+2)	-	8E-10	-	-
		D, see ^{170}Hf	1E+3	3E+2	1E-7	-	1E-5	1E-4

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
72	Hafnium-180m	W, see ¹⁷⁰ Hf	-	6E+2	3E-7	8E-10	-	-
		D, see ¹⁷⁰ Hf	7E+3	2E+4	9E-6	3E-8	1E-4	1E-3
72	Hafnium-181	W, see ¹⁷⁰ Hf	-	3E+4	1E-5	4E-8	-	-
		D, see ¹⁷⁰ Hf	1E+3	2E+2	7E-8	-	2E-5	2E-4
72	Hafnium-182m ²		-	Bone surf (4E+2)	-	6E-10	-	-
		W, see ¹⁷⁰ Hf	-	4E+2	2E-7	6E-10	-	-
		D, see ¹⁷⁰ Hf	4E+4	9E+4	4E-5	1E-7	5E-4	5E-3
72	Hafnium-182	W, see ¹⁷⁰ Hf	-	1E+5	6E-5	2E-7	-	-
		D, see ¹⁷⁰ Hf	2E+2	8E-1	3E-10	-	-	-
72	Hafnium-183 ²		-	Bone surf (4E+2)	-	2E-12	5E-6	5E-5
		W, see ¹⁷⁰ Hf	-	3E+0	1E-9	-	-	-
		D, see ¹⁷⁰ Hf	2E+4	5E+4	2E-5	6E-8	3E-4	3E-3
72	Hafnium-184	W, see ¹⁷⁰ Hf	-	6E+4	2E-5	8E-8	-	-
		D, see ¹⁷⁰ Hf	2E+3	8E+3	3E-6	1E-8	3E-5	3E-4
73	Tantalum-172 ²	W, all compounds except those given for Y	4E+4	1E+5	5E-5	2E-7	5E-4	5E-3
		Y, elemental Ta, oxides, hydroxides, halides, carbides, nitrates, and nitrides	-	1E+5	4E-5	1E-7	-	-
73	Tantalum-173	W, see ¹⁷² Ta	7E+3	2E+4	8E-6	3E-8	9E-5	9E-4
		Y, see ¹⁷² Ta	-	2E+4	7E-6	2E-8	-	-
73	Tantalum-174 ²	W, see ¹⁷² Ta	3E+4	1E+5	4E-5	1E-7	4E-4	4E-3
		Y, see ¹⁷² Ta	-	9E+4	4E-5	1E-7	-	-
73	Tantalum-175	W, see ¹⁷² Ta	6E+3	2E+4	7E-6	2E-8	8E-5	8E-4
		Y, see ¹⁷² Ta	-	1E+4	6E-6	2E-8	-	-
73	Tantalum-176	W, see ¹⁷² Ta	4E+3	1E+4	5E-6	2E-8	5E-5	5E-4
		Y, see ¹⁷² Ta	-	1E+4	5E-6	2E-8	-	-
73	Tantalum-177	W, see ¹⁷² Ta	1E+4	2E+4	8E-6	3E-8	2E-4	2E-3
		Y, see ¹⁷² Ta	-	2E+4	7E-6	2E-8	-	-
73	Tantalum-178	W, see ¹⁷² Ta	2E+4	9E+4	4E-5	1E-7	2E-4	2E-3
		Y, see ¹⁷² Ta	-	7E+4	3E-5	1E-7	-	-
73	Tantalum-179	W, see ¹⁷² Ta	2E+4	5E+3	2E-6	8E-9	3E-4	3E-3
		Y, see ¹⁷² Ta	-	9E+2	4E-7	1E-9	-	-
73	Tantalum-180m	W, see ¹⁷² Ta	2E+4	7E+4	3E-5	9E-8	3E-4	3E-3
		Y, see ¹⁷² Ta	-	6E+4	2E-5	8E-8	-	-
73	Tantalum-180	W, see ¹⁷² Ta	1E+3	4E+2	2E-7	6E-10	2E-5	2E-4
		Y, see ¹⁷² Ta	-	2E+1	1E-8	3E-11	-	-
73	Tantalum-182m ²	W, see ¹⁷² Ta	2E+5	5E+5	2E-4	8E-7	-	-
			St wall (2E+5)	-	-	-	3E-3	3E-2
		Y, see ¹⁷² Ta	-	4E+5	2E-4	6E-7	-	-
73	Tantalum-182	W, see ¹⁷² Ta	8E+2	3E+2	1E-7	5E-10	1E-5	1E-4

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
73	Tantalum-183	Y, see ^{172}Ta	-	1E+2	6E-8	2E-10	-	-
		W, see ^{172}Ta	9E+2	1E+3	5E-7	2E-9	-	-
73	Tantalum-184	LLI wall (1E+3)	-	-	-	-	2E-5	2E-4
		Y, see ^{172}Ta	-	1E+3	4E-7	1E-9	-	-
73	Tantalum-185 ²	W, see ^{172}Ta	2E+3	5E+3	2E-6	8E-9	3E-5	3E-4
		Y, see ^{172}Ta	-	5E+3	2E-6	7E-9	-	-
73	Tantalum-186 ²	W, see ^{172}Ta	3E+4	7E+4	3E-5	1E-7	4E-4	4E-3
		Y, see ^{172}Ta	-	6E+4	3E-5	9E-8	-	-
74	Tungsten-176	W, see ^{172}Ta	5E+4	2E+5	1E-4	3E-7	-	-
		St wall (7E+4)	-	-	-	-	1E-3	1E-2
74	Tungsten-177	Y, see ^{172}Ta	-	2E+5	9E-5	3E-7	-	-
		D, all compounds	1E+4	5E+4	2E-5	7E-8	1E-4	1E-3
74	Tungsten-178	D, all compounds	2E+4	9E+4	4E-5	1E-7	3E-4	3E-3
74	Tungsten-179 ²	D, all compounds	5E+3	2E+4	8E-6	3E-8	7E-5	7E-4
74	Tungsten-181	D, all compounds	5E+5	2E+6	7E-4	2E-6	7E-3	7E-2
74	Tungsten-185	D, all compounds	2E+4	3E+4	1E-5	5E-8	2E-4	2E-3
		LLI wall (3E+3)	-	-	-	-	4E-5	4E-4
74	Tungsten-187	D, all compounds	2E+3	9E+3	4E-6	1E-8	3E-5	3E-4
74	Tungsten-188	D, all compounds	4E+2	1E+3	5E-7	2E-9	-	-
		LLI wall (5E+2)	-	-	-	-	7E-6	7E-5
75	Rhenium-177 ²	D, all compounds except those given for W	9E+4	3E+5	1E-4	4E-7	-	-
		St wall (1E+5)	-	-	-	-	2E-3	2E-2
75	Rhenium-178 ²	W, oxides, hydroxides, and nitrates	-	4E+5	1E-4	5E-7	-	-
		D, see ^{177}Re	7E+4	3E+5	1E-4	4E-7	-	-
75	Rhenium-181	St wall (1E+5)	-	-	-	-	1E-3	1E-2
		W, see ^{177}Re	-	3E+5	1E-4	4E-7	-	-
75	Rhenium-182 (12.7 h)	D, see ^{177}Re	5E+3	9E+3	4E-6	1E-8	7E-5	7E-4
		W, see ^{177}Re	-	9E+3	4E-6	1E-8	-	-
75	Rhenium-182 (64.0 h)	D, see ^{177}Re	7E+3	1E+4	5E-6	2E-8	9E-5	9E-4
		W, see ^{177}Re	-	2E+4	6E-6	2E-8	-	-
75	Rhenium-184m	D, see ^{177}Re	1E+3	2E+3	1E-6	3E-9	2E-5	2E-4
		W, see ^{177}Re	-	2E+3	9E-7	3E-9	-	-
75	Rhenium-184	D, see ^{177}Re	2E+3	3E+3	1E-6	4E-9	3E-5	3E-4
		W, see ^{177}Re	-	4E+2	2E-7	6E-10	-	-
75	Rhenium-186m	D, see ^{177}Re	2E+3	4E+3	1E-6	5E-9	3E-5	3E-4
		W, see ^{177}Re	-	1E+3	6E-7	2E-9	-	-
75	Rhenium-186m	D, see ^{177}Re	1E+3	2E+3	7E-7	-	-	-
		St wall (2E+3)	-	St wall (2E+3)	-	3E-9	2E-5	2E-4
		W, see ^{177}Re	-	2E+2	6E-8	2E-10	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
75	Rhenium-186	D, see ¹⁷⁷ Re	2E+3	3E+3	1E-6	4E-9	3E-5	3E-4
		W, see ¹⁷⁷ Re	-	2E+3	7E-7	2E-9	-	-
75	Rhenium-187	D, see ¹⁷⁷ Re	6E+5	8E+5	4E-4	-	8E-3	8E-2
			-	St wall (9E+5)	-	1E-6	-	-
		W, see ¹⁷⁷ Re	-	1E+5	4E-5	1E-7	-	-
75	Rhenium-188m ²	D, see ¹⁷⁷ Re	8E+4	1E+5	6E-5	2E-7	1E-3	1E-2
		W, see ¹⁷⁷ Re	-	1E+5	6E-5	2E-7	-	-
75	Rhenium-188	D, see ¹⁷⁷ Re	2E+3	3E+3	1E-6	4E-9	2E-5	2E-4
		W, see ¹⁷⁷ Re	-	3E+3	1E-6	4E-9	-	-
75	Rhenium-189	D, see ¹⁷⁷ Re	3E+3	5E+3	2E-6	7E-9	4E-5	4E-4
		W, see ¹⁷⁷ Re	-	4E+3	2E-6	6E-9	-	-
76	Osmium-180 ²	D, all compounds except those given for W and Y	1E+5	4E+5	2E-4	5E-7	1E-3	1E-2
		W, halides and nitrates	-	5E+5	2E-4	7E-7	-	-
		Y, oxides and hydroxides	-	5E+5	2E-4	6E-7	-	-
76	Osmium-181 ²	D, see ¹⁸⁰ Os	1E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, see ¹⁸⁰ Os	-	5E+4	2E-5	6E-8	-	-
		Y, see ¹⁸⁰ Os	-	4E+4	2E-5	6E-8	-	-
76	Osmium-182	D, see ¹⁸⁰ Os	2E+3	6E+3	2E-6	8E-9	3E-5	3E-4
		W, see ¹⁸⁰ Os	-	4E+3	2E-6	6E-9	-	-
		Y, see ¹⁸⁰ Os	-	4E+3	2E-6	6E-9	-	-
76	Osmium-185	D, see ¹⁸⁰ Os	2E+3	5E+2	2E-7	7E-10	3E-5	3E-4
		W, see ¹⁸⁰ Os	-	8E+2	3E-7	1E-9	-	-
		Y, see ¹⁸⁰ Os	-	8E+2	3E-7	1E-9	-	-
76	Osmium-189m	D, see ¹⁸⁰ Os	8E+4	2E+5	1E-4	3E-7	1E-3	1E-2
		W, see ¹⁸⁰ Os	-	2E+5	9E-5	3E-7	-	-
		Y, see ¹⁸⁰ Os	-	2E+5	7E-5	2E-7	-	-
76	Osmium-191m	D, see ¹⁸⁰ Os	1E+4	3E+4	1E-5	4E-8	2E-4	2E-3
		W, see ¹⁸⁰ Os	-	2E+4	8E-6	3E-8	-	-
		Y, see ¹⁸⁰ Os	-	2E+4	7E-6	2E-8	-	-
76	Osmium-191	D, see ¹⁸⁰ Os	2E+3	2E+3	9E-7	3E-9	-	-
			LLI wall (3E+3)	-	-	-	3E-5	3E-4
		W, see ¹⁸⁰ Os	-	2E+3	7E-7	2E-9	-	-
		Y, see ¹⁸⁰ Os	-	1E+3	6E-7	2E-9	-	-
76	Osmium-193	D, see ¹⁸⁰ Os	2E+3	5E+3	2E-6	6E-9	-	-
			LLI wall (2E+3)	-	-	-	2E-5	2E-4
		W, see ¹⁸⁰ Os	-	3E+3	1E-6	4E-9	-	-
		Y, see ¹⁸⁰ Os	-	3E+3	1E-6	4E-9	-	-
76	Osmium-194	D, see ¹⁸⁰ Os	4E+2	4E+1	2E-8	6E-11	-	-
			LLI wall (6E+2)	-	-	-	8E-6	8E-5
		W, see ¹⁸⁰ Os	-	6E+1	2E-8	8E-11	-	-
		Y, see ¹⁸⁰ Os	-	8E+0	3E-9	1E-11	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
77	Iridium-182 ²	D, all compounds except those given for W and Y	4E+4	1E+5	6E-5	2E-7	-	-
			St wall (4E+4)	-	-	-	6E-4	6E-3
77	Iridium-184	W, halides, nitrates, and metallic iridium	-	2E+5	6E-5	2E-7	-	-
		Y, oxides and hydroxides	-	1E+5	5E-5	2E-7	-	-
77	Iridium-182	D, see ¹⁸² Ir	8E+3	2E+4	1E-5	3E-8	1E-4	1E-3
		W, see ¹⁸² Ir	-	3E+4	1E-5	5E-8	-	-
		Y, see ¹⁸² Ir	-	3E+4	1E-5	4E-8	-	-
77	Iridium-185	D, see ¹⁸² Ir	5E+3	1E+4	5E-6	2E-8	7E-5	7E-4
		W, see ¹⁸² Ir	-	1E+4	5E-6	2E-8	-	-
		Y, see ¹⁸² Ir	-	1E+4	4E-6	1E-8	-	-
77	Iridium-186	D, see ¹⁸² Ir	2E+3	8E+3	3E-6	1E-8	3E-5	3E-4
		W, see ¹⁸² Ir	-	6E+3	3E-6	9E-9	-	-
		Y, see ¹⁸² Ir	-	6E+3	2E-6	8E-9	-	-
77	Iridium-187	D, see ¹⁸² Ir	1E+4	3E+4	1E-5	5E-8	1E-4	1E-3
		W, see ¹⁸² Ir	-	3E+4	1E-5	4E-8	-	-
		Y, see ¹⁸² Ir	-	3E+4	1E-5	4E-8	-	-
77	Iridium-188	D, see ¹⁸² Ir	2E+3	5E+3	2E-6	6E-9	3E-5	3E-4
		W, see ¹⁸² Ir	-	4E+3	1E-6	5E-9	-	-
		Y, see ¹⁸² Ir	-	3E+3	1E-6	5E-9	-	-
77	Iridium-189	D, see ¹⁸² Ir	5E+3	5E+3	2E-6	7E-9	-	-
		LLI wall (5E+3)	-	-	-	-	7E-5	7E-4
		W, see ¹⁸² Ir	-	4E+3	2E-6	5E-9	-	-
		Y, see ¹⁸² Ir	-	4E+3	1E-6	5E-9	-	-
77	Iridium-190m ²	D, see ¹⁸² Ir	2E+5	2E+5	8E-5	3E-7	2E-3	2E-2
		W, see ¹⁸² Ir	-	2E+5	9E-5	3E-7	-	-
		Y, see ¹⁸² Ir	-	2E+5	8E-5	3E-7	-	-
77	Iridium-190	D, see ¹⁸² Ir	1E+3	9E+2	4E-7	1E-9	1E-5	1E-4
		W, see ¹⁸² Ir	-	1E+3	4E-7	1E-9	-	-
		Y, see ¹⁸² Ir	-	9E+2	4E-7	1E-9	-	-
77	Iridium-192m	D, see ¹⁸² Ir	3E+3	9E+1	4E-8	1E-10	4E-5	4E-4
		W, see ¹⁸² Ir	-	2E+2	9E-8	3E-10	-	-
		Y, see ¹⁸² Ir	-	2E+1	6E-9	2E-11	-	-
77	Iridium-192	D, see ¹⁸² Ir	9E+2	3E+2	1E-7	4E-10	1E-5	1E-4
		W, see ¹⁸² Ir	-	4E+2	2E-7	6E-10	-	-
		Y, see ¹⁸² Ir	-	2E+2	9E-8	3E-10	-	-
77	Iridium-194m	D, see ¹⁸² Ir	6E+2	9E+1	4E-8	1E-10	9E-6	9E-5
		W, see ¹⁸² Ir	-	2E+2	7E-8	2E-10	-	-
		Y, see ¹⁸² Ir	-	1E+2	4E-8	1E-10	-	-
77	Iridium-194	D, see ¹⁸² Ir	1E+3	3E+3	1E-6	4E-9	1E-5	1E-4
		W, see ¹⁸² Ir	-	2E+3	9E-7	3E-9	-	-
		Y, see ¹⁸² Ir	-	2E+3	8E-7	3E-9	-	-
77	Iridium-195m	D, see ¹⁸² Ir	8E+3	2E+4	1E-5	3E-8	1E-4	1E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
77	Iridium-195	W, see ¹⁸² Ir	-	3E+4	1E-5	4E-8	-	-
		Y, see ¹⁸² Ir	-	2E+4	9E-6	3E-8	-	-
		D, see ¹⁸² Ir	1E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, see ¹⁸² Ir	-	5E+4	2E-5	7E-8	-	-
		Y, see ¹⁸² Ir	-	4E+4	2E-5	6E-8	-	-
78	Platinum-186	D, all compounds	1E+4	4E+4	2E-5	5E-8	2E-4	2E-3
78	Platinum-188	D, all compounds	2E+3	2E+3	7E-7	2E-9	2E-5	2E-4
78	Platinum-189	D, all compounds	1E+4	3E+4	1E-5	4E-8	1E-4	1E-3
78	Platinum-191	D, all compounds	4E+3	8E+3	4E-6	1E-8	5E-5	5E-4
78	Platinum-193m	D, all compounds	3E+3	6E+3	3E-6	8E-9	-	-
78	Platinum-193	D, all compounds	LLI wall (3E+4)	-	-	-	4E-5	4E-4
			4E+4	2E+4	1E-5	3E-8	-	-
78	Platinum-195m	D, all compounds	LLI wall (5E+4)	-	-	-	6E-4	6E-3
			2E+3	4E+3	2E-6	6E-9	-	-
78	Platinum-197m ²	D, all compounds	LLI wall (2E+3)	-	-	-	3E-5	3E-4
			2E+4	4E+4	2E-5	6E-8	2E-4	2E-3
78	Platinum-197	D, all compounds	3E+3	1E+4	4E-6	1E-8	4E-5	4E-4
78	Platinum-199 ²	D, all compounds	5E+4	1E+5	6E-5	2E-7	7E-4	7E-3
78	Platinum-200	D, all compounds	1E+3	3E+3	1E-6	5E-9	2E-5	2E-4
79	Gold-193	D, all compounds except those given for W and Y	9E+3	3E+4	1E-5	4E-8	1E-4	1E-3
		W, halides and nitrates	-	2E+4	9E-6	3E-8	-	-
		Y, oxides and hydroxides	-	2E+4	8E-6	3E-8	-	-
79	Gold-194	D, see ¹⁹³ Au	3E+3	8E+3	3E-6	1E-8	4E-5	4E-4
		W, see ¹⁹³ Au	-	5E+3	2E-6	8E-9	-	-
		Y, see ¹⁹³ Au	-	5E+3	2E-6	7E-9	-	-
79	Gold-195	D, see ¹⁹³ Au	5E+3	1E+4	5E-6	2E-8	7E-5	7E-4
		W, see ¹⁹³ Au	-	1E+3	6E-7	2E-9	-	-
		Y, see ¹⁹³ Au	-	4E+2	2E-7	6E-10	-	-
79	Gold-198m	D, see ¹⁹³ Au	1E+3	3E+3	1E-6	4E-9	1E-5	1E-4
		W, see ¹⁹³ Au	-	1E+3	5E-7	2E-9	-	-
		Y, see ¹⁹³ Au	-	1E+3	5E-7	2E-9	-	-
79	Gold-198	D, see ¹⁹³ Au	1E+3	4E+3	2E-6	5E-9	2E-5	2E-4
		W, see ¹⁹³ Au	-	2E+3	8E-7	3E-9	-	-
		Y, see ¹⁹³ Au	-	2E+3	7E-7	2E-9	-	-
79	Gold-199	D, see ¹⁹³ Au	3E+3	9E+3	4E-6	1E-8	-	-
		LLI wall (3E+3)	-	-	-	4E-5	4E-4	
		W, see ¹⁹³ Au	-	4E+3	2E-6	6E-9	-	-
		Y, see ¹⁹³ Au	-	4E+3	2E-6	5E-9	-	-
		D, see ¹⁹³ Au	1E+3	4E+3	1E-6	5E-9	2E-5	2E-4
79	Gold-200m	W, see ¹⁹³ Au	-	3E+3	1E-6	4E-9	-	-
		Y, see ¹⁹³ Au	-	2E+4	1E-6	3E-9	-	-
		D, see ¹⁹³ Au	3E+4	6E+4	3E-5	9E-8	4E-4	4E-3
		W, see ¹⁹³ Au	-	8E+4	3E-5	1E-7	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
79	Gold-201 ²	Y, see ¹⁹³ Au	-	7E+4	3E-5	1E-7	-	-
		D, see ¹⁹³ Au	7E+4	2E+5	9E-5	3E-7	-	-
		St wall (9E+4)	-	-	-	-	1E-3	1E-2
80	Mercury-193m	W, see ¹⁹³ Au	-	2E+5	1E-4	3E-7	-	-
		Y, see ¹⁹³ Au	-	2E+5	9E-5	3E-7	-	-
		Vapor	-	8E+3	4E-6	1E-8	-	-
		Organic D	4E+3	1E+4	5E-6	2E-8	6E-5	6E-4
		D, sulfates	3E+3	9E+3	4E-6	1E-8	4E-5	4E-4
80	Mercury-193	W, oxides, hydroxides, halides, nitrates, and sulfides	-	8E+3	3E-6	1E-8	-	-
		Vapor	-	3E+4	1E-5	4E-8	-	-
		Organic D	2E+4	6E+4	3E-5	9E-8	3E-4	3E-3
		D, see ^{193m} Hg	2E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, see ^{193m} Hg	-	4E+4	2E-5	6E-8	-	-
80	Mercury-194	Vapor	-	3E+1	1E-8	4E-11	-	-
		Organic D	2E+1	3E+1	1E-8	4E-11	2E-7	2E-6
		D, see ^{193m} Hg	8E+2	4E+1	2E-8	6E-11	1E-5	1E-4
		W, see ^{193m} Hg	-	1E+2	5E-8	2E-10	-	-
		Vapor	-	4E+3	2E-6	6E-9	-	-
80	Mercury-195m	Organic D	3E+3	6E+3	3E-6	8E-9	4E-5	4E-4
		D, see ^{193m} Hg	2E+3	5E+3	2E-6	7E-9	3E-5	3E-4
		W, see ^{193m} Hg	-	4E+3	2E-6	5E-9	-	-
		Vapor	-	3E+4	1E-5	4E-8	-	-
		Organic D	2E+4	5E+4	2E-5	6E-8	2E-4	2E-3
80	Mercury-195	D, see ^{193m} Hg	1E+4	4E+4	1E-5	5E-8	2E-4	2E-3
		W, see ^{193m} Hg	-	3E+4	1E-5	5E-8	-	-
		Vapor	-	5E+3	2E-6	7E-9	-	-
		Organic D	4E+3	9E+3	4E-6	1E-8	5E-5	5E-4
		D, see ^{193m} Hg	3E+3	7E+3	3E-6	1E-8	4E-5	4E-4
80	Mercury-197	W, see ^{193m} Hg	-	5E+3	2E-6	7E-9	-	-
		Vapor	-	8E+3	4E-6	1E-8	-	-
		Organic D	7E+3	1E+4	6E-6	2E-8	9E-5	9E-4
		D, see ^{193m} Hg	6E+3	1E+4	5E-6	2E-8	8E-5	8E-4
		W, see ^{193m} Hg	-	9E+3	4E-6	1E-8	-	-
80	Mercury-199m ²	Vapor	-	8E+4	3E-5	1E-7	-	-
		Organic D	6E+4	2E+5	7E-5	2E-7	-	-
		St wall (1E+5)	-	-	-	-	1E-3	1E-2
		D, see ^{193m} Hg	6E+4	1E+5	6E-5	2E-7	8E-4	8E-3
		W, see ^{193m} Hg	-	2E+5	7E-5	2E-7	-	-
80	Mercury-203	Vapor	-	8E+2	4E-7	1E-9	-	-
		Organic D	5E+2	8E+2	3E-7	1E-9	7E-6	7E-5
		D, see ^{193m} Hg	2E+3	1E+3	5E-7	2E-9	3E-5	3E-4
		W, see ^{193m} Hg	-	1E+3	5E-7	2E-9	-	-
		D, all compounds	5E+4	2E+5	6E-5	2E-7	-	-
81	Thallium-194m ²		St wall (7E+4)	-	-	-	1E-3	1E-2

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
81	Thallium-194 ²	D, all compounds	3E+5	6E+5	2E-4	8E-7	-	-
			St wall (3E+5)	-	-	-	4E-3	4E-2
81	Thallium-195 ²	D, all compounds	6E+4	1E+5	5E-5	2E-7	9E-4	9E-3
81	Thallium-197	D, all compounds	7E+4	1E+5	5E-5	2E-7	1E-3	1E-2
81	Thallium-198m ²	D, all compounds	3E+4	5E+4	2E-5	8E-8	4E-4	4E-3
81	Thallium-198	D, all compounds	2E+4	3E+4	1E-5	5E-8	3E-4	3E-3
81	Thallium-199	D, all compounds	6E+4	8E+4	4E-5	1E-7	9E-4	9E-3
81	Thallium-200	D, all compounds	8E+3	1E+4	5E-6	2E-8	1E-4	1E-3
81	Thallium-201	D, all compounds	2E+4	2E+4	9E-6	3E-8	2E-4	2E-3
81	Thallium-202	D, all compounds	4E+3	5E+3	2E-6	7E-9	5E-5	5E-4
81	Thallium-204	D, all compounds	2E+3	2E+3	9E-7	3E-9	2E-5	2E-4
82	Lead-195m ²	D, all compounds	6E+4	2E+5	8E-5	3E-7	8E-4	8E-3
82	Lead-198	D, all compounds	3E+4	6E+4	3E-5	9E-8	4E-4	4E-3
82	Lead-199 ²	D, all compounds	2E+4	7E+4	3E-5	1E-7	3E-4	3E-3
82	Lead-200	D, all compounds	3E+3	6E+3	3E-6	9E-9	4E-5	4E-4
82	Lead-201	D, all compounds	7E+3	2E+4	8E-6	3E-8	1E-4	1E-3
82	Lead-202m	D, all compounds	9E+3	3E+4	1E-5	4E-8	1E-4	1E-3
82	Lead-202	D, all compounds	1E+2	5E+1	2E-8	7E-11	2E-6	2E-5
82	Lead-203	D, all compounds	5E+3	9E+3	4E-6	1E-8	7E-5	7E-4
82	Lead-205	D, all compounds	4E+3	1E+3	6E-7	2E-9	5E-5	5E-4
82	Lead-209	D, all compounds	2E+4	6E+4	2E-5	8E-8	3E-4	3E-3
82	Lead-210	D, all compounds	6E-1	2E-1	1E-10	-	-	-
			Bone surf (1E+0)	Bone surf (4E-1)	-	6E-13	1E-8	1E-7
82	Lead-211 ²	D, all compounds	1E+4	6E+2	3E-7	9E-10	2E-4	2E-3
82	Lead-212	D, all compounds	8E+1	3E+1	1E-8	5E-11	-	-
			Bone surf (1E+2)	-	-	-	2E-6	2E-5
82	Lead-214 ²	D, all compounds	9E+3	8E+2	3E-7	1E-9	1E-4	1E-3
83	Bismuth-200 ²	D, nitrates	3E+4	8E+4	4E-5	1E-7	4E-4	4E-3
		W, all other compounds	-	1E+5	4E-5	1E-7	-	-
83	Bismuth-201 ²	D, see ²⁰⁰ Bi	1E+4	3E+4	1E-5	4E-8	2E-4	2E-3
		W, see ²⁰⁰ Bi	-	4E+4	2E-5	5E-8	-	-
83	Bismuth-202 ²	D, see ²⁰⁰ Bi	1E+4	4E+4	2E-5	6E-8	2E-4	2E-3
		W, see ²⁰⁰ Bi	-	8E+4	3E-5	1E-7	-	-
83	Bismuth-203	D, see ²⁰⁰ Bi	2E+3	7E+3	3E-6	9E-9	3E-5	3E-4
		W, see ²⁰⁰ Bi	-	6E+3	3E-6	9E-9	-	-
83	Bismuth-205	D, see ²⁰⁰ Bi	1E+3	3E+3	1E-6	3E-9	2E-5	2E-4
		W, see ²⁰⁰ Bi	-	1E+3	5E-7	2E-9	-	-
83	Bismuth-206	D, see ²⁰⁰ Bi	6E+2	1E+3	6E-7	2E-9	9E-6	9E-5
		W, see ²⁰⁰ Bi	-	9E+2	4E-7	1E-9	-	-
83	Bismuth-207	D, see ²⁰⁰ Bi	1E+3	2E+3	7E-7	2E-9	1E-5	1E-4
		W, see ²⁰⁰ Bi	-	4E+2	1E-7	5E-10	-	-
83	Bismuth-210m	D, see ²⁰⁰ Bi	4E+1	5E+0	2E-9	-	-	-
			Kidneys (6E+1)	Kidneys (6E+0)	-	9E-12	8E-7	8E-6
		W, see ²⁰⁰ Bi	-	7E-1	3E-10	9E-13	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration
			Oral Ingestion	Inhalation		Air $\mu\text{Ci/ml}$	Water $\mu\text{Ci/ml}$	$\mu\text{Ci/ml}$
				ALI μCi	ALI μCi			
83	Bismuth-210	D, see ^{200}Bi	8E+2	2E+2	1E-7	-	1E-5	1E-4
			-	Kidneys (4E+2)	-	5E-10	-	-
		W, see ^{200}Bi	-	3E+1	1E-8	4E-11	-	-
83	Bismuth-212 ²	D, see ^{200}Bi	5E+3	2E+2	1E-7	3E-10	7E-5	7E-4
		W, see ^{200}Bi	-	3E+2	1E-7	4E-10	-	-
83	Bismuth-213 ²	D, see ^{200}Bi	7E+3	3E+2	1E-7	4E-10	1E-4	1E-3
		W, see ^{200}Bi	-	4E+2	1E-7	5E-10	-	-
83	Bismuth-214 ²	D, see ^{200}Bi	2E+4	8E+2	3E-7	1E-9	-	-
			St wall (2E+4)	-	-	-	3E-4	3E-3
		W, see ^{200}Bi	-	9E-2	4E-7	1E-9	-	-
84	Polonium-203 ²	D, all compounds except those given for W	3E+4	6E+4	3E-5	9E-8	3E-4	3E-3
		W, oxides, hydroxides, and nitrates	-	9E+4	4E-5	1E-7	-	-
84	Polonium-205 ²	D, see ^{203}Po	2E+4	4E+4	2E-5	5E-8	3E-4	3E-3
		W, see ^{203}Po	-	7E+4	3E-5	1E-7	-	-
84	Polonium-207	D, see ^{203}Po	8E+3	3E+4	1E-5	3E-8	1E-4	1E-3
		W, see ^{203}Po	-	3E+4	1E-5	4E-8	-	-
84	Polonium-210	D, see ^{203}Po	3E+0	6E-1	3E-10	9E-13	4E-8	4E-7
		W, see ^{203}Po	-	6E-1	3E-10	9E-13	-	-
85	Astatine-207 ²	D, halides	6E+3	3E+3	1E-6	4E-9	8E-5	8E-4
		W	-	2E+3	9E-7	3E-9	-	-
85	Astatine-211	D, halides	1E+2	8E+1	3E-8	1E-10	2E-6	2E-5
		W	-	5E+1	2E-8	8E-11	-	-
86	Radon-220	With daughters removed	-	2E+4	7E-6	2E-8	-	-
		With daughters present	-	2E+1	9E-9	3E-11	-	-
				(or 12 working level months)		(or 1.0 working level)		
86	Radon-222	With daughters removed	-	1E+4	4E-6	1E-8	-	-
		With daughters present	-	1E+2	3E-8	1E-10	-	-
				(or 4 working level months)		(or 0.33 working level)		
87	Francium-222 ²	D, all compounds	2E+3	5E+2	2E-7	6E-10	3E-5	3E-4
87	Francium-223 ²	D, all compounds	6E+2	8E+2	3E-7	1E-9	8E-6	8E-5
88	Radium-223	W, all compounds	5E+0	7E-1	3E-10	9E-13	-	-
			Bone surf (9E+0)	-	-	-	1E-7	1E-6
88	Radium-224	W, all compounds	8E+0	2E+0	7E-10	2E-12	-	-
			Bone surf (2E+1)	-	-	-	2E-7	2E-6
88	Radium-225	W, all compounds	8E+0	7E-1	3E-10	9E-13	-	-
			Bone surf (2E+1)	-	-	-	2E-7	2E-6
88	Radium-226	W, all compounds	2E+0	6E-1	3E-10	9E-13	-	-
			Bone surf (5E+0)	-	-	-	6E-8	6E-7

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
88	Radium-227 ²	W, all compounds	2E+4	1E+4	6E-6	-	-	-
			Bone surf (2E+4)	Bone surf (2E+4)	-	3E-8	3E-4	3E-3
88	Radium-228	W, all compounds	2E+0	1E+0	5E-10	2E-12	-	-
			Bone surf (4E+0)	-	-	-	6E-8	6E-7
89	Actinium-224	D, all compounds except those given for W and Y	2E+3	3E+1	1E-8	-	-	-
			LLI wall (2E+3)	Bone surf (4E+1)	-	5E-11	3E-5	3E-4
		W, halides and nitrates	-	5E+1	2E-8	7E-11	-	-
		Y, oxides and hydroxides	-	5E+1	2E-8	6E-11	-	-
89	Actinium-225	D, see ²²⁴ Ac	5E+1	3E-1	1E-10	-	-	-
			LLI wall (5E+1)	Bone surf (5E-1)	-	7E-13	7E-7	7E-6
		W, see ²²⁴ Ac	-	6E-1	3E-10	9E-13	-	-
		Y, see ²²⁴ Ac	-	6E-1	3E-10	9E-13	-	-
89	Actinium-226	D, see ²²⁴ Ac	1E+2	3E+0	1E-9	-	-	-
			LLI wall (1E+2)	Bone surf (4E+0)	-	5E-12	2E-6	2E-5
		W, see ²²⁴ Ac	-	5E+0	2E-9	7E-12	-	-
		Y, see ²²⁴ Ac	-	5E+0	2E-9	6E-12	-	-
89	Actinium-227	D, see ²²⁴ Ac	2E-1	4E-4	2E-13	-	-	-
			Bone surf (4E-1)	Bone surf (8E-4)	-	1E-15	5E-9	5E-8
		W, see ²²⁴ Ac	-	2E-3	7E-13	-	-	-
			-	Bone surf (3E-3)	-	4E-15	-	-
		Y, see ²²⁴ Ac	-	4E-3	2E-12	6E-15	-	-
89	Actinium-228	D, see ²²⁴ Ac	2E+3	9E+0	4E-9	-	3E-5	3E-4
			-	Bone surf (2E+1)	-	2E-11	-	-
		W, see ²²⁴ Ac	-	4E+1	2E-8	-	-	-
			-	Bone surf (6E+1)	-	8E-11	-	-
		Y, see ²²⁴ Ac	-	4E+1	2E-8	6E-11	-	-
90	Thorium-226 ²	W, all compounds except those given for Y	5E+3	2E+2	6E-8	2E-10	-	-
			St wall (5E+3)	-	-	-	7E-5	7E-4
		Y, oxides and hydroxides	-	1E+2	6E-8	2E-10	-	-
90	Thorium-227	W, see ²²⁶ Th	1E+2	3E-1	1E-10	5E-13	2E-6	2E-5
		Y, see ²²⁶ Th	-	3E-1	1E-10	5E-13	-	-
90	Thorium-228	W, see ²²⁶ Th	6E+0	1E-2	4E-12	-	-	-
			Bone surf (1E+1)	Bone surf (2E-2)	-	3E-14	2E-7	2E-6
		Y, see ²²⁶ Th	-	2E-2	7E-12	2E-14	-	-
90	Thorium-229	W, see ²²⁶ Th	6E-1	9E-4	4E-13	-	-	-
			Bone surf (1E+0)	Bone surf (2E-3)	-	3E-15	2E-8	2E-7
		Y, see ²²⁶ Th	-	2E-3	1E-12	-	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
90	Thorium-230	W, see ²²⁶ Th	-	Bone surf (3E-3)	-	4E-15	-	-
			4E+0	6E-3	3E-12	-	-	-
		Y, see ²²⁶ Th	Bone surf (9E+0)	Bone surf (2E-2)	-	2E-14	1E-7	1E-6
90	Thorium-231	W, see ²²⁶ Th	-	2E-2	6E-12	-	-	-
			4E+3	6E+3	3E-6	9E-9	5E-5	5E-4
		Y, see ²²⁶ Th	-	6E+3	3E-6	9E-9	-	-
90	Thorium-232	W, see ²²⁶ Th	7E-1	1E-3	5E-13	-	-	-
			Bone surf (2E+0)	Bone surf (3E-3)	-	4E-15	3E-8	3E-7
		Y, see ²²⁶ Th	-	3E-3	1E-12	-	-	-
90	Thorium-234	W, see ²²⁶ Th	-	Bone surf (4E-3)	-	6E-15	-	-
			3E+2	2E+2	8E-8	3E-10	-	-
		Y, see ²²⁶ Th	LLI wall (4E+2)	-	-	-	5E-6	5E-5
91	Protactinium-227 ²	W, all compounds except those given for Y	-	2E+2	6E-8	2E-10	-	-
			4E+3	1E+2	5E-8	2E-10	5E-5	5E-4
		Y, oxides and hydroxides	-	1E+2	4E-8	1E-10	-	-
91	Protactinium-228	W, see ²²⁷ Pa	1E+3	1E+1	5E-9	-	2E-5	2E-4
			-	Bone surf (2E+1)	-	3E-11	-	-
		Y, see ²²⁷ Pa	-	1E+1	5E-9	2E-11	-	-
91	Protactinium-230	W, see ²²⁷ Pa	6E+2	5E+0	2E-9	7E-12	-	-
			Bone surf (9E+2)	-	-	-	1E-5	1E-4
		Y, see ²²⁷ Pa	-	4E+0	1E-9	5E-12	-	-
91	Protactinium-231	W, see ²²⁷ Pa	2E-1	2E-3	6E-13	-	-	-
			Bone surf (5E-1)	Bone surf (4E-3)	-	6E-15	6E-9	6E-8
		Y, see ²²⁷ Pa	-	4E-3	2E-12	-	-	-
91	Protactinium-232	W, see ²²⁷ Pa	-	Bone surf (6E-3)	-	8E-15	-	-
			1E+3	2E+1	9E-9	-	2E-5	2E-4
		Y, see ²²⁷ Pa	-	Bone surf (6E+1)	-	8E-11	-	-
91	Protactinium-233	W, see ²²⁷ Pa	-	6E+1	2E-8	-	-	-
			1E+3	7E+2	3E-7	1E-9	-	-
		Y, see ²²⁷ Pa	LLI wall (2E+3)	-	-	-	2E-5	2E-4
91	Protactinium-234	W, see ²²⁷ Pa	-	6E+2	2E-7	8E-10	-	-
			2E+3	8E+3	3E-6	1E-8	3E-5	3E-4
		Y, see ²²⁷ Pa	-	7E+3	3E-6	9E-9	-	-
92	Uranium-230	D, UF ₆ , UO ₂ F ₂ , UO ₂ (NO ₃) ₂	4E+0	4E-1	2E-10	-	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers	
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml	
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml		
				ALI μCi	ALI μCi				DAC μCi/ml
92	Uranium-231		Bone surf (6E+0)	Bone surf (6E-1)	-	8E-13	8E-8	8E-7	
		W, UO ₃ , UF ₄ , UCl ₄	-	4E-1	1E-10	5E-13	-	-	
		Y, UO ₂ , U ₃ O ₈	-	3E-1	1E-10	4E-13	-	-	
		D, see ²³⁰ U	5E+3	8E+3	3E-6	1E-8	-	-	
92	Uranium-232		LLI wall (4E+3)	-	-	-	6E-5	6E-4	
		W, see ²³⁰ U	-	6E+3	2E-6	8E-9	-	-	
		Y, see ²³⁰ U	-	5E+3	2E-6	6E-9	-	-	
		D, see ²³⁰ U	2E+0	2E-1	9E-11	-	-	-	
92	Uranium-233		Bone surf (4E+0)	Bone surf (4E-1)	-	6E-13	6E-8	6E-7	
		W, see ²³⁰ U	-	4E-1	2E-10	5E-13	-	-	
		Y, see ²³⁰ U	-	8E-3	3E-12	1E-14	-	-	
		D, see ²³⁰ U	1E+1	1E+0	5E-10	-	-	-	
92	Uranium-234 ³		Bone surf (2E+1)	Bone surf (2E+0)	-	3E-12	3E-7	3E-6	
		W, see ²³⁰ U	-	7E-1	3E-10	1E-12	-	-	
		Y, see ²³⁰ U	-	4E-2	2E-11	5E-14	-	-	
		D, see ²³⁰ U	1E+1	1E+0	5E-10	-	-	-	
92	Uranium-235 ³		Bone surf (2E+1)	Bone surf (2E+0)	-	3E-12	3E-7	3E-6	
		W, see ²³⁰ U	-	7E-1	3E-10	1E-12	-	-	
		Y, see ²³⁰ U	-	4E-2	2E-11	5E-14	-	-	
		D, see ²³⁰ U	1E+1	1E+0	6E-10	-	-	-	
92	Uranium-236		Bone surf (2E+1)	Bone surf (2E+0)	-	3E-12	3E-7	3E-6	
		W, see ²³⁰ U	-	8E-1	3E-10	1E-12	-	-	
		Y, see ²³⁰ U	-	4E-2	2E-11	6E-14	-	-	
		D, see ²³⁰ U	1E+1	1E+0	5E-10	-	-	-	
92	Uranium-237		Bone surf (2E+1)	Bone surf (2E+0)	-	3E-12	3E-7	3E-6	
		W, see ²³⁰ U	-	8E-1	3E-10	1E-12	-	-	
		Y, see ²³⁰ U	-	4E-2	2E-11	6E-14	-	-	
		D, see ²³⁰ U	2E+3	3E+3	1E-6	4E-9	-	-	
92	Uranium-238 ³		LLI wall (2E+3)	-	-	-	3E-5	3E-4	
		W, see ²³⁰ U	-	2E+3	7E-7	2E-9	-	-	
		Y, see ²³⁰ U	-	2E+3	6E-7	2E-9	-	-	
		D, see ²³⁰ U	1E+1	1E+0	6E-10	-	-	-	
92	Uranium-239 ²		Bone surf (2E+1)	Bone surf (2E+0)	-	3E-12	3E-7	3E-6	
		W, see ²³⁰ U	-	8E-1	3E-10	1E-12	-	-	
		Y, see ²³⁰ U	-	4E-2	2E-11	6E-14	-	-	
		D, see ²³⁰ U	7E+4	2E+5	8E-5	3E-7	9E-4	9E-3	
92	Uranium-240		W, see ²³⁰ U	-	2E+5	7E-5	2E-7	-	-
		Y, see ²³⁰ U	-	2E+5	6E-5	2E-7	-	-	
		D, see ²³⁰ U	1E+3	4E+3	2E-6	5E-9	2E-5	2E-4	

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
92	Uranium-natural ³	W, see ²³⁰ U	-	3E+3	1E-6	4E-9	-	-
		Y, see ²³⁰ U	-	2E+3	1E-6	3E-9	-	-
		D, see ²³⁰ U	1E+1	1E+0	5E-10	-	-	-
93	Neptunium-232 ²		Bone surf (2E+1)	Bone surf (2E+0)	-	3E-12	3E-7	3E-6
		W, see ²³⁰ U	-	8E-1	3E-10	9E-13	-	-
		Y, see ²³⁰ U	-	5E-2	2E-11	9E-14	-	-
93	Neptunium-233 ²	W, all compounds	1E+5	2E+3	7E-7	-	2E-3	2E-2
			-	Bone surf (5E+2)	-	6E-9	-	-
93	Neptunium-234	W, all compounds	8E+5	3E+6	1E-3	4E-6	1E-2	1E-1
93	Neptunium-234	W, all compounds	2E+3	3E+3	1E-6	4E-9	3E-5	3E-4
93	Neptunium-235	W, all compounds	2E+4	8E+2	3E-7	-	-	-
93	Neptunium-236 (1.15E+5 y)	W, all compounds	LLI wall (2E+4)	Bone surf (1E+3)	-	2E-9	3E-4	3E-3
			3E+0	2E-2	9E-12	-	-	-
93	Neptunium-236 (22.5 h)	W, all compounds	Bone surf (6E+0)	Bone surf (5E-2)	-	8E-14	9E-8	9E-7
			3E+3	3E+1	1E-8	-	-	-
93	Neptunium-237	W, all compounds	Bone surf (4E+3)	Bone surf (7E+1)	-	1E-10	5E-5	5E-4
			5E-1	4E-3	2E-12	-	-	-
93	Neptunium-238	W, all compounds	Bone surf (1E+0)	Bone surf (1E-2)	-	1E-14	2E-8	2E-7
			1E+3	6E+1	3E-8	-	2E-5	2E-4
93	Neptunium-239	W, all compounds	-	Bone surf (2E+2)	-	2E-10	-	-
			2E+3	2E+3	9E-7	3E-9	-	-
93	Neptunium-240 ²	W, all compounds	LLI wall (2E+3)	-	-	-	2E-5	2E-4
			2E+4	8E+4	3E-5	1E-7	3E-4	3E-3
94	Plutonium-234	W, all compounds except PuO ₂	8E+3	2E+2	9E-8	3E-10	1E-4	1E-3
		Y, PuO ₂	-	2E+2	8E-8	3E-10	-	-
94	Plutonium-235 ²	W, see ²³⁴ Pu	9E+5	3E+6	1E-3	4E-6	1E-2	1E-1
		Y, see ²³⁴ Pu	-	3E+6	1E-3	3E-6	-	-
94	Plutonium-236	W, see ²³⁴ Pu	2E+0	2E-2	8E-12	-	-	-
			Bone surf (4E+0)	Bone surf (4E-2)	-	5E-14	6E-8	6E-7
		Y, see ²³⁴ Pu	-	4E-2	2E-11	6E-14	-	-
94	Plutonium-237	W, see ²³⁴ Pu	1E+4	3E+3	1E-6	5E-9	2E-4	2E-3
		Y, see ²³⁴ Pu	-	3E+3	1E-6	4E-9	-	-
94	Plutonium-238	W, see ²³⁴ Pu	9E-1	7E-3	3E-12	-	-	-
			Bone surf (2E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
		Y, see ²³⁴ Pu	-	2E-2	8E-12	2E-14	-	-
94	Plutonium-239	W, see ²³⁴ Pu	8E-1	6E-3	3E-12	-	-	-
			Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
		Y, see ²³⁴ Pu	-	2E-2	7E-12	-	-	-
		-	Bone surf (2E-2)	-	2E-14	-	-	

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
94	Plutonium-240	W, see ²³⁴ Pu	8E-1	6E-3	3E-12	-	-	-
		Y, see ²³⁴ Pu	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
94	Plutonium-241	W, see ²³⁴ Pu	4E+1	3E-1	1E-10	-	-	-
		Y, see ²³⁴ Pu	Bone surf (7E+1)	Bone surf (6E-1)	-	8E-13	1E-6	1E-5
94	Plutonium-242	W, see ²³⁴ Pu	8E-1	7E-3	3E-12	-	-	-
		Y, see ²³⁴ Pu	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
94	Plutonium-243	W, see ²³⁴ Pu	2E+4	4E+4	2E-5	5E-8	2E-4	2E-3
		Y, see ²³⁴ Pu	-	4E+4	2E-5	5E-8	-	-
94	Plutonium-244	W, see ²³⁴ Pu	8E-1	7E-3	3E-12	-	-	-
		Y, see ²³⁴ Pu	Bone surf (2E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
94	Plutonium-245	W, see ²³⁴ Pu	2E+3	5E+3	2E-6	6E-9	3E-5	3E-4
		Y, see ²³⁴ Pu	-	4E+3	2E-6	6E-9	-	-
94	Plutonium-246	W, see ²³⁴ Pu	4E+2	3E+2	1E-7	4E-10	-	-
		Y, see ²³⁴ Pu	LLI wall (4E+2)	-	-	-	6E-6	6E-5
95	Americium-237 ²	W, all compounds	8E+4	3E+5	1E-4	4E-7	1E-3	1E-2
95	Americium-238 ²	W, all compounds	4E+4	3E+3	1E-6	-	5E-4	5E-3
		Y, see ²³⁴ Pu	-	Bone surf (6E+3)	-	9E-9	-	-
95	Americium-239	W, all compounds	5E+3	1E+4	5E-6	2E-8	7E-5	7E-4
95	Americium-240	W, all compounds	2E+3	3E+3	1E-6	4E-9	3E-5	3E-4
95	Americium-241	W, all compounds	8E-1	6E-3	3E-12	-	-	-
		Y, see ²³⁴ Pu	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
95	Americium-242m	W, all compounds	8E-1	6E-3	3E-12	-	-	-
		Y, see ²³⁴ Pu	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
95	Americium-242	W, all compounds	4E+3	8E+1	4E-8	-	5E-5	5E-4
		Y, see ²³⁴ Pu	-	Bone surf (9E+1)	-	1E-10	-	-
95	Americium-243	W, all compounds	8E-1	6E-3	3E-12	-	-	-
		Y, see ²³⁴ Pu	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
95	Americium-244m ²	W, all compounds	6E+4	4E+3	2E-6	-	-	-

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
95	Americium-244	W, all compounds	St wall (8E+4)	Bone surf (7E+3)	-	1E-8	1E-3	1E-2
			3E+3	2E+2	8E-8	-	4E-5	4E-4
95	Americium-245	W, all compounds	-	Bone surf (3E+2)	-	4E-10	-	-
			3E+4	8E+4	3E-5	1E-7	4E-4	4E-3
95	Americium-246m ²	W, all compounds	5E+4	2E+5	8E-5	3E-7	-	-
95	Americium-246 ²	W, all compounds	St wall (6E+4)	-	-	-	8E-4	8E-3
			3E+4	1E+5	4E-5	1E-7	4E-4	4E-3
96	Curium-238	W, all compounds	2E+4	1E+3	5E-7	2E-9	2E-4	2E-3
96	Curium-240	W, all compounds	6E+1	6E-1	2E-10	-	-	-
96	Curium-241	W, all compounds	Bone surf (8E+1)	Bone surf (6E-1)	-	9E-13	1E-6	1E-5
			1E+3	3E+1	1E-8	-	2E-5	2E-4
96	Curium-242	W, all compounds	-	Bone surf (4E+1)	-	5E-11	-	-
			3E+1	3E-1	1E-10	-	-	-
96	Curium-243	W, all compounds	Bone surf (5E+1)	Bone surf (3E-1)	-	4E-13	7E-7	7E-6
			1E+0	9E-3	4E-12	-	-	-
96	Curium-244	W, all compounds	Bone surf (2E+0)	Bone surf (2E-2)	-	2E-14	3E-8	3E-7
			1E+0	1E-2	5E-12	-	-	-
96	Curium-245	W, all compounds	Bone surf (3E+0)	Bone surf (2E-2)	-	3E-14	3E-8	3E-7
			7E-1	6E-3	3E-12	-	-	-
96	Curium-246	W, all compounds	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
			7E-1	6E-3	3E-12	-	-	-
96	Curium-247	W, all compounds	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
			8E-1	6E-3	3E-12	-	-	-
96	Curium-248	W, all compounds	Bone surf (1E+0)	Bone surf (1E-2)	-	2E-14	2E-8	2E-7
			2E-1	2E-3	7E-13	-	-	-
96	Curium-249 ²	W, all compounds	Bone surf (4E-1)	Bone surf (3E-3)	-	4E-15	5E-9	5E-8
			5E+4	2E+4	7E-6	-	7E-4	7E-3
96	Curium-250	W, all compounds	-	Bone surf (3E+4)	-	4E-8	-	-
			4E-2	3E-4	1E-13	-	-	-
97	Berkelium-245	W, all compounds	Bone surf (6E-2)	Bone surf (5E-4)	-	8E-16	9E-10	9E-9
			2E+3	1E+3	5E-7	2E-9	3E-5	3E-4
97	Berkelium-246	W, all compounds	3E+3	3E+3	1E-6	4E-9	4E-5	4E-4
97	Berkelium-247	W, all compounds	5E-1	4E-3	2E-12	-	-	-
97	Berkelium-249	W, all compounds	Bone surf (1E+0)	Bone surf (9E-3)	-	1E-14	2E-8	2E-7
			2E+2	2E+0	7E-10	-	-	-
97	Berkelium-250	W, all compounds	Bone surf (5E+2)	Bone surf (4E+0)	-	5E-12	6E-6	6E-5
			9E+3	3E+2	1E-7	-	1E-4	1E-3

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
98	Californium-244 ²	W, all compounds except those given for Y	-	Bone surf (7E+2)	-	1E-9	-	-
			3E+4	6E+2	2E-7	8E-10	-	-
98	Californium-246	Y, oxides and hydroxides	-	6E+2	2E-7	8E-10	-	-
		W, see ²⁴⁴ Cf	4E+2	9E+0	4E-9	1E-11	5E-6	5E-5
98	Californium-248	Y, see ²⁴⁴ Cf	-	9E+0	4E-9	1E-11	-	-
		W, see ²⁴⁴ Cf	8E+0	6E-2	3E-11	-	-	-
98	Californium-249	Y, see ²⁴⁴ Cf	-	1E-1	4E-11	1E-13	-	-
		W, see ²⁴⁴ Cf	5E-1	4E-3	2E-12	-	-	-
98	Californium-250	Y, see ²⁴⁴ Cf	-	1E-2	4E-12	-	-	-
		W, see ²⁴⁴ Cf	1E+0	9E-3	4E-12	-	-	-
98	Californium-251	Y, see ²⁴⁴ Cf	-	3E-2	1E-11	4E-14	-	-
		W, see ²⁴⁴ Cf	5E-1	4E-3	2E-12	-	-	-
98	Californium-252	Y, see ²⁴⁴ Cf	-	3E-2	1E-11	5E-14	-	-
		W, see ²⁴⁴ Cf	2E+0	2E-2	8E-12	-	-	-
98	Californium-253	Y, see ²⁴⁴ Cf	-	3E-2	1E-11	5E-14	-	-
		W, see ²⁴⁴ Cf	2E+2	2E+0	8E-10	3E-12	-	-
98	Californium-254	Y, see ²⁴⁴ Cf	-	2E+0	7E-10	2E-12	-	-
		W, see ²⁴⁴ Cf	2E+0	2E-2	9E-12	3E-14	3E-8	3E-7
99	Einsteinium-250	W, all compounds	4E+4	5E+2	2E-7	-	6E-4	6E-3
99	Einsteinium-251	W, all compounds	-	Bone surf (1E+3)	-	2E-9	-	-
			7E+3	9E+2	4E-7	-	1E-4	1E-3
99	Einsteinium-253	W, all compounds	2E+2	1E+0	6E-10	2E-12	2E-6	2E-5
99	Einsteinium-254m	W, all compounds	3E+2	1E+1	4E-9	1E-11	-	-
			LLI wall (3E+2)	-	-	-	4E-6	4E-5
99	Einsteinium-254	W, all compounds	8E+0	7E-2	3E-11	-	-	-
			Bone surf (2E+1)	Bone surf (1E-1)	-	2E-13	2E-7	2E-6

Atomic No.	Radionuclide	Class	Table I Occupational Values			Table II Effluent Concentration		Table III Releases to Sewers
			Col. 1	Col. 2	Col. 3	Col. 1	Col. 2	Monthly Average Concentration μCi/ml
			Oral Ingestion	Inhalation		Air μCi/ml	Water μCi/ml	
				ALI μCi	ALI μCi			
100	Fermium-252	W, all compounds	5E+2	1E+1	5E-9	2E-11	6E-6	6E-5
100	Fermium-253	W, all compounds	1E+3	1E+1	4E-9	1E-11	1E-5	1E-4
100	Fermium-254	W, all compounds	3E+3	9E+1	4E-8	1E-10	4E-5	4E-4
100	Fermium-255	W, all compounds	5E+2	2E+1	9E-9	3E-11	7E-6	7E-5
100	Fermium-257	W, all compounds	2E+1	2E-1	7E-11	-	-	-
			Bone surf (4E+1)	Bone surf (2E-1)	-	3E-13	5E-7	5E-6
101	Mendelevium-257	W, all compounds	7E+3	8E+1	4E-8	-	1E-4	1E-3
			-	Bone surf (9E+1)	-	1E-10	-	-
101	Mendelevium-258	W, all compounds	3E+1	2E-1	1E-10	-	-	-
			Bone surf (5E+1)	Bone surf (3E-1)	-	5E-13	6E-7	6E-6
-	Any single radionuclide not listed above with decay mode other than alpha emission or spontaneous fission and with radioactive half-life less than ((2)) <u>two</u> hours	Submersion ¹	-	2E+2	1E-7	1E-9	-	-
-	Any single radionuclide not listed above with decay mode other than alpha emission or spontaneous fission and with radioactive half-life greater than ((2)) <u>two</u> hours	-	2E-1	1E-10	1E-12	1E-8	1E-7
-	Any single radionuclide not listed above that decays by alpha emission or spontaneous fission, or any mixture for which either the identity or the concentration of any radionuclide in the mixture is not known	-	4E-4	2E-13	1E-15	2E-9	2E-8

FOOTNOTES:

¹"Submersion" means that values given are for submersion in a hemispherical semi-infinite cloud of airborne material.

²These radionuclides have radiological half-lives of less than ((2)) two hours. The total effective dose equivalent received during operations with these radionuclides might include a significant contribution from external exposure. The DAC values for all radionuclides, other than those designated Class "Submersion," are based upon the committed effective dose equivalent due to the intake of the radionuclide into the body and do NOT include potentially significant contributions to dose equivalent from external exposures. The licensee may substitute 1E-7 µCi/ml for the listed DAC to account for the submersion dose prospectively, but should use individual monitoring devices or other radiation measuring instruments that measure external exposure to demonstrate compliance with the limits. (See WAC 246-221-015(5).)

³For soluble mixtures of U-238, U-234, and U-235 in air, chemical toxicity may be the limiting factor (see WAC 246-221-010(5)). If the percent by weight (enrichment) of U-235 is not greater than ((5)) five, the concentration value for a 40-hour workweek is 0.2 milligrams uranium per cubic meter of air average. For any enrichment, the product of the average concentration and time of exposure during a 40-hour workweek shall not exceed 8E-3 (SA) µCi-hr/ml, where SA is the specific activity of the uranium inhaled. The specific activity for natural uranium is 6.77E-7 curies per gram U. The specific activity for other mixtures of U-238, U-235, and U-234, if not known, shall be:

$$SA = 3.6E-7 \text{ curies/gram U, U-depleted}$$

$$SA = [0.4 + 0.38 (\text{enrichment}) + 0.0034 (\text{enrichment})^2] E-6, \text{ enrichment} \geq 0.72 \text{ where enrichment is the percentage by weight of U-235, expressed as percent.}$$

NOTE:

1. If the identity of each radionuclide in a mixture is known but the concentration of one or more of the radionuclides in the mixture is not known, the DAC for the mixture shall be the most restrictive DAC of any radionuclide in the mixture.
2. If the identity of each radionuclide in the mixture is not known, but it is known that certain radionuclides specified in this appendix are not present in the mixture, the inhalation ALI, DAC, and effluent and sewage concentrations for the mixture are the lowest values specified in this appendix for any radionuclide that is not known to be absent from the mixture; or

If it is known that Ac-227-D and Cm-250-W are not present	-	7E-4	3E-13	-	-	-
If, in addition, it is known that Ac-227-W,Y, Th-229-W,Y, Th-230-W, Th-232-W,Y, Pa-231-W,Y, Np-237-W, Pu-239-W, Pu-240-W, Pu-242-W, Am-241-W, Am-242m-W, Am-243-W, Cm-245-W, Cm-246-W, Cm-247-W, Cm-248-W, Bk-247-W, Cf-249-W, and Cf-251-W are not present	-	7E-3	3E-12	-	-	-
If, in addition, it is known that Sm-146-W, Sm-147-W, Gd-148-D,W, Gd-152-D,W, Th-228-W,Y, Th-230-Y, U-232-Y, U-233-Y, U-234-Y, U-235-Y, U-236-Y, U-238-Y, Np-236-W, Pu-236-W,Y, Pu-238-W,Y, Pu-239-Y, Pu-240-Y, Pu-242-Y, Pu-244-W,Y, Cm-243-W, Cm-244-W, Cf-248-W, Cf-249-Y, Cf-250-W,Y, Cf-251-Y, Cf-252-W,Y, and Cf-254-W,Y are not present	-	7E-2	3E-11	-	-	-
If, in addition, it is known that Pb-210-D, Bi-210m-W, Po-210-D,W, Ra-223-W, Ra-225-W, Ra-226-W, Ac-225-D,W,Y, Th-227-W,Y, U-230-D,W,Y, U-232-D,W, Pu-241-W, Cm-240-W, Cm-242-W, Cf-248-Y, Es-254-W, Fm-257-W, and Md-258-W are not present	-	7E-1	3E-10	-	-	-
If, in addition, it is known that Si-32-Y, Ti-44-Y, Fe-60-D, Sr-90-Y, Zr-93-D, Cd-113m-D, Cd-113-D, In-115-D,W, La-138-D, Lu-176-W, Hf-178m-D,W, Hf-182-D,W, Bi-210m-D, Ra-224-W, Ra-228-W, Ac-226-D,W,Y, Pa-230-W,Y, U-233-D,W, U-234-D,W, U-235-D,W, U-236-D,W, U-238-D,W, Pu-241-Y, Bk-249-W, Cf-253-W,Y, and Es-253-W are not present	-	7E+0	3E-9	-	-	-
If it is known that Ac-227-D,W,Y, Th-229-W,Y, Th-232-W,Y, Pa-231-W,Y, Cm-248-W, and Cm-250-W are not present	-	-	-	1E-14	-	-
If, in addition, it is known that Sm-146-W, Gd-148-D,W, Gd-152-D, Th-228-W,Y, Th-230-W,Y, U-232-Y, U-233-Y, U-234-Y, U-235-Y, U-236-Y, U-238-Y, U-Nat-Y, Np-236-W, Np-237-W, Pu-236-W,Y, Pu-238-W,Y, Pu-239-W,Y, Pu-240-W,Y, Pu-242-W,Y, Pu-244-W,Y, Am-241-W, Am-242m-W, Am-243-W, Cm-243-W, Cm-244-W, Cm-245-W, Cm-246-W, Cm-247-W, Bk-247-W, Cf-249-W,Y, Cf-250-W,Y, Cf-251-W,Y, Cf-252-W,Y, and Cf-254-W,Y are not present	-	-	-	1E-13	-	-
If, in addition, it is known that Sm-147-W, Gd-152-W, Pb-210-D, Bi-210m-W, Po-210-D,W, Ra-223-W, Ra-225-W, Ra-226-W, Ac-225-D,W,Y, Th-227-W,Y, U-230-D,W,Y, U-232-D,W, U-Nat-W, Pu-241-W, Cm-240-W, Cm-242-W, Cf-248-W,Y, Es-254-W, Fm-257-W, and Md-258-W are not present	-	-	-	-	1E-12	-
If, in addition, it is known that Fe-60, Sr-90, Cd-113m, Cd-113, In-115, I-129, Cs-134, Sm-145, Sm-147, Gd-148, Gd-152, Hg-194 (organic), Bi-210m, Ra-223, Ra-224, Ra-225, Ac-225, Th-228, Th-230, U-233, U-234, U-235, U-236, U-238, U-Nat, Cm-242, Cf-248, Es-254, Fm-257, and Md-258 are not present	-	-	-	-	1E-6	1E-5

3. If a mixture of radionuclides consists of uranium and its daughters in ore dust (10 µm AMAD particle distribution assumed) prior to chemical separation of the uranium from the ore, the following values may be used for the DAC of the mixture: 6E-11 µCi of gross alpha activity from uranium-238, uranium-234, thorium-230, and radium-226 per milliliter of air; 3E-11 µCi of natural uranium per milliliter of air; or 45 micrograms of natural uranium per cubic meter of air.

4. If the identity and concentration of each radionuclide in a mixture are known, the limiting values should be derived as follows: Determine, for each radionuclide in the mixture, the ratio between the concentration present in the mixture and the concentration otherwise established in this section for the specific radionuclide when not in a mixture. The sum of such ratios for all of the radionuclides in the mixture may not exceed "1" (i.e., "unity").

Example: If radionuclides "A," "B," and "C" are present in concentrations CA, CB, and CC, and if the applicable DACs are DAC_A, DAC_B, and DAC_C, respectively, then the concentrations shall be limited so that the following relationship exists:

$$\frac{C_A}{DAC_A} + \frac{C_B}{DAC_B} + \frac{C_C}{DAC_C} \leq 1$$