# DANCESAFE FENTANYL TEST STRIPS

# **DANCESAFE**FENTANYL TEST STRIPS

## DANCESAFE FENTANYL TEST STRIPS

### WHY SHOULD I TEST MY DRUGS FOR FENTANYL?

- Fentanyl is a powerful opioid. It's often found in pressed pills or powder.
- If you know fentanyl might be in your drugs, it can help you make decisions that lower your risk of overdose.



## WHAT SHOULD I KNOW ABOUT FENTANYL TEST STRIPS?

- Fentanyl isn't always mixed evenly into drugs. A single negative test does not mean your entire supply is fentanyl-free.
- Test strips should be used in combination with other harm reduction strategies:
  - Start with a smaller dose or test shot.
  - $\,^\circ\,$  Take turns using with a friend.
  - Call a friend or the Never Use Alone Hotline (877-696-1996) if you use alone.
  - · Carry naloxone and know how to use it.



 If you know fentanyl might be in your drugs, it can help you make decisions that lower your risk of overdose.

in pressed pills

or powder.



## WHAT SHOULD I KNOW ABOUT FENTANYL TEST STRIPS?

- Fentanyl isn't always mixed evenly into drugs. A single negative test does not mean your entire supply is fentanyl-free.
- Test strips should be used in combination with other harm reduction strategies:
  - Start with a smaller dose or test shot.
  - $\,^\circ\,$  Take turns using with a friend.
  - Call a friend or the Never Use Alone Hotline (877-696-1996) if you use alone.
  - · Carry naloxone and know how to use it.



DOH 150-230, September 2023. To request this document in another format, call 1-800-525-0127. Dead for hard of hearing customers, please call 711 (Washington Relay) or email doh.information@doh.wa.gov.

### WHY SHOULD I TEST MY DRUGS FOR FENTANYL?

- Fentanyl is a powerful opioid. It's often found in pressed pills or powder.
- If you know fentanyl might be in your drugs, it can help you make decisions that lower your risk of overdose.



## WHAT SHOULD I KNOW ABOUT FENTANYL TEST STRIPS?

- Fentanyl isn't always mixed evenly into drugs. A single negative test does not mean your entire supply is fentanyl-free.
- Test strips should be used in combination with other harm reduction strategies:
  - Start with a smaller dose or test shot.
  - Take turns using with a friend.
  - Call a friend or the Never Use Alone Hotline (877-696-1996) if you use alone.
  - Carry naloxone and know how to use it.



DOH 150-230, September 2023. To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email doh.information@doh.wa.gov.


# HOW TO USE DANCESAFE FENTANYL TEST STRIPS

# HOW TO USE DANCESAFE FENTANYL TEST STRIPS

# HOW TO USE DANCESAFE FENTANYL TEST STRIPS

#### 1. PREPARE YOUR SAMPLE

- To test residue left in a cooker:
  - Add 2mL of clean water (2 full syringes) to your cooker. Mix well.
- To test pills, crystal, and powders:
  - Crush up your drugs and mix them up.
  - Add 10mg of drugs (about the size of a match head or grain of rice) in a cooker.



 Add 1mL of water (about 20 drops or 1 full syringe) for every 10mg of drugs. Mix well.

#### 2. DIP THE STRIP FOR 15 SECONDS.

Hold the yellow end of the test strip.
 Don't dip it past the dotted area.

#### 3. WAIT 3 MINUTES FOR RESULTS.

 Place the strip face up on a flat surface while you wait.

### **READING RESULTS**

One red line means the test is positive. You may have fentanyl in your drugs.

1 RED LINE = POSITIVE for FENTANYL

N FEN FEN FEN FEN FEN FEN FEN

Two red lines means the test is negative. Look closely for a second line – it can be very faint!


2 RED LINES = NEGATIVE for FENTANYL:

N FEN FEN FEN FEN FEN FEN FEN

If there is only a red line on the left or no red lines appear, the test is INVALID. Redo the test with a new strip.

#### 1. PREPARE YOUR SAMPLE

- To test residue left in a cooker:
  - Add 2mL of clean water (2 full syringes) to your cooker. Mix well.
- To test pills, crystal, and powders:
  - Crush up your drugs and mix them up.
  - Add 10mg of drugs (about the size of a match head or grain of rice) in a cooker.



Add 1mL of water

 (about 20 drops or 1 full syringe) for
 every 10mg of drugs. Mix well.

#### 2. DIP THE STRIP FOR 15 SECONDS.

Hold the yellow end of the test strip.
 Don't dip it past the dotted area.

#### 3. WAIT 3 MINUTES FOR RESULTS.

 Place the strip face up on a flat surface while you wait.

### **READING RESULTS**

One red line means the test is positive. You may have fentanyl in your drugs.

1 RED LINE = POSITIVE for FENTANYL

N FEN FEN FEN FEN FEN FEN FEN

Two red lines means the test is negative. Look closely for a second line – it can be very faint!


2 RED LINES = NEGATIVE for FENTANYL:

N FEN FEN FEN FEN FEN FEN FEN FEN

If there is only a red line on the left or no red lines appear,
the test is INVALID. Redo the test with a new strip.

#### 1. PREPARE YOUR SAMPLE

- To test residue left in a cooker:
  - Add 2mL of clean water (2 full syringes) to your cooker. Mix well.
- · To test pills, crystal, and powders:
  - Crush up your drugs and mix them up.
  - Add 10mg of drugs (about the size of a match head or grain of rice) in a cooker.



 Add 1mL of water (about 20 drops or 1 full syringe) for every 10mg of drugs. Mix well.

#### 2. DIP THE STRIP FOR 15 SECONDS.

Hold the yellow end of the test strip.
 Don't dip it past the dotted area.

#### 3. WAIT 3 MINUTES FOR RESULTS.

• Place the strip face up on a flat surface while you wait.

### **READING RESULTS**

One red line means the test is positive. You may have fentanyl in your drugs.

1 RED LINE = POSITIVE for FENTANYL

N FEN FEN FEN FEN FEN FEN FEN FEN

Two red lines means the test is negative. Look closely for a second line – it can be very faint!

2 RED LINES = NEGATIVE for FENTANYL:

N FEN FEN FEN FEN FEN FEN FEN

If there is only a red line on the left or no red lines appear,
the test is INVALID. Redo the test with a new strip.