

Carbapenem-Resistant *Pseudomonas aeruginosa* (CRPA)

A drug-resistant germ that spreads in healthcare facilities



What is *Pseudomonas aeruginosa*?

The germ, *Pseudomonas aeruginosa*, is a type of bacteria that can cause life-threatening infections in patients with serious medical problems in hospitals and nursing homes. When these germs become resistant to strong antibiotics such as carbapenems, they are harder to treat and even more dangerous. Carbapenem-resistant *Pseudomonas aeruginosa* is called CRPA for short.

What is colonization?

Patients can become colonized with CRPA. Colonization means people carry CRPA in or on their bodies but don't have any illness symptoms. Colonization with CRPA can last for many months and even years. People who are colonized can still spread these germs to others.

Why is CRPA a problem?

CRPA can spread quickly from person to person in healthcare facilities on healthcare workers' hands, healthcare equipment, and surfaces, if they are not cleaned properly. CRPA infections can be difficult to treat and even cause death.

Who is more likely to get CRPA?

CRPA mainly affects patients who have long-term, serious medical problems. People who have spent a lot of time in healthcare facilities are at the highest risk of becoming colonized or developing an infection with CRPA. Other things that increase the risk of CRPA infection include:

- Being on a mechanical ventilator (breathing machine) for a long time.
- Having tubes going into the body, such as urinary catheters, central venous catheters, and feeding or breathing tubes.

How does CRPA spread to patients?

Patients with infection or colonization with CRPA can have the germs on their skin and in body fluids that can contaminate their living space, clothing, and medical equipment. CRPA can contaminate healthcare workers' hands, clothing, and other items in the healthcare environment – like doorknobs, bed rails, stethoscopes, blood pressure cuffs, and thermometers—and can quickly spread to other patients.

How can we prevent the spread of CRPA?

To prevent the spread of CRPA in healthcare facilities, healthcare workers should:

- Know how germs spread and make sure they do not carry them to other patients on their clothing, hands, or equipment.
- Wash or sanitize hands before and after caring for patients or touching items in the patient's room.

- Use the right personal protective equipment when caring for patients who can spread infections – often this includes gowns and gloves.
- Use hospital grade disinfectants for disinfecting areas and items that might have CRPA germs on them and make sure that environmental services staff know how to correctly use these products.
- Clean and disinfect shared equipment before and after use on a patient.

To prevent the spread of CRPA in the home:

- Always clean your hands with soap and water:
 - Before preparing or eating food.
 - After using the bathroom.
 - Before and after touching any wounds or medical devices that you may have.
- Do not share personal care items like towels and washcloths with other people.
- Cover any wounds with a bandage whenever possible.
- No special cleaning is required in your home:
 - Your clothing may be washed in the usual manner, along with the rest of the household laundry.
 - All eating utensils and dishes can be washed the way you normally do.
- You do not need to take special precautions around your pets.

Who Should be Tested for CRPA?

Public Health recommends testing patients for CRPA if they may have been exposed to another person with CRPA infection or colonization. The testing is to see if they have the CRPA germ on their body.

Should healthcare workers and family members be tested to see if they have CRPA?

Public Health does not recommend testing healthy family members or healthcare workers who care for patients with CRPA infection or colonization. In general, healthy people are not at risk from CRPA and do not need to be tested. However, healthcare facilities may decide to screen household contacts of patients with CRE if they become ill and are admitted to a healthcare facility.



DOH 420-502 December 2024

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 ([Washington Relay](#)) or email doh.information@doh.wa.gov.