SEATTLE CHILDREN'S HOSPITAL ANTIBIOGRAM 2020

NUMBERS ARE PERCENT SUSCEPTIBLE

GRAM POSITIVE ORGANISMS (Some antimicrobial agents known to lack clinical efficacy are not included.)	Number tested (U = Urine)	Ampicillin	Amox-Clavulanate	Cefazolin	Cefepime	Ceftriaxone	Ciprofloxacin	Clindamycin NU	Doxycycline	Erythromycin NU	Gentamicin	Gent Synergy	Levofloxacin	Linezolid	Meropenem	Nitrofurantoin U	Oxacillin	Penicillin (Parenteral Oral)	Rifampin	Trim-Sulfa	Vancomycin
Staphylococcus aureus (mSsa) not CF	813 (U27)		97	100		89	79	72		42	98			99			100	28	98	98	100 ^a
Staphylococcus aureus (mRsa) not CF ^b	186 (U9)		0	0		0	21	64		16	97			100			0	0	99	91	100
Staphylococcus aureus (CF) mSsa and mRsa	255		86	87		72	74	71		43	96			100			88	20	96	99	100
Staphylococcus epidermidis	166 (U51)		39	39		39		33			79			98		100	39	12	97	48	100
Streptococcus pyogenes	61				98	100		53		73								100			100
Streptococcus pneumoniae ^c	58 (CSF1) ^d					94		91	35	69			98	100	82 ^e			96 66	100	73	100
Viridans streptococci ^f	45				84	83		86		42			78					48			100
Enterococcus faecalis	201 (U161)	100										91	99	92		97					100
Enterococcus faecium ⁹	25 (U14)	64										88	50	87		28					76

One isolate of VISA (Vancomycin Intermediate Staph aureus) was recovered in 2020.

mRsa: Methicillin resistant Staphylococcus aureus. Non-CF mRsa rate for 2020 = 19%

mSsa: Methicillin susceptible Staphylococcus aureus

Seattle Children's Hospital

4800 Sand Point Way N.E. MS: OC.8.720

Seattle, Washington 98105

Sean Murphy, M.D./Ph.D. Microbiology Laboratory Director

Microbiology Laboratory: (206) 987-2585

U: Only urinary isolates included Only non-urinary isolates included NU:

2020 Seattle Children's Hospital All Rights Reserved



Methicillin resistance was tested also by latex agglutination of the PBP and/or molecular assay for mecA gene.

^{6%} of ceftriaxone MICs and 4% / 25% of penicillin (parenteral/oral) MICs were in the 'INTERMEDIATE' range.

CSF breakpoint for CRO Susceptible is <=0.5. CSF breakpoint for Pen IV Susceptible is <=0.064.

Number of meropenem tests was insufficient for antibiogram threshold. In 2019, there was 100% susceptibility to meropenem.

^{8%} of cefepime MICs, 7% of ceftriaxone MICs, and 44% of penicillin MICs were in the 'INTERMEDIATE' range.

g Number of E faecium tested is insufficient for antibiogram threshold