

Specimen Collection and Submission Instructions Carbapenem-resistant Enterobacterales (CRE) Carbapenemase Testing Page 1 of 2

Specimen Type	Collection Time	Collection Frequency	Collection Procedures	Transport Media	Shipping & Handling (S&H)*
Escherichia coli, Klebsiella oxytoca, Klebsiella pneumoniae, and Enterobacter spp that are resistant to imipenem, meropenem, doripenem, or ertapenem by standard susceptibility testing methods. Minimum inhibitory concentrations: o doripenem ≥4 µg/ml, or o imipenem ≥4 µg/ml, or o meropenem ≥4 µg/ml, or o ertapenem ≥2 µg/ml	NA	NA	NA	Choc, HIA, BHI slant or plate. Note: Plate may be used <i>only</i> if transported by courier.	Transport device: Choc, HIA, BHI slant or plate. Note: Plate may be used only if transported by courier. Rejection Criteria:



Specimen Collection and Submission Instructions Carbapenem-resistant Enterobacterales (CRE) Carbapenemase Testing Page 2 of 2

General Rejection Criteria (for additional details, see S&H)

• Failure to follow specific S&H requirements.

NOTE:

Specimens will not be processed until ALL of the following information are provided on requisition form:

- Patient name, patient's county of residence, and second identifier.
- Two matching identifiers on both the tube and the laboratory requisition form.
- Specimen type, date of collection, and test requested.
- Submitter name, address, and telephone/FAX numbers.
- Reason for testing at WAPHL.

Collection kit availability

Not available.

Comments

None.