

Table 1: Summary of Guidance for PrEP Use

	Men Who Have Sex with Men	Heterosexual Women and Men	Persons Who Inject Drugs
Detecting substantial risk of acquiring HIV infection	HIV-positive sexual partner Recent bacterial STI [†] High number of sex partners History of inconsistent or no condom use Commercial sex work	HIV-positive sexual partner Recent bacterial STI [‡] High number of sex partners History of inconsistent or no condom use Commercial sex work In high HIV prevalence area or network	HIV-positive injecting partner Sharing injection equipment
Clinically eligible	Documented negative HIV test result before prescribing PrEP No signs/symptoms of acute HIV infection Normal renal function; no contraindicated medications Documented hepatitis B virus infection and vaccination status		
Prescription	Daily, continuing, oral doses of TDF/FTC (Truvada), ≤90-day supply		
Other services	Follow-up visits at least every 3 months to provide the following: HIV test, medication adherence counseling, behavioral risk reduction support, side effect assessment, STI symptom assessment At 3 months and every 6 months thereafter, assess renal function Every 3-6 months, test for bacterial STIs		
	Do oral/rectal STI testing	For women, assess pregnancy intent Pregnancy test every 3 months	Access to clean needles/syringes and drug treatment services

STI: sexually transmitted infection

[†] Gonorrhea, chlamydia, syphilis for MSM including those who inject drugs

[‡] Gonorrhea, syphilis for heterosexual women and men including those who inject drugs