Adapted from WA DOH Form 347-102

Fax Page 1 To: Whitman County Health Department (509) 397-6239 (Confidential FAX line)



CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Report STIs within 3 work days (WAC 246-101-101/301)

| PATIENT INFORMATION | | | | | | | | | | | | | | |
|---|--|---|---|---|---|------------------------------|---------------------------|--|---|--------------------------|------------|------------|-------------|---|
| LAST NAME | | FIRST NAME | | | | MIDDLE NAME | | DAT | DATE OF BIRTH | | | | | |
| | | | | | | | M | | мо | D DAY | | | YR | |
| ADDRESS (Unhoused or unstably housed | | | in the past 3 months) | | СІТҮ | | | | STATE | | ZIP COD | E | | |
| | | | | | | | | | | | | | | |
| TELEPHONE EMAIL | | | | ENGLISH SPEAKING? Yes No | | | lo *instructions pg. 3 | ^{on} C | DIAGNOS | IS DA | TE | | | |
| () | | | Prefer | | Prefer | red Language | | | • | MO DAY YR | | YR | | |
| SEX ASSIGNED GENDER IDENTITY | | ПТҮ | ETHNICITY | | | RACE CATEGORY (check all th | | that | apply)*: | | *In | structions | s on page 3 | |
| AT BIRTH 🗌 Male | | Transgender MTF | | Hispanic or | | | | | | | | Asiar | | |
| Male Female | | Transgender FTM | | Non-Hispanic | | American Indian / Alaska N | | a Nat | ☐ Other Native ☐ Unknown | | | | | |
| Female Female Female Genderquee | | | | Unknown Refused | | □ Native Hawaiian / Other Pa | | | | | | | | |
| ☐ Intersex Genderqueer ☐ Refused ☐ Refused | | | | EXTENDED RACE COD | | | | CE CODE(S) | *: F | R | | R | R | R |
| CURRENTLY | REASON FOR E | XAM (check one) | GENDER OF SEX PARTNERS (check all that apply): HIV ST | | | HIV STATU | | | | | | | | |
| PREGNANT? | | | | _ 0 | | | | • | • | | ON P | | | |
| □ Yes □ Symptomati | | | | _ 0 | | | | | / diagnosis at this visit* ☐ Yes e HIV test at this visit ☐ No | | | | | |
| □ No □ Unk. □ NA | □ Routine Exar | | | onbinary / 🔅 🗌 Other enderqueer 🔅 🗌 Unknow | | | | test (unknown status) | | | | | | |
| DIAGNOSIS - D | | | 1 | | | | | l | | | | | I | |
| GONORRHEA (I | | | | | | | | | PHIL | | , | | | |
| DIAGNOSIS (check one) | | SITES (all that apply): | | TREATMENT (check all prescribed): | | | | STAGE (check one): | | | | | | |
| | Asymptomatic Symptomatic, Uncomplicated | | | | Ceftriaxone: 250 mg 500 mg 1 g Cefixime: 400 mg 800 mg | | | | Primary (Chancre, etc.) Secondary (Rash, etc.) | | | | | |
| Pelvic Inflammatory Disease | | · — · · · · · · · · · · · · · · · · · · | | \square Azithromycin: \square 1 g \square 2 g | | | | | Latent (< | | | | | |
| Ophthalmia Disseminated | | Rectum | | Doxycycline: 100 mg BID x 7 d | | | | Javs — | | Unknown Duration or Late | | | | |
| Other Compli | | | | Gentamicin: 240 mg | | | | Congenital | | | | | | |
| | | | | | Gemifloxacin: 320 mg | | | | MANIFESTATIONS (check all that apply): | | | | | |
| Date Tested: | | Other: | | Date Prescribed: | | | | | □ Neurologic □ Otic □ Ocular □ Tertiary | | | | | |
| CHLAMYDIA (la | b confirmed) | 1 | | | | | | TR | EATN | /IENT (ch | eck d | one): | | |
| DIAGNOSIS (che | | SITES (all that apply): | | TREATMENT (check all prescribed): | | | | Bicillin L - A: 2.4 MU IM x 1 | | | | | | |
| Asymptomati | | Cervix | | Azithromycin: 1 g | | | | 2.4 MU IM x 3 | | | | | | |
| Symptomatic Pelvic Inflam | , Uncomplicated | | | Doxycycline: 100 mg BID x 7 days | | | Do | хусус | | | ng BID x 1 | | | |
| 🗌 Ophthalmia | | | | Levofloxacin: 500 mg daily x 7 days Other: | | | _ | ☐ 100 mg BID x 28 days | | | | | | |
| Other Compl | ications: | Pharynx | | | | | | Benzathine 50,000 units/kg IM x 1 PCN-G: 50,000 units/kg IM x 3 | | | | | | |
| | | U Vagina | | | | | - | | | | | | 5 | |
| Date Tested: | | Ocular | | Date F | Date Prescribed: | | | | Aqueous 18-24 MU/day IV Crystalline for 10-14 days | | | | | |
| HERPES SIMPLE | X | | | | OTHER | DISEASES | | Pe | nicilli | | | o _ : aayo | | |
| DIAGNOSIS LABORATORY CONFIRMATION Chancroid Other: | | | | | | | | | | | | | | |
| Genital (initial infection only) | | | | | 🔲 Granuloma In | | | | ite Pr | escribed: | | | | |
| | | Lymphogranuloma Venereum | | | | | | | | | | | | |
| | PARTNER TREATMENT PLAN (check one or more options) Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2 | | | | | | | | | | | | | |
| for additional information). | | | | | | | | | | | | | | |
| · | | of partners treat | | - | | | | | | | | | | |
| Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis | | | | | | | | | | | | | | |
| REPORTING CLINIC INFORMATION | | | | | | | | | | | | | | |
| DATE FACILITY NAME DIAGNOSING CLINICIAN | | | | | | | | | | | | | | |
| ADDRESS CITY STATE ZIP | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | |
| PERSON COMPL | ETING FORM | | | TELEPHONE | | | EMAIL | MAIL | | | | | | |
| Thank you for reporting an STI. All i | | | | (|) | ••• • | 1 1.1 .1 | | <i>c. ,</i> | | | | | |
| | Thank | you for reportin | g an STI. | All infor | mation | will be man | aged with the s | strictest con | nder | mality. | | | | |

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile.

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Whitman County Health Department may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Whitman County Health Department recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Whitman County Health Department: (509) 397-6280.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs)[†]

Alternatives for uncomplicated infections of the cervix, urethra, or rectum:[‡]

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose OR

Cefixime 800 mg orally as a single dose[†]

⁺ If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.

[‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days **OR**

Ofloxacin 300 mg PO BID for 7 days **OR**

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>https://www.cdc.gov/std/treatment/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

DOH 347-102, updated 12/19/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

List of Preferred Languages:

Instructions: Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

| (L01) Amharic | (L18) Karen | (L35) Sign languages |
|-----------------------------|-------------------------|-----------------------------------|
| (L02) Arabic | (L19) Khmer/Cambodian | (L36) Somali |
| (LO3) Balochi/Baluchi | (L20) Kinyarwanda | (L37) Spanish/Castilian |
| (LO4) Burmese | (L21) Korean | (L38) Swahili/Kiswahili |
| (L05) Cantonese | (L22) Kosraean | (L39) Tagalog |
| (L06) Chinese (unspecified) | (L23) Lao | (L40) Tamil |
| (L07) Chamorro | (L24) Mandarin | (L41) Telugu |
| (L08) Chuukese | (L25) Marshallese | (L42) Thai |
| (L09) Dari | (L26) Mixteco | (L43) Tigrinya |
| (L10) Farsi/Persian | (L27) Nepali | (L44) Ukrainian |
| (L11) Fijian | (L28) Oromo | (L45) Urdu |
| (L12) Filipino/Pilipino | (L29) Panjabi/Punjabi | (L46) Vietnamese |
| (L13) French | (L30) Pashto | (L77) Other language |
| (L14) German | (L31) Portuguese | (L88) Patient declined to respond |
| (L15) Hindi | (L32) Romanian/Rumanian | (L99) Unknown |
| (L16) Hmong | (L33) Russian | |
| (L17) Japanese | (L34) Samoan | |
| | | |

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

Extended Race Codes:

| (R01) Afghan | (R22) Fijian | (R43) Mexican/Mexican American | (R64) Tongan |
|---------------------------------|-----------------------------|--------------------------------|------------------|
| (R02) Afro-Caribbean | (R23) Filipino | (R44) Middle Eastern | (R65) Ugandan |
| (R03) Alaska Native | (R24) First Nations | (R45) Mien | (R66) Ukrainian |
| (R04) American Indian | (R25) Guamanian or Chamorro | (R46) Moroccan | (R67) Vietnamese |
| (R05) Arab | (R26) Hmong/Mong | (R47) Native Hawaiian | (R68) White |
| (R06) Asian | (R27) Indigenous-Latina/o/x | (R48) Nepalese | (R69) Yemeni |
| (R07) Asian Indian | (R28) Indonesian | (R49) North African | (R77) Other race |
| (R08) Bamar/Burman/Burmese | (R29) Iranian | (R50) Oromo | (R99) Unknown |
| (R09) Bangladeshi | (R30) Iraqi | (R51) Pacific Islander | |
| (R10) Bhutanese | (R31) Japanese | (R52) Pakistani | |
| (R11) Black or African American | (R32) Jordanian | (R53) Puerto Rican | |
| (R12) Central American | (R33) Karen | (R54) Romanian/Rumanian | |
| (R13) Cham | (R34) Kenyan | (R55) Russian | |
| (R14) Chicana/o or Chicanx | (R35) Khmer/Cambodian | (R56) Samoan | |
| (R15) Chinese | (R36) Korean | (R57) Saudi Arabian | |
| (R16) Congolese | (R37) Kuwaiti | (R58) Somali | |
| (R17) Cuban | (R38) Lao | (R59) South African | |
| (R18) Dominican | (R39) Lebanese | (R60) South American | |
| (R19) Egyptian | (R40) Malaysian | (R61) Syrian | |
| (R20) Eritrean | (R41) Marshallese | (R62) Taiwanese | |
| (R21) Ethiopian | (R42) Mestizo | (R63) Thai | |
| | | | |