

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019

Helpful Hints

Hospital Reports:

Astria - Sunnyside Community Hospital
Astria – Toppenish Community Hospital
Cascade Valley Hospital
Confluence Health – Central Washington Hospital
Coulee Medical Center
EvergreenHealth Kirkland
Forks Community Hospital
Grays Harbor Community Hospital
Island Hospital
Jefferson Healthcare
Kadlec Regional Medical Center
Kittitas Valley Healthcare
Lake Chelan Community Hospital
Legacy Salmon Creek Medical Center
Madigan Army Medical Center
Mason General Hospital
Mid-Valley Hospital
MultiCare – Auburn Medical Center
MultiCare – Capital Medical Center
MultiCare – Covington Medical Center
MultiCare – Deaconess Hospital
MultiCare – Good Samaritan Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019

MultiCare – Tacoma General Hospital
MultiCare – Valley Hospital
Naval Hospital – Bremerton
Naval Hospital – Oak Harbor
Newport Hospital
North Valley Hospital
Olympic Medical Center
Othello Community Hospital
Overlake Medical Center
PeaceHealth – Southwest Family Birth Center
PeaceHealth – St John Medical Center
PeaceHealth - St Joseph Hospital – Bellingham
Prosser Memorial Hospital
Providence – Centralia Hospital
Providence – Everett Regional Medical Center
Providence – Holy Family Hospital
Providence – Mount Carmel Hospital
Providence – Sacred Heart Medical Center
Providence – St Mary Medical Center
Providence – St Peter Hospital
Pullman Regional Hospital
Samaritan Healthcare
Skagit Valley Hospital
Swedish Medical Center – Ballard
Swedish Medical Center – Edmonds

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019

Swedish Medical Center – First Hill

Swedish Medical Center – Issaquah

Three Rivers Hospital

Trios Health Hospital

University of Washington Medical Center

University of Washington Medical Center – Northwest

University of Washington Medical Center – Valley Medical Center

Virginia Mason Franciscan Health – St Anne Hospital

Virginia Mason Franciscan Health – St Elizabeth Hospital

Virginia Mason Franciscan Health – St Francis Hospital

Virginia Mason Franciscan Health – St Joseph Medical Center - Tacoma

Virginia Mason Franciscan Health – St Michael Medical Center

WhidbeyHealth Medical Center

Whitman Hospital and Medical Center

Yakima Valley Memorial

Helpful Hints



Newborn Hearing Training Curriculum



<http://infanthearing.org/nhstc>

- Interactive, web-based training
- Separate modules for self-paced learning
- Tracks pre-tests and post-tests for personnel record of competencies
- Free and staff can earn a Certificate of Completion and/or Continuing Education Units (CEUs)

- Use a reliable tracking system to ensure babies get needed screens, rescreens, and diagnostic referrals.
- Provide annual competency training for all screeners to reinforce adequate techniques and knowledge. Check out the NCHAM resource to the left.
- Perform a rescreen before discharge.
- Do a maximum of two screen attempts per test type per ear.
- Find out the name of the clinic where the infant will be seen for well child checks.
- Get at least 2 contact numbers or email addresses for families needing follow-up.
- Give families all results both verbally and in writing (the Department of Health (DOH) has templates) in their preferred language.
- Set up appointments for outpatient rescreens or audiology before discharge.
- Rescreen both ears even if only one ear referred.
- Use a script found in the DOH protocol book for standardized explanations to parents.
- Say "did not pass" rather than "failed."
- Understand and assess for risk factors.

Reporting to DOH

- **For the initial screen, use the original pink card from the 1st blood spot.**
- **If the original pink card is unavailable, use a blue card.**
- **Mark corrections clearly – strike through the error and circle the correction.**
- **In the Outpatient Provider field on the card, write in the name of clinic or the clinic number of where the infant will be seen.**
- **A designated person should review all cards for completion before sending to DOH.**
- **Mail hearing screening cards to DOH weekly.**
- **Use 'Notes' to indicate a missed screen, NICU status, transferred, expired, or other**

Astria – Sunnyside Community Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Astria – Toppenish Community Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Astria – Toppenish Community Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Cascade Valley Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Cascade Valley Hospital **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Confluence Health – Central Washington Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Coulee Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Coulee Medical Center **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



EvergreenHealth Kirkland

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Forks Community Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Forks Community Hospital **14** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Grays Harbor Community Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Island Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Island Hospital **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Jefferson Healthcare

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Jefferson Healthcare **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Kadlec Regional Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Kittitas Valley Healthcare

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Kittitas Valley Healthcare **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Lake Chelan Community Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Lake Chelan Community Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Legacy Salmon Creek Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Madigan Army Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Mason General Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Mason General Hospital **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Mid-Valley Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss

Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare – Auburn Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare – Capital Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.


Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare – Covington Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare – Deaconess Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare - Good Samaritan Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare – Tacoma General Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



MultiCare – Valley Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Naval Hospital – Bremerton

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Naval Hospital – Oak Harbor

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Newport Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Newport Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



North Valley Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

North Valley Hospital **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Olympic Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Olympic Medical Center **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Othello Community Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Overlake Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



PeaceHealth – Southwest Family Birth Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



PeaceHealth – St. John Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

PeaceHealth St. John Med. Center **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



PeaceHealth – St. Joseph Hospital - Bellingham

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Prosser Memorial Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – Centralia Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – Everett Regional Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – Holy Family Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – Mount Carmel Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen




Referred Initial Hearing Screen



Recommend Refer Rate to Not Exceed 5%

Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Providence Mount Carmel Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – Sacred Heart Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.


Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.


Diagnosed with Hearing Loss




Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – St. Mary Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Providence – St. Peter Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Providence St. Peter Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Pullman Regional Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Samaritan Healthcare

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Samaritan Healthcare **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Skagit Valley Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Skagit Valley Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Swedish Medical Center - Ballard

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Swedish Medical Center - Edmonds

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Swedish Medical Center – First Hill

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Swedish Medical Center – First Hill **2** Washington State **2**

EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.


Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Swedish Medical Center - Issaquah

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen




Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Three Rivers Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss

Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Three Rivers Hospital **2** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Trios Health Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss

Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



University of Washington Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



University of Washington Medical Center - Northwest

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital's screening numbers with the State's overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



University of Washington Medical Center – Valley Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.


Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.


Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.


Diagnosed with Hearing Loss




Diagnosed with Hearing Loss by 3 Months



No Hearing Loss




Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Virginia Mason Franciscan Health – St. Anne Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Virginia Mason Franciscan Health St. Anne **1** Washington State **2**

EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.


Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Virginia Mason Franciscan Health – St. Elizabeth Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss

Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Virginia Mason St. Elizabeth Hospital **2** Washington State **2**

EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Virginia Mason Franciscan Health – St. Francis Hospital

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Virginia Mason Franciscan St. Francis **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Virginia Mason Franciscan Health – St. Joseph Medical Center Tacoma

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)

Virginia Mason St. Joseph MC Tacoma **1** Washington State **2**

EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Virginia Mason Franciscan Health – St. Michael Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019




Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.

Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals

Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.

Diagnosed with Hearing Loss



Diagnosed with Hearing Loss by 3 Months



No Hearing Loss



Pending Diagnosis



Lost -- No Diagnosis



Diagnosis Status is 'Other'



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



WhidbeyHealth Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Whitman Hospital and Medical Center

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss

Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Yakima Valley Memorial

Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019



Washington State
Department of Health
EHDDI Program

Phone: 1-888-923-4334
Email: ehddi2@doh.wa.gov
Web: www.doh.wa.gov/earlyhearingloss


Newborn Hearing Screening

Hospital newborn hearing screening programs are essential for the early identification of infants who are deaf or hard of hearing. With your help, the Early Hearing Detection, Diagnosis, and Intervention (EHDDI) program can accurately track infants born in Washington State to ensure they are screened for hearing loss and receive appropriate follow-up care. The following table compares your hospital’s screening numbers with the State’s overall statistics.


Received Hearing Screen



Referred Initial Hearing Screen



Lost after Missed or Referred Initial Hearing Screen



Average Age of Initial Screen (days)



EHDDI Goals


Infants receive hearing screen before hospital discharge or by **1** month of age.

Infants who not pass two hearing screens have a diagnostic evaluation before they are **3** months of age.

Infants identified as deaf or hard of hearing start early intervention services by **6** months of age.

Diagnostic Evaluation

Infants who are identified as deaf or hard of hearing early and enrolled into early intervention services by six months of age are more likely to have age appropriate language and social-emotional development. The table below summarizes information reported to the EHDDI program for infants who were referred for diagnostic testing.



*Other: Infants whose parents declined, deceased infants, medically fragile infants and infants that have moved out of state.

The EHDDI program appreciates the work you do to ensure infants receive quality screening and care. We hope this information assists you in your efforts to make your UNHS program as successful as possible.

Guidelines for an Optimal Test

- Quiet room
- Quiet sleeping baby
- Ideally, conducted after baby has been fed
- Clean diaper
- Sign posted to inform others a hearing test is in progress
- Establish test times that are optimal for nurses, parents, and babies' schedules



Universal Newborn Hearing Screening (UNHS) Hospital Summary Report for Infants Born in 2019

To request this document in another format, call 1-800-525-0127. Deaf or hard of hearing customers, please call 711 (Washington Relay) or email civil.rights@doh.wa.gov.



DOH 344-076 July 2021