

# Fax Page 1 To: Adams County Public Health

Adapted from WA DOH Form 347-102

Ritzville: (509) 659-4109 / Othello: (509) 331-0030 (Confidential FAX line) CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Report STIs within 3 work days (WAC 246-101-101/301)

	PATIENT INFORMATION													
LAST NAME			FIRST NAME			MIDDLE NAME DATE OF BIRTH								
						1			мо	l	DAY	1	YR	
ADDRESS ( Unhoused or unstably housed			in the past 3 months)		СІТҮ				STATE		ZIP COD	DE		
TELEPHONE		EMAIL			ENGLIS	SH SPEAKING	6? 🗆 Yes 🗌 N	o *instructions	<sup>s on</sup> C		IS DA	ATE		
()					Prefer	red Language	e (Code*: L	pg. 3		10	DA	Y	YR	
SEX ASSIGNED	GENDER IDEN	ΓΙΤΥ		ETHNIC			RACE CATEGO	, RY (check all						s on page 3
AT BIRTH	□ Male	Transgen	der MTF	☐ Hisp	anic or	Latina/o/x	🗌 White			,		🗌 Asiai	n	
□ Male □ Female □ Transgen			der FTM 🗌 Non-Hispanic				Black     Other       American Indian / Alaska Native     Unknown							
Female	□ Nonbinary ,						□ American Indian / Alaska Native □ Onknown							
☐ Intersex Genderqueer ☐ Refused			Refused		EXTENDED RACE CODE		•							
CURRENTLY	REASON FOR E	XAM (check one)	GEND	ER OF S	EX PAR	TNERS (check	all that apply):	HIV STATU				Case Report		RENTLY
PREGNANT?	Exposed to I	Infection				Previo	us po	sitive				rEP?		
🗌 Yes	Symptomati	с	🗌 🗌 Fen			Transge	nder FTM		ew HIV diagnosis at this visit*					
□ No □ Unk. □ NA	□ Routine Exa	m (No Symptoms	·/	nbinary		Other		-	Negative HIV test at this visit I No Did not test (unknown status) I NA					
			Ger	nderque			V11			UNIKIUW	11 36			
DIAGNOSIS - I GONORRHEA (I								sv	PHILI	c				
DIAGNOSIS (ch		SITES (all that	apply):	TREATM	AENT (c	heck all pres	cribed).				ne).			
Asymptomati		Cervix	It apply):       TREATMENT (check all prescribed):         Ceftriaxone:       250 mg       500 mg       1 g         Ceftriaxone:       400 mg       800 mg				□1g □	STAGE (check one):						
Symptomatic	, Uncomplicated	🗌 Urethra							ndary (Ra					
<ul> <li>Pelvic Inflami</li> <li>Ophthalmia</li> </ul>	matory Disease	Urine	Azithromycin: 1 g 2 g						Latent (<					
Disseminated	k	Pharynx	<ul> <li>Doxycycline: 100 mg BID x 7 days</li> <li>Gentamicin: 240 mg</li> </ul>					<ul> <li>Unknown Duration or Late</li> <li>Congenital</li> </ul>						
🗌 Other Compl	ications:	🗌 Vagina	Gemifloxacin: 320 mg				м	MANIFESTATIONS (check all that apply):						
		Ocular	□ Other:					□ Neurologic □ Otic □ Ocular □ Tertiary						
Date Tested:		Other:	Date Prescribed:											
CHLAMYDIA (la				TOFATA		heek ell mus	entile e al \.			/IENT (ch		•		
DIAGNOSIS (ch	-	SITES (all that	t apply): TREATMENT (check all prescribed):  Azithromycin: 1 g Doxycycline: 100 mg BID x 7 days Levofloxacin: 500 mg daily x 7 days Other:				Bio	Bicillin L - A: 2.4 MU IM x 1 2.4 MU IM x 3 Doxycycline: 100 mg BID x 14 days 100 mg BID x 28 days Benzathine 50,000 units/kg IM x 1						
_ / !	c, Uncomplicated						De							
Pelvic Inflam	matory Disease	🗌 Urine												
<ul> <li>Ophthalmia</li> <li>Other Compl</li> </ul>	lications:	Rectum					Ве							
Other Complications:     □ Pharynx     □ Vagina							PC	PCN-G:         50,000 units/kg IM x 3           Aqueous         18-24 MU/day IV				3		
🗌 Ocular														
Date Tested: Other:			Date Prescribed:					Crystalline for 10-14 days Penicillin G:						
HERPES SIMPLE	EX	LABORATORY (	ONEIDAAA			DISEASES			her:					
	al infection only)		UNFIRIVIA		Cha	ncroid nuloma Ingui	nale		_					
Neonatal		🗌 No			Lym	phogranulon	na Venereum	Da	ate Pr	escribed:				
PARTNER TREATMENT PLAN (check one or more options)														
for additional inf	ormation).	treatment by eith	C				:	•		0		•	•	
for additional information).  In-person evaluation - Number of partners treated following medical evaluation:														
<ul> <li>Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis</li> </ul>														
REPORTING CLINIC INFORMATION														
	FACILITY N/	AME				DIAGN	IOSING CLINICI	AN						
DATE														
ADDRESS						CITY			STATE		Z	ΊP		
	LETING FORM			TEL	EPHON			EMAIL			Z	ÎP		
ADDRESS	LETING FORM			TEL							Z	(IP		

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile.

# **Gonorrhea or Chlamydia Infection: Partner Treatment**

#### All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Adams County Public Health may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Adams County Public Health recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Adams County Public Health: Ritzville: (509) 659-3315/ Othello: (509) 488-2031.

# Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

# **RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS\***

### **GONORRHEA** -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs)<sup> $\dagger$ </sup>

#### Alternatives for uncomplicated infections of the cervix, urethra, or rectum:<sup>‡</sup>

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose OR

Cefixime 800 mg orally as a single dose<sup>†</sup>

<sup>+</sup> If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.

<sup>‡</sup> Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

## CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

#### Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days **OR** 

Ofloxacin 300 mg PO BID for 7 days **OR** 

Levofloxacin 500 mg PO for 7 days

# SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

## SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

\* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>https://www.cdc.gov/std/treatment/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

## List of Preferred Languages:

**Instructions:** Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

#### Language Identity Codes:

(L01) Amharic	(L18) Karen	(L35) Sign languages
(LO2) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(LO4) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown
(L16) Hmong	(L33) Russian	
(L17) Japanese	(L34) Samoan	

### **Race Category and Identity Instructions:**

**Instructions:** The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

#### **Extended Race Codes:**

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	