

PATIENT INFORMATION

Fax Page 1 To: **Clallam County Dept. of Health & Human Services** Port Angeles: (360) 452-4492 (Confidential FAX line) CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Report STIs within 3 work days (WAC 246-101-101/301)

LAST NAME			FIRST NAME			MIDDLE NAME		DATE OF BIRTH				
								мо	I	DAY		YR
ADDRESS (Unhoused or unstably housed			d in the past 3 months) CITY					STATE ZIP COD		E		
TELEPHONE		EMAIL		ENGL	SH SPEAKING	6? 🗆 Yes 🗌 No	*instructions	on D		S DAT	ΓE	
()			Pre		ferred Language (Code*: L		pg. 3)	м	0	DAY	Í	YR
() SEX ASSIGNED AT BIRTH Alle Female Intersex Refused CURRENTLY PREGNANT? Yes No Unk. NA DIAGNOSIS - E	Exposed to I Symptomatic Routine Exar	☐ Transgend ☐ Transgend / ☐ Other er ☐ Refused XAM (check one): nfection	der MTF	HNICITY Hispanic or Non-Hispan Unknown Refused OF SEX PAF	r Latina/o/x nic	RACE CATEGORY White Black American India Native Hawaii EXTENDED RACE all that apply): r nder MTF nder FTM	an / Alaska an / Other	that a Pacif *: R S Is pos V dia re HIV	ve ic Islander *Submit HIV/ sitive gnosis at / test at t	r /AIDS Ca this V	*Ins *Ins Asian Other Unkno Refus R asse Report visit* sit	structions on page 3
GONORRHEA (I							SY	PHILI	S			
 Asymptomatic Symptomatic, Uncomplicated Pelvic Inflammatory Disease Ophthalmia Disseminated Other Complications: Date Tested: CHLAMYDIA (lab confirmed) DIAGNOSIS (check one) Asymptomatic Symptomatic, Uncomplicated 		□ Cervix □ Ceftr □ Urethra □ Cefix □ Urine □ Azith □ Rectum □ Doxy □ Pharynx □ Gent. □ Ocular □ Other □ Other: □ Date Pro SITES (all that apply): TREATM □ Cervix □ Azith □ Urethra □ Doxy		Ceftriaxone Cefixime: Azithromyci Doxycycline Gentamicin Gemifloxaci Other: ate Prescribe	time: 400 mg 800 mg promycin: 1 g 2 g vcycline: 100 mg BID x 7 days ramicin: 240 mg ifloxacin: 320 mg er:			 STAGE (check one): Primary (Chancre, etc.) Secondary (Rash, etc.) Early Latent (< 1 year) Unknown Duration or Late Congenital MANIFESTATIONS (check all that apply): Neurologic Otic Ocular Tertiary TREATMENT (check one): Bicillin L - A: 2.4 MU IM x 1 2.4 MU IM x 3 Doxycycline: 100 mg BID x 14 days 100 mg BID x 28 days 				
Other Complications:		 Rectum Pharynx Vagina Ocular Other: 	Other: Date Prescribed: OTHER DISEASES			PCI Aq Cry	N-G: ueous vstallii	athine 50,000 units/kg IM x 1 G: 50,000 units/kg IM x 3				
		🗌 No		DN Cha Gra Lyn	ncroid nuloma Inguinale			Other: Date Prescribed:				
		(check one or										
for additional info In-person eva	ormation). Aluation - Number ered treatment* -	of partners treat	ed following r ers for whom	nedical evalu provider pre	uation:	scribing medication	Turn	over fo	or Partner Ti EPT) medi	reatme icatio	ent Plan In n pack to	structions D
REPORTING CLINIC INFORMATION												
DATE	FACILITY NA	AME			DIAGN	IOSING CLINICIAN						
ADDRESS					CITY		S	TATE		ZII	P	
PERSON COMPLETING FORM				TELEPHONE EM			EMAIL	/AIL				

Thank you for reporting an STI. All information will be managed with the strictest confidentiality.

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile.

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Clallam County Dept. of Health & Human Services may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Clallam County Dept. of Health & Human Services recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Clallam County Dept. of Health & Human Services: Port Angeles: (360) 565-2612.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs)^{\dagger}

Alternatives for uncomplicated infections of the cervix, urethra, or rectum:[‡]

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose OR

Cefixime 800 mg orally as a single dose[†]

⁺ If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.

[‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days **OR**

Ofloxacin 300 mg PO BID for 7 days **OR**

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>https://www.cdc.gov/std/treatment/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

List of Preferred Languages:

Instructions: Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

(L01) Amharic	(L18) Karen	(L35) Sign languages
(LO2) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(LO4) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown
(L16) Hmong	(L33) Russian	
(L17) Japanese	(L34) Samoan	

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

Extended Race Codes:

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	