

Fax Page 1 To: Grant County Health District (509) 764-2813 (Confidential FAX line)

Adapted from WA DOH Form 347-102

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Report STIs within 3 work days (WAC 246-101-101/301)

PATIENT INFORMATION														
LAST NAME			FIRST NAME			MIDDLE NAM	E DATE OF BIRTH							
									MO DAY YR			YR		
ADDRESS (Unhoused or unstably housed			in the past 3 months)		CITY				STATE		ZIP COD	E		
TELEPHONE		EMAIL			ENGLI	SH SPEAKING	6? 🗌 Yes 🗌 N	o *instruction	s on C		IS DA	ATE		
()					Prefer	red Languag	e (Code*: L	pg. 3	•	10	DA	Y	YR	
SEX ASSIGNED	GENDER IDEN	ΓΙΤΥ		ETHNI			RACE CATEGO	RY (check al	l that	apply)*:		*In	structions	on page 3
AT BIRTH	🗌 Male	🗌 Transgeno	der MTF	🗌 Hisp	oanic or	Latina/o/x	U White					🗌 Asiar		
Male	□ Male □ Female □ Transgen		der FTM 🛛 🗌 Non-Hispani			nic 🛛 🗌 Black		ndian / Alask	Alaska Native 🗌 Unknown					
Female Intersex Genderqueer Refused		Unknown				Native Hawaiian / Other Pacific Islander			er	Refus				
Refused	Genderque	er 🗌 Refused			iseu		EXTENDED RA	CE CODE(S))*: F	R		R	R	R
CURRENTLY	REASON FOR E	XAM (check one)	GEND	DER OF S	EX PAR	TNERS (check	all that apply):	HIV STATU	JS	*Submit HI	//AIDS	Case Report	CURF	RENTLY
PREGNANT?	Exposed to	nfection		☐ Male ☐ Transger			nder MTF 🗌 Previous positive ON PrEP?							
Yes	^{′es} ☐ Symptomatic ☐ Female ☐ 1			_ 0	nsgender FTM IN New HIV diagnosis at this visit* IN Yes ner IN Negative HIV test at this visit IN NO									
□ No □ Unk. □ NA	□ Routine Exa	m (No Symptoms	., .	onbinary nderque		Other Unknov	vn	Did no						
				mucrqui			vii							
GONORRHEA (I								SY	'PHILI	S				
DIAGNOSIS (ch		SITES (all that	apply):	TREAT	VIENT (c	heck all pres	cribed):			check or	ne):			
Asymptomati	ic	🗌 Cervix	\square Ceftriaxone: \square 250 mg \square 500 mg \square 1 g				ח1g □	Primary (Chancre, etc.)						
Symptomatic	, Uncomplicated	Urethra	Cefixime: 400 mg 800 mg						ndary (Ra Latent (<					
Ophthalmia	liatory Disease	Rectum	□ Azithromycin: □ 1 g □ 2 g □ Doxycycline: □ 100 mg BID x 7 days						own Dura					
Disseminated		Pharynx	Gentamicin: 240 mg					Congenital						
🗌 Other Compl	ications:	Vagina	🔲 Gemifloxacin: 🔲 320 mg				м	MANIFESTATIONS (check all that apply):						
Coular		Ocular	Other: Date Prescribed:				🗆	□ Neurologic □ Otic □ Ocular □ Tertiary						
Date Tested:	h and firm a d)		i	Date P	rescribe	a:						,		
CHLAMYDIA (la DIAGNOSIS (ch	-	SITES (all that	apply):	TREAT	MENT (c	heck all pres	cribed):			/IENT (ch		•		
Asymptomat	-	Cervix	Azithromycin: 1 g					Bicillin L - A: 2.4 MU IM x 1 2.4 MU IM x 3						
	, Uncomplicated		Doxycycline: 100 mg BID x 7 days				Do	Doxycycline: 🗌 100 mg BID x 14 days						
 Pelvic Inflam Ophthalmia 	matory Disease	Urine	 Levofloxacin: 500 mg daily x 7 days Other: 				□ 100 mg BID x 28 days							
Other Compl	ications:						Benzathine 🗌 50,000 units/kg IM x 1							
Vagina							PC	PCN-G: 50,000 units/kg IM x 3						
Date Tested:		Date Prescribed:					Aqueous I 18-24 MU/day IV Crystalline for 10-14 days							
Date Tested: Other: Other:			OTHER DISEASES					Crystalline for 10-14 days Penicillin G:						
DIAGNOSIS		LABORATORY C	ONFIRMA	TION				Ot	ther: _					
Genital (initia	I infection only)				🗌 Gra	nuloma Ingui		D	ate Pr	escribed:				
Neonatal DAPTNER TRE		l 🗌 No	more on	tions		phogranulon	a venereum			coenocu.				
	PARTNER TREATMENT PLAN (check one or more options) Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2													
for additional information).														
	In-person evaluation - Number of partners treated following medical evaluation:													
				 Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis 										
Patient-delive	ered treatment*	Number of partn	ers for wh										its with	<i>,</i> ,
Patient-delive	ered treatment* - the patient to the	Number of partneir partneir partner (s):	ers for wh										its with	
Patient-delive delivered by t	ered treatment* - the patient to the	Number of partnering partnering partneries (s):	ers for wh			d treatment is		d for men w					its with	
Patient-delive delivered by t REPORTING C	ered treatment* the patient to the LINIC INFORM	Number of partnering partnering partneries (s):	ers for wh			d treatment is	not recommende	ad for men w		ve sex with	h mer		its with	
Patient-delive delivered by 1 REPORTING CI DATE	ered treatment* - the patient to the LINIC INFORM FACILITY N/	Number of partnering partnering partneries (s):	ers for wh	*Patient-	delivered	d treatment is DIAGN CITY	not recommende	ad for men w	state	ve sex with	h mer	n or patier	its with	
Patient-delive delivered by 1 REPORTING CI DATE ADDRESS	ered treatment* - the patient to the LINIC INFORM FACILITY N/	Number of partnering partnering partneries (s):	ers for wh	*Patient-		d treatment is DIAGN CITY	not recommende	ad for men w	state	ve sex with	h mer	n or patier	its with	

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile.

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Grant County Health District may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Grant County Health District recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Grant County Health District: (509) 766-7960 ext.13.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs)^{\dagger}

Alternatives for uncomplicated infections of the cervix, urethra, or rectum:[‡]

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose OR

Cefixime 800 mg orally as a single dose[†]

⁺ If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.

[‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days **OR**

Ofloxacin 300 mg PO BID for 7 days **OR**

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>https://www.cdc.gov/std/treatment/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

List of Preferred Languages:

Instructions: Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

(L01) Amharic	(L18) Karen	(L35) Sign languages
(LO2) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(LO4) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown
(L16) Hmong	(L33) Russian	
(L17) Japanese	(L34) Samoan	

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

Extended Race Codes:

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	