

# Fax Page 1 To:

# **Kitsap County Health District**

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Adapted from WA DOH Form 347-102 (360) 813-1168 (Confidential FAX line)

		N.	port 511	S WILIIII	3 WO	rk days (w	AC 246-101-10	11/301)					
PATIENT INFO	RMATION												
LAST NAME			FIRST NAME			MIDDLE NAME		DAT	DATE OF BIRTH				
								МО		DAY		YR	
ADDRESS ( Unhoused or unstably housed			in the past 3 months) CITY						STATE		ZIP COD	E	
TELEPHONE EMAIL			ENGLISH SPEAKING?			6? ☐ Yes ☐ N	O *instruction pg. 3	s on D	IAGNOS	IS DA	NTE		
( )			Preferred Language			e (Code*: L	)	M	10	DA	Y	YR	
SEX ASSIGNED AT BIRTH  Male Female Intersex	BIRTH     □ Male     □ Transgender M       Male     □ Female     □ Transgender F       Female     □ Nonbinary /     □ Other			r MTF Hispanic or Latina/o/x r FTM Non-Hispanic Unknown			RACE CATEGORY (check all that apply)*:  White Asian Black Other American Indian / Alaska Native Unknown Native Hawaiian / Other Pacific Islander Refused						
Refused				E			EXTENDED RACE CODE(S)*: R R R R				R R		
CURRENTLY PREGNANT?  Yes No Unk. NA	☐ Exposed to I	osed to Infection ptomatic utine Exam (No Symptoms)			DER OF SEX PARTNERS (check a lale Transgen male Transgen onbinary / Other landerqueer Unknown		nder MTF nder FTM	☐ New H	· · · · · · · · · · · · · · · · · · ·			□ No	
DIAGNOSIS - D	DISEASE												
GONORRHEA (la	ab confirmed)							S۱	SYPHILIS				
DIAGNOSIS (check one)  Asymptomatic  Symptomatic, Uncomplicated  Pelvic Inflammatory Disease  Ophthalmia  Disseminated  Other Complications:		SITES (all that apply):  Cervix Urethra Urine Rectum Pharynx Vagina Coular Other:		TREATMENT (check all prescribed):  Ceftriaxone: 250 mg 500 Cefixime: 400 mg 800 Azithromycin: 1 g 2 g Doxycycline: 100 mg BID x 7 dd Gentamicin: 240 mg Gemifloxacin: 320 mg Other:			500 mg [ 800 mg ] 2 g BID x 7 days	1 g	STAGE (check one):  Primary (Chancre, etc.)  Secondary (Rash, etc.)  Early Latent (< 1 year)  Unknown Duration or Late  Congenital  MANIFESTATIONS (check all that apply):  Neurologic Otic Ocular Tertiany				
Date Tested:		! — other	i	Date Pre	escribe	d:							
CHLAMYDIA (lab confirmed)  DIAGNOSIS (check one)  Asymptomatic  Symptomatic, Uncomplicated  Pelvic Inflammatory Disease  Ophthalmia  Other Complications:  Planynx  Vagina			TREATMENT (check all pres   Azithromycin:   1 g   Doxycycline:   100 mg   Levofloxacin:   500 mg   Other:			g BID x 7 days	Bi Do Be	TREATMENT (check one):  Bicillin L - A:				8 days g IM x 1	
Date Tested: Ocular Other:			Date Prescribed:					Aqueous 18-24 MU/day IV Crystalline for 10-14 days			IV		
HERPES SIMPLEX DIAGNOSIS  Genital (initial infection only) Neonatal  LABORATORY CO			OTHER DISEASES				O:	Crystalline for 10-14 days Penicillin G: Other:  Date Prescribed:					
PARTNER TREATMENT PLAN (check one or more options)													
Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2 for additional information).  In-person evaluation - Number of partners treated following medical evaluation:  Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s):  *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis													
REPORTING CLINIC INFORMATION													
DATE													
ADDRESS				CITY					STATE ZIP				
PERSON COMPLETING FORM				TELEPHONE ( )			EMAIL	EMAIL					

#### PARTNER MANAGEMENT PLAN INSTRUCTIONS

# **Gonorrhea or Chlamydia Infection: Partner Treatment**

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Kitsap County Health District may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Kitsap County Health District recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Kitsap County Health District: (360) 728-2235.

#### Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

# RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS\*

# **GONORRHEA -- Uncomplicated**

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs) $^{\dagger}$ 

## Alternatives for uncomplicated infections of the cervix, urethra, or rectum: ‡

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose  $\bf OR$  Cefixime 800 mg orally as a single dose  $^{\dagger}$ 

- † If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.
- <sup>‡</sup> Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

#### **CHLAMYDIA** -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

#### Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days OR

Ofloxacin 300 mg PO BID for 7 days OR

Levofloxacin 500 mg PO for 7 days

#### SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

# **SYPHILIS -- LATE OR UNKNOWN DURATION**

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

<sup>\*</sup> Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<a href="https://www.cdc.gov/std/treatment/default.htm">https://www.cdc.gov/std/treatment/default.htm</a>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

# **List of Preferred Languages:**

Instructions: Complete the Preferred Language section if the patient's preferred language is not English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

# **Language Identity Codes:**

(L01) Amharic	(L18) Karen	(L35) Sign languages
(L02) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(L04) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown

(L33) Russian

(L34) Samoan

# **Race Category and Identity Instructions:**

Instructions: The patient's race may either be reported in the general Race Category section or the Extended Race section on page 1 of the case report. When completing the Extended Race section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the extended race section. Unknown race can be documented within either one of the race sections.

## **Extended Race Codes:**

(L16) Hmong

(L17) Japanese

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	