

## Fax Page 1 To:

# **Skamania County Health Department**

(509) 427-0188 (Confidential FAX line)

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Adapted from WA DOH Form 347-102

		R	eport ST	ls within	1 3 wo	rk days (W	AC 246-101-10:	1/301)					
PATIENT INFO	RMATION												
			FIRST NAME					DATI MO	DATE OF BIRTH				
455556 /	7							<u>                                     </u>			DAY		YR
ADDRESS ( Unhoused or unstably housed in			in the pa	in the past 3 months) CITY						STATE		ZIP COD	Ē
TELEPHONE		EMAIL		ENGLISH SPEAKING?			G? ☐ Yes ☐ No	? ☐ Yes ☐ No *instructions on DIAGNOSIS DATE					
( )			Preferred Language			e (Code*: L )		м	MO DAY YR		YR		
SEX ASSIGNED	GENDER IDENT	TITY				RACE CATEGORY (check all that apply)*:					structions on page 3		
AT BIRTH ☐ Male ☐ Transgend							<ul><li>☐ White</li><li>☐ Black</li></ul>			☐ Asian ☐ Other			
☐ Male ☐ Female ☐ Transgende						nic	☐ American Indian / Alaska N						
☐ Female ☐ Nonbinary / ☐ Other ☐ Intersex ☐ Genderqueer ☐ Refused			☐ Unknown ☐ Refused				☐ Native Hawaiian / Other P			<del>_</del>			
☐ Intersex Genderqueer ☐ Refused ☐ Refused				EXTENDED RAC			CE CODE(S)	CODE(S)*: R R R R				R R	
CURRENTLY	REASON FOR E	XAM (check one):	GENI	DER OF SI	EX PAR	TNERS (check	all that apply):	HIV STATU	<u></u>	*Submit HIV	/AIDS	Case Report	CURRENTLY
PREGNANT?	REGNANT?							☐ Previou	Previous positive ON PrEP?				
☐ Yes	Symptomatic	С	☐ Female ☐ Tra			_	~			diagnosis at this visit*			
□ No	☐ Routine Exar	n (No Symptoms				☐ Other	I			HIV test at this visit ☐ No est (unknown status) ☐ NA			1 —
☐ Unk. ☐ NA			G	enderque	er	Unknov	vn	□ Dia not	test	unknow	n Sta	atus)	□NA
DIAGNOSIS - DIAGNORRHEA (I								CV	211111				
DIAGNOSIS (che		CITES (all that	annlul.					SYPHILIS					
☐ Asymptomati	•	Cervix				REATMENT (check all prescribed):    Ceftriaxone: □ 250 mg □ 500 mg □ 1 g			STAGE (check one):  Primary (Chancre, etc.)				
Symptomatic	□ Urethra	를 ** : '' : ''			☐ Cefixime: ☐ 250 mg ☐ 500 mg			Secon	dary (Ras	sh, et	tc.)		
Pelvic Inflamr	natory Disease	☐ Urine			☐ Azithromycin: ☐ 1 g ☐ 2 g				☐ Early Latent (< 1 year)				
☐ Ophthalmia ☐ Disseminated		Rectum	☐ Doxycycline: ☐ 100 mg I					BID X / days		known Duration or Late ngenital			
☐ Other Compli		☐ Pharynx☐ Vagina	☐ Gentamicin: ☐ 240 mg ☐ Gemifloxacin: ☐ 320 mg			-							
		Ocular	☐ Gemilioxacin: ☐ 320 mg			IVIAINI			IIFESTATIONS (check all that apply):				
Date Tested:		☐ Other:	Othor:		ate Prescribed:			'	☐ Neurologic ☐ Otic ☐ Ocular ☐ Tertiary				
CHLAMYDIA (la	b confirmed)							TR	EATIV	IENT (ch	eck o	one):	
DIAGNOSIS (che		SITES (all that	TREATMENT (check all prescribed):			Bic	Bicillin L - A: ☐ 2.4 MU IM x 1						
Asymptomatic		☐ Cervix		☐ Azithromycin: ☐ 1 g				☐ 2.4 MU IM x 3					
<ul><li>Symptomatic, Uncomplicated</li><li>Pelvic Inflammatory Disease</li></ul>		☐ Urethra☐ Urine		☐ Doxycycline: ☐ 100 mg E☐ Levofloxacin: ☐ 500 mg G☐ Doxycycline: ☐ 50						ycycline: 🔲 100 mg BID x 14 days			
Ophthalmia		I □ Orine		l □ Levofloxacin: □ 500 mg d						$\Box$ 100 mg BID x 28 days			
☐ Other Complications:		☐ Pharynx							Benzathine 50,000 units/kg IM x PCN-G: 50,000 units/kg IM x				
——————— ☐ Vagina												_	
Date Tested:		☐ Ocular ☐ Other:		Date Prescribed:				Aqueous					
HERPES SIMPLE	x	- other				R DISEASES		'	nicillir		01 10	0-14 days	
DIAGNOSIS LABORATORY CONFIRMATION Chancroid Other:													
☐ Genital (initial infection only) ☐ Yes☐ Neonatal ☐ No			☐ Granuloma Inguinale ☐ Lymphogranuloma Venereum			Da	Date Prescribed:						
PARTNER TREATMENT PLAN (check one or more options)													
Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2 for additional information).    In-person evaluation - Number of partners treated following medical evaluation:    Turn over for Partner Treatment Plan Instructions   Turn over for Partner Treatment Plan Instructions													
☐ In-person eva	luation - Number	of partners treat	ed followi	ng medica	al evalu	ation:	!	Turn	over fo	r Partner T	reatn	nent Plan In	structions
☐ Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis													
REPORTING CLINIC INFORMATION													
DATE	TE FACILITY NAME DIAGNOSING CLINICIAN												
ADDRESS				CITY				STATE			Z	ZIP	
PERSON COMPLETING FORM			TEL	TELEPHONE EN			EMAIL	I ΛΑΙL					
					/								

#### PARTNER MANAGEMENT PLAN INSTRUCTIONS

## **Gonorrhea or Chlamydia Infection: Partner Treatment**

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Skamania County Health Department may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Skamania County Health Department recommends you refer all MSM patients and all patients with syphilis or newly diagnosed HIV to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Skamania County Health Department: (509) 427-3881.

## Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

## RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS\*

## **GONORRHEA -- Uncomplicated**

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs) $^{\dagger}$ 

## Alternatives for uncomplicated infections of the cervix, urethra, or rectum: ‡

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose  $\bf OR$  Cefixime 800 mg orally as a single  $\bf dose^{\dagger}$ 

- † If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.
- <sup>‡</sup> Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

#### **CHLAMYDIA** -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR**Azithromycin 1g PO as a single dose

#### Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days  ${\bf OR}$ 

Ethylsuccinate 800 mg PO QID for 7 days OR

Ofloxacin 300 mg PO BID for 7 days OR

Levofloxacin 500 mg PO for 7 days

#### SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

## **SYPHILIS -- LATE OR UNKNOWN DURATION**

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

<sup>\*</sup> Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<a href="https://www.cdc.gov/std/treatment/default.htm">https://www.cdc.gov/std/treatment/default.htm</a>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

## **List of Preferred Languages:**

Instructions: Complete the Preferred Language section if the patient's preferred language is not English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

## **Language Identity Codes:**

(L01) Amharic	(L18) Karen	(L35) Sign languages
(L02) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(L04) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown

(L33) Russian

(L34) Samoan

## **Race Category and Identity Instructions:**

Instructions: The patient's race may either be reported in the general Race Category section or the Extended Race section on page 1 of the case report. When completing the Extended Race section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the extended race section. Unknown race can be documented within either one of the race sections.

## **Extended Race Codes:**

(L16) Hmong

(L17) Japanese

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	