

# Fax Page 1 To: Wahkiakum County Department of Health (360) 795-6143 (Confidential FAX line)

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Report STIs within 3 work days (WAC 246-101-101/301)

PATIENT INFO	RMATION						1						PATIENT INFORMATION							
LAST NAME			FIRST NAME			MIDDLE NAM	1E	DATE OF BIRTH												
									мо	DAY		YR								
ADDRESS ( Unhoused or unstably housed			in the past 3 months)		СІТҮ				STATE		ZIP COD	E								
TELEPHONE EMAIL			ENGLISH SPEAKIN			SH SPEAKING	6?  Yes No *instructions on DIAGNOSIS DAT		TE											
( )		Prefer		red Language	ge (Code*: L )			0	DA	Y I	YR									
SEX ASSIGNED	GENDER IDEN	ΓΙΤΥ		ETHNICITY RACE CATEGORY (che			RY (check all	that	apply)*:		*in	structions on page 3								
AT BIRTH			ler MTF					🗌 Asian												
🗆 Male	] Male 🛛 🗌 Female 🔅 Transgen		der FTM 🛛 🗆 Non-Hispan			ic	Black			☐ Other Jaska Native ☐ Unknown										
Female Nonbinary / Other		Unknown				□ Native Hawaiian / Other Pacific														
□ Intersex Genderqueer □ Refused □ Refused			EXTENDED RACE COD			CE CODE(S)	E(S)*: R R R R R													
CURRENTLY	REASON FOR E	XAM (check one)	GEND	DER OF SE	X PAR	TNERS (check	all that apply):	HIV STATU	S	*Submit HI\	//AIDS	Case Report	CURRENTLY							
PREGNANT?							Previou	revious positive ON PrEP?												
🗌 Yes	es Symptomatic					Transge				liagnosis at this visit*										
□ No □ Unk. □ NA			-	Negative HIV test at this visit     Image: No       Did not test (unknown status)     Image: No																
			Ge	nderque	ei	Unknov			i iest	UNKIOW	ni Sta	itusj								
DIAGNOSIS - E GONORRHEA (I								CV	PHIL	16										
		SITES (all that		TREATM		hock all proc	cribod).													
Asymptomati				t apply): TREATMENT (check all prescribed):					STAGE (check one):											
Symptomatic	, Uncomplicated	🗌 Urethra	$\Box Cefixime: \Box 400 mg \Box 800 mg$					Seco	ndary (Ra	sh, et	tc.)									
Pelvic Inflammatory Disease     Urine						n: 🗌 1 g	🗌 2 g			Latent (<										
Ophthalmia     Disseminated     Disseminated		Pharynx	Doxycycline: 100 mg BID x 7 days					Unknown Duration or Late												
🗌 Other Compli	ications:	Vagina	☐ Gentamicin: ☐ 240 mg ☐ Gemifloxacin: ☐ 320 mg					-												
		🗌 Ocular	Other:					MANIFESTATIONS (check all that apply):												
Date Tested:		Other:		Date Prescribed:					□ Neurologic □ Otic □ Ocular □ Tertiary											
CHLAMYDIA (la								TR	EAT	VENT (ch	eck d	one):								
DIAGNOSIS (check one) SITES (all tha							Bio	Bicillin L - A: 🗌 2.4 MU IM x 1												
Asymptomatic Cervix Symptomatic, Uncomplicated Ureth			Azithromycin: 1 g					2.4 MU IM x 3												
Pelvic Inflam			Levofloxacin: 500 mg daily x 7 days				Do	Doxycycline: 100 mg BID x 14 days 100 mg BID x 28 days												
Ophthalmia	ications	🗌 Rectum	□ Other:			Be	Benzathine $\Box$ 50,000 units/kg IM x 1													
Other Complications:  Pharynx							N-G:			00 units/k										
└── Vagina └── Vagina └── Ocular						Aq	Aqueous 18-24 MU/day IV													
Date Tested: Other:		Date Prescribed:					ystall	ine		0-14 days										
HERPES SIMPLE	X		·		OTHER	R DISEASES			nicilli											
DIAGNOSIS LABORATORY (			CONFIRMATION Chancroid					Other:												
	in inflection only)					phogranulom		Da	ite Pr	escribed:										
PARTNER TRE	ATMENT PLAN	(check one or	more op	tions)																
Providers should	manage partner	treatment by eith			in-per	son or by pres	cribing medicat	•		-		•								
for additional information).																				
In-person evaluation - Number of partners treated following medical evaluation:																				
delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis																				
REPORTING CLINIC INFORMATION																				
DATE FACILITY NAME DIAGNOSING CLINICIAN																				
ADDRESS						CITY		9	STATE	E	Z	IP								
PERSON COMPLETING FORM						-														
PERSON COMPL				TELEPHONE EN			EMAIL	AIL												
	Thank	you for reportin	ng an STI.	All inforn	nation	will be mana	aged with the s	trictest con	fider	ntiality.										

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile.

### **Gonorrhea or Chlamydia Infection: Partner Treatment**

#### All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Wahkiakum County Department of Health may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Wahkiakum County Department of Health recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Wahkiakum County Department of Health: (360) 795-6207.

## **Other STIs: Partner Treatment**

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

## **RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS\***

#### **GONORRHEA** -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs)<sup> $\dagger$ </sup>

#### Alternatives for uncomplicated infections of the cervix, urethra, or rectum:<sup>‡</sup>

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose OR

Cefixime 800 mg orally as a single dose<sup>†</sup>

<sup>+</sup> If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.

<sup>‡</sup> Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

#### CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

#### Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days **OR** 

Ofloxacin 300 mg PO BID for 7 days **OR** 

Levofloxacin 500 mg PO for 7 days

## SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

## SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

\* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>https://www.cdc.gov/std/treatment/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

### List of Preferred Languages:

**Instructions:** Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

#### Language Identity Codes:

(L01) Amharic	(L18) Karen	(L35) Sign languages
(LO2) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(LO4) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown
(L16) Hmong	(L33) Russian	
(L17) Japanese	(L34) Samoan	

#### **Race Category and Identity Instructions:**

**Instructions:** The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

#### **Extended Race Codes:**

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	