

Fax Page 1 To: Walla Walla County Health Department (509) 524-2642 (Confidential FAX line)

CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT

Report STIs within 3 work days (WAC 246-101-101/301)

PATIENT INFORMATION														
LAST NAME			FIRST NAME			MIDDLE NAM	/IE	DATE OF BIRTH						
							M		мо	D DAY YR		YR		
ADDRESS (Unhoused or unstably housed			l in the past 3 months)		СІТҮ				STATE		ZIP COD	E		
TELEPHONE							? Yes No *instructions on		s on C					
							pg. 3			MO DAY YR				
SEX ASSIGNED GENDER IDENTITY				Preferred Language (Code*: L) ETHNICITY RACE CATEGORY (che			, RY (check al		t apply)*: *Instructions on pa			on page 3		
AT BIRTH					Latina/o/x		(Asian			1			
🗆 Male						ic Black			Other					
Female Nonbinary Other			Unknown				□ American Indian / A							
□ Intersex Genderqueer □ Refused □ Refused			Refused EXTENDED RACE COD							R	R	R		
CURRENTLY	REASON FOR E	XAM (check one)	GEND	DER OF SE	EX PAR	TNERS (check	all that apply):	HIV STATU				Case Report	<u> </u>	RENTLY
PREGNANT?							Previo	revious positive ON PrEP?						
🗌 Yes	Yes Symptomatic		🗌 Female			Transge				diagnosis at this visit*				s
	□ No □ Routine Exam (No								ative HIV test at this visit I No not test (unknown status) I NA					
🗌 Unk. 🗌 NA			Ge	enderque	er	Unknov	vn	Did no	t test	(unknov	vn sta	itus)		4
DIAGNOSIS - E									DI					
GONORRHEA (I				TDEATA		h e els ell surs			PHIL					·
Asymptomati			at apply): TREATMENT (check all prescribed):					STAGE (check one):						
Symptomatic			☐ Ceftriaxone: ☐ 250 mg ☐ 500 mg ☐ 1 g ☐ Cefixime: ☐ 400 mg ☐ 800 mg						ndary (Ra					
Pelvic Inflammatory Disease		Urine	\square Azithromycin: \square 1 g \square 2 g						Early Latent (< 1 year)					
 Ophthalmia Disseminated 		Rectum	um 🗌 Doxycycline: 🗌 100 mg BID x 7 days						Unknown Duration or Late					
Other Compli		Pharynx												
		Ocular	Gemifloxacin: 320 mg				M	MANIFESTATIONS (check all that apply):						
Date Tested:		Date Prescribed:				0	□ Neurologic □ Otic □ Ocular □ Tertiary							
CHLAMYDIA (la	b confirmed)	i	i					ТБ	FAT	MENT (ch	eck o	ne).		
DIAGNOSIS (ch		SITES (all that	apply): TREATMENT (check all prescribed):						TREATMENT (check one): Bicillin L - A: 2.4 MU IM x 1					
Asymptomat		🗌 Cervix	☐ Azithromycin: ☐ 1 g					□ 2.4 MU IM x 3						
Symptomatic			Doxycycline: 100 mg BID x 7 days			Do	Doxycycline: 🗆 100 mg BID x 14 days							
 Pelvic Inflammatory Disease Ophthalmia 		□ Urine □ Rectum	 Levofloxacin: 500 mg daily x 7 days Other: 				□ 100 mg BID x 28 days							
Other Complications:			U Other:				Benzathine 🗌 50,000 units/kg IM x 1							
Vagina							PC	PCN-G: 50,000 units/kg IM x 3				3		
Deta Testada		Date Prescribed:					Aqueous I 18-24 MU/day IV Crystalline for 10-14 days							
Date Tested: Other:								nicilli		for 10	0-14 days			
HERPES SIMPLE DIAGNOSIS	:X	LABORATORY (ONEIRMA			DISEASES			her:					
Genital (initial infection only)						ncroid nuloma Inguinale							-	
🗌 Neonatal	-	🗌 No				phogranulor		Da	ate Pr	escribed:				
PARTNER TREATMENT PLAN (check one or more options)														
Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2 for additional information).														
	for additional information). In-person evaluation - Number of partners treated following medical evaluation: Turn over for Partner Treatment Plan Instructions													
		Number of partr						l partner the	rapy	(EPT) mee	dicatio	on pack to	o be	
delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis														
REPORTING CLINIC INFORMATION														
DATE FACILITY NAME DIAGNOSING CLINICIAN														
ADDRESS CITY STATE ZIP														
PERSON COMPL	PERSON COMPLETING FORM			TELI	EPHON ۱	E		EMAIL						
	There	Nou for some	0 0 0 CTI	All : f)	will he me		trictost and	f ial	Hallt				
Thank you for reporting an STI. All information will be managed with the strictest confidentiality.														

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile.

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Walla Walla County Health Department may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Walla Walla County Health Department recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Walla Walla County Health Department: (509) 524-2650.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs)^{\dagger}

Alternatives for uncomplicated infections of the cervix, urethra, or rectum:[‡]

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose OR

Cefixime 800 mg orally as a single dose[†]

⁺ If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.

[‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days **OR**

Ofloxacin 300 mg PO BID for 7 days **OR**

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<u>https://www.cdc.gov/std/treatment/default.htm</u>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

DOH 347-102, updated 12/14/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

List of Preferred Languages:

Instructions: Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

(L01) Amharic	(L18) Karen	(L35) Sign languages
(LO2) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(LO4) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown
(L16) Hmong	(L33) Russian	
(L17) Japanese	(L34) Samoan	

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

Extended Race Codes:

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	