


Sexually Transmitted Disease Profile

Clark County 2014



Disease Control and Health Statistics
Infectious Disease Assessment Unit



Sexually Transmitted Disease Profile

Clark County 2014



For persons with disabilities, this document is available in other formats. To submit a request, please call 1-800-525-0127 (TDD/TYY 1-800-833-6388).

Washington State Department of Health
Disease Control and Health Statistics
Infectious Disease Assessment Unit
Olympia, Washington
(360) 236-3455

Contents

Introduction.....1

Data Sources, Definitions, and Limitations1

County STD Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, 2014.4

Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates per 100,000 population, 1995-20144

Figure 2. Age-specific Chlamydia Incidence Rates by Gender, 2014.....5

Figure 3. Chlamydia Incidence Rates by Gender, 1995-20145

Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates per 100,000 population, 1995-20146

Figure 5. Age-specific Gonorrhea Incidence Rates by Gender, 20146

Figure 6. Gonorrhea Incidence Rates by Gender, 1995-20147

Primary & Secondary Syphilis

Figure 7. Primary & Secondary Syphilis Cases and Incidence Rates, 1995-2014.7

Figure 8. Age-specific Primary & Secondary Syphilis Incidence Rates by Gender, 20148

Figure 9. Primary & Secondary Syphilis Incidence Rates by Gender, 1995-2014.....8

Genital Herpes

Figure 10. Genital Herpes Cases and Incidence Rates per 100,000 population 1995-2014.....9

Figure 11. Age-specific Genital Herpes Incidence Rates by Gender, 20149

Figure 12. Genital Herpes Incidence Rates by Gender, 1995-2014.10

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group, 2005-201411

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group, 2005-2014..... 13

Table 4. Primary & Secondary Syphilis Cases and Incidence Rates by Gender and Age
Group, 2005-2014.....15

Table 5. Genital Herpes Cases and Incidence Rates by Gender and Age Group, 2005-2014..17

Introduction

Sexually transmitted infections continue to be the most frequently diagnosed and reported notifiable conditions in Washington State. This report describes the sexually transmitted infection burden in Clark County. Data are presented for the more commonly reported diseases of chlamydial infection, gonorrhea, primary and secondary syphilis, and genital herpes when at least twenty (20) cases of disease were diagnosed in 2014 and the corresponding rates are statistically stable. The incidence rates, including those by age and sex, are presented graphically. The report concludes with tables containing a decade of historical data by age group and gender.

Data Sources, Definitions and Limitations

Cases: Surveillance cases are the number of new episodes of disease (not unique persons) diagnosed in a given year. Cases are identified and submitted by health care providers to local health jurisdictions and entered into the Washington State Department of Health Public Health Information Management System – Sexually Transmitted Diseases (PHIMS-STD) data system. To be included in surveillance reporting, each case must meet disease definitions (see below). Data presented in this report represent new cases of infection diagnosed during a given year and reported as of March 2, 2015.

Confidence Intervals: A confidence interval is a range around a measurement that suggests how precise the measurement is. The confidence interval displays both the possible range around the estimate and also how stable the estimate is. Wider confidence intervals in relation to the estimate itself indicate instability. A confidence interval is always qualified by a particular confidence level, usually expressed as a percentage. In this report, 95% confidence intervals are used.

Disease Definitions:

- | | |
|----------------------|--|
| Chancroid | – A sexually transmitted infection caused by the bacterium <i>Haemophilus ducreyi</i> that may include the symptoms of painful genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of <i>H. ducreyi</i> from a clinical specimen. |
| Chlamydia (CT) | – A sexually transmitted infection caused by the bacterium <i>Chlamydia trachomatis</i> that may include the symptoms of swelling and pain of internal sexual organs, however the infection often has no symptoms in women. Cases are defined by laboratory detection of <i>C. trachomatis</i> from a clinical specimen. |
| Genital Herpes (HSV) | – A sexually transmitted infection caused by the herpes simplex viruses type 1 and type 2 that may include the symptoms of blisters or sores in the genital area. Cases are defined by laboratory detection of herpes simplex virus (HSV1 or HSV2) or positive antibody response from a clinical specimen. Reportable cases include only adult genital initial infection and neonatal infection. |
| Gonorrhea (GC) | – A sexually transmitted infection caused by the bacterium <i>Neisseria gonorrhoeae</i> that may include the symptoms of swelling and pain of internal sexual organs, however the infection sometimes has no symptoms. |

Cases are defined by laboratory detection of the bacterium *N. gonorrhoeae* from a clinical specimen.

Granuloma Inguinale (GI) – A sexually transmitted infection caused by the bacterium *Klebsiella granulomatis* that may include the symptoms of slowly increasing genital sores and swollen pelvic lymph nodes. Cases are defined by microscopic examination of a clinical specimen.

Lymphogranuloma Venereum (LGV) – A sexually transmitted infection caused by three strains of *Chlamydia trachomatis* that may include the symptoms of genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of the L1, L2 and L3 serovars of *C. trachomatis* from a clinical specimen.

Syphilis – A sexually transmitted infection caused by the bacterium *Treponema pallidum* that may include many kinds of symptoms or none at all depending upon the stage of disease. Cases are defined and assigned a stage by a combination of positive blood tests, symptoms, and history of previous treatment. The U.S. Centers for Disease Control and Prevention (CDC) provides guidelines with additional details of surveillance definitions and staging criteria. The stages of primary and secondary (P&S) syphilis are grouped together for analysis in this report.

Primary – identified by the presence of one or many painless sores.

Secondary – identified by the presence of a rash on one or more areas of the body, often with fever, fatigue or other symptoms at the same time.

Other Stages – additional stages of syphilis include, early latent, latent, late latent, and congenital. See CDC guidelines for specific criteria: www.cdc.gov/std/

Incidence Rates: Incidence rates in this report are calculated as the number of new episodes of a disease (not unique persons) diagnosed in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates are not presented when there were fewer than 20 cases of disease reported and the calculated relative standard error was greater than thirty percent (>30%) as this indicates statistical instability.

Limitations: The data presented in this report may be subject to a number of limiting factors. Clinically diagnosed cases (without laboratory confirmation) may be missed through public health surveillance systems. Depending upon diagnosing practices, completeness of reporting may vary by the source of health care. In addition, the diagnosing practitioner is responsible for providing the case information including the patient demographic data items of age and gender upon which many of the analyses in this report depend. Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. Also, small increases or decreases in numbers from year to year can look large if the actual number of cases is small. Care should be taken in interpreting these data in light of known limitations.

Population: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM) April 1, 2014. This data was accessed March 3, 2015 from <http://www.ofm.wa.gov/pop/>

Tabular Data: The data tables are provided in hopes that community and local partners will use these historical data as a resource for future health planning. Users of these data should keep in mind that small numbers often produce rates that can be unstable from year to year and may not be reliable when comparing across strata within a particular year. Data tables for additional years previous are available upon request.

Anyone with specific questions about how these data should be interpreted is encouraged to contact the Infectious Disease Assessment Unit at 360-236-3455.

Clark County STD Disease Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, Clark County, 2014


Disease	County Cases	County Rate§	WA State Rate
Chlamydia	1,534	346.4	376.7
Gonorrhea	208	47.0	88.1
P&S Syphilis	20	4.5	4.8
Genital Herpes	193	43.6	29.9
Chancroid/GI/LGV	0		
Total	1,955		

§ Crude incidence rate per 100,000 population.

* Rates are suppressed for counts under 20 with a corresponding RSE >30% due to statistical instability.


Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates* per 100,000 population, Clark County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 2. Age-specific Chlamydia Incidence Rates* by Gender, Clark County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 3. Chlamydia Incidence Rates by Gender*, Clark County, 1995-2015



* Gender-specific crude incidence with 95% confidence intervals.


Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates* per 100,000 population, Clark County, 1995-2014.




* Crude incidence rate with 95% confidence intervals.

Figure 5. Age-specific Gonorrhea Incidence Rates* by Gender, Clark County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 6. Gonorrhea Incidence Rates by Gender*, Clark County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.


Primary and Secondary Syphilis

Figure 7. Primary and Secondary Syphilis Cases and Incidence Rates* per 100,000 population, Clark County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 8. Age-specific Primary and Secondary Syphilis Incidence Rates* by Gender, Clark County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 9. Primary and Secondary Syphilis Incidence Rates by Gender*, Clark County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.


Genital Herpes

Figure 10. Genital Herpes Cases and Incidence Rates* per 100,000 population, Clark County, 1995-2014




* Crude incidence rate with 95% confidence intervals.

Figure 11. Age-specific Genital Herpes Incidence Rates* by Gender, Clark County, 2014



* Age-specific crude incidence rate with 95% confidence intervals.

Figure 12. Genital Herpes Incidence Rates by Gender*, Clark County, 1995-2014



* Gender-specific crude incidence with 95% confidence intervals.

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2005	0-9	1	1.7	0	0.0	1	3.5
	10-14	5	15.7	0	0.0	5	32.4
	15-19	243	851.1	35	238.5	208	1,499.0
	20-24	351	1,438.2	99	820.7	252	2,041.8
	25-29	156	649.7	59	496.1	97	800.4
	30-34	65	242.4	31	232.5	34	252.2
	35-39	37	126.6	13	88.7	24	164.9
	40-44	18	57.5	11	69.9	7	45.0
	45+	11	7.8	8	11.9	3	4.1
	Missing	14	0.0	3	0.0	11	0.0
	All Ages	901	228.3	259	132.5	642	322.5
2006	0-9	1	1.7	1	3.3	0	0.0
	10-14	3	9.4	0	0.0	3	19.3
	15-19	267	909.1	33	218.6	234	1,639.3
	20-24	303	1,207.4	80	644.4	223	1,758.5
	25-29	155	617.3	46	371.3	109	857.1
	30-34	67	252.2	26	197.1	41	306.6
	35-39	38	125.9	14	92.3	24	160.0
	40-44	12	38.6	8	51.1	4	25.9
	45+	9	6.2	5	7.2	4	5.2
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	855	211.2	213	106.2	642	314.4
2007	0-9	2	3.3	2	6.5	0	0.0
	10-14	4	12.4	0	0.0	4	25.6
	15-19	293	975.8	40	259.1	253	1,734.2
	20-24	334	1,315.7	68	542.1	266	2,071.3
	25-29	142	546.7	37	288.9	105	797.6
	30-34	58	217.9	24	181.8	34	253.3
	35-39	29	94.4	9	58.2	20	131.2
	40-44	15	49.0	9	58.4	6	39.5
	45+	10	6.6	7	9.7	3	3.8
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	889	215.4	196	95.9	691	331.8
2008	0-9	1	1.7	0	0.0	1	3.4
	10-14	15	46.5	1	6.0	14	89.2
	15-19	381	1,255.0	40	256.1	341	2,313.9
	20-24	352	1,385.4	70	557.9	282	2,193.0
	25-29	164	614.8	46	349.4	118	873.3
	30-34	69	255.3	21	156.9	48	351.7
	35-39	30	97.7	9	58.2	21	137.7
	40-44	19	62.9	10	65.6	9	60.0
	45+	25	16.0	15	20.1	10	12.3
	Missing	4	0.0	0	0.0	4	0.0
	All Ages	1061	253.2	212	102.2	848	400.8
2009	0-9	0	0.0	0	0.0	0	0.0
	10-14	13	39.9	1	6.0	12	75.9
	15-19	461	1,521.9	59	378.6	402	2,732.8
	20-24	486	1,921.8	108	864.6	378	2,953.6
	25-29	186	688.0	53	398.0	133	969.6
	30-34	87	315.5	30	219.9	57	409.3
	35-39	40	132.2	23	151.4	17	112.8
	40-44	19	63.2	9	59.3	10	67.3
	45+	22	13.8	14	18.3	8	9.6
	Missing	1	0.0	1	0.0	0	0.0
	All Ages	1315	310.3	298	142.1	1017	475.1

Continued Table 2. Chlamydia

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2010	0-9	0	0	0.0	0	0.0	
	10-14	15	45.7	0	0.0	15	94.2
	15-19	422	1,405.7	69	446.4	353	2,423.8
	20-24	500	2,050.6	103	845.8	397	3,252.8
	25-29	190	719.2	60	462.6	130	966.7
	30-34	115	404.0	38	272.6	77	530.1
	35-39	62	208.8	25	169.7	37	247.3
	40-44	21	70.0	10	66.3	11	73.7
	45+	23	14.1	12	15.5	11	12.9
	Missing	0	0.0	0	0.0	0	0.0
All Ages		1348	316.9	317	151.0	1031	478.6
2011	0-9	0	0.0	0	0.0	0	0.0
	10-14	9	27.4	0	0.0	9	56.5
	15-19	460	1,553.6	68	446.0	392	2,729.3
	20-24	587	2,405.3	134	1,100.0	453	3,706.2
	25-29	230	874.8	71	548.9	159	1,190.2
	30-34	116	399.1	42	295.1	74	498.9
	35-39	43	148.6	22	153.4	21	143.8
	40-44	26	85.2	14	91.2	12	79.1
	45+	19	11.5	9	11.4	10	11.5
	Missing	0	0.0	0	0.0	0	0.0
All Ages		1490	348.1	360	170.4	1130	521.4
2012	0-9	3	5.1	0	0.0	3	10.5
	10-14	5	15.3	0	0.0	5	31.6
	15-19	420	1,396.4	49	317.1	371	2,537.0
	20-24	560	2,133.3	124	941.0	436	3,335.0
	25-29	198	757.2	60	462.5	138	1,047.3
	30-34	101	354.3	38	272.6	63	432.5
	35-39	42	147.8	20	142.6	22	152.8
	40-44	25	80.6	12	76.9	13	84.4
	45+	34	20.1	23	28.4	11	12.4
	Missing	0	0.0	0	0.0	0	0.0
All Ages		1388	321.9	326	153.1	1062	486.4
2013	0-9	0	0.0	0	0.0	0	0.0
	10-14	14	43.3	1	6.0	13	82.7
	15-19	399	1,330.6	63	408.1	336	2,309.3
	20-24	528	1,936.8	107	785.4	421	3,087.0
	25-29	217	830.9	77	593.5	140	1,065.3
	30-34	133	464.9	51	362.2	82	564.5
	35-39	70	246.6	27	193.0	43	298.7
	40-44	27	87.4	14	90.3	13	84.5
	45+	28	16.1	13	15.7	15	16.6
	Missing	5	0.0	0	0.0	5	0.0
All Ages		1421	326.3	353	164.1	1068	484.5
2014	0-9	0	0.0	0	0.0	0	0.0
	10-14	11	34.0	3	18.1	8	50.8
	15-19	375	1,240.4	44	282.2	331	2,261.0
	20-24	563	2,038.2	136	986.1	427	3,087.4
	25-29	273	1,040.6	83	637.6	190	1,437.5
	30-34	138	476.2	57	398.8	81	551.6
	35-39	77	267.0	33	231.9	44	301.0
	40-44	54	176.2	27	176.0	27	176.5
	45+	43	24.0	34	39.6	9	9.6
	Missing	0	0.0	0	0.0	0	0.0
All Ages		1534	346.4	417	190.7	1117	498.3

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2005	0-9	0	0.0	0	0.0	0
	10-14	2	6.3	0	0.0	2
	15-19	30	105.1	3	20.4	27
	20-24	60	245.9	35	290.1	25
	25-29	39	162.4	17	142.9	22
	30-34	21	78.3	10	75.0	11
	35-39	14	47.9	11	75.0	3
	40-44	14	44.7	9	57.2	5
	45+	15	10.7	12	17.9	3
	Missing	1	0.0	1	0.0	0
All Ages		196	49.7	98	50.1	98
2006	0-9	0	0.0	0	0.0	0
	10-14	2	6.2	0	0.0	2
	15-19	25	85.1	9	59.6	16
	20-24	48	191.3	25	201.4	23
	25-29	31	123.5	20	161.4	11
	30-34	14	52.7	7	53.1	7
	35-39	13	43.1	8	52.7	5
	40-44	9	29.0	4	25.6	5
	45+	5	3.4	4	5.7	1
	Missing	1	0.0	0	0.0	1
All Ages		148	36.6	77	38.4	71
2007	0-9	1	1.7	0	0.0	1
	10-14	1	3.1	0	0.0	1
	15-19	37	123.2	10	64.8	27
	20-24	50	197.0	19	151.5	31
	25-29	27	104.0	18	140.5	9
	30-34	23	86.4	12	90.9	11
	35-39	7	22.8	5	32.3	2
	40-44	2	6.5	2	13.0	0
	45+	11	7.3	8	11.1	3
	Missing	0	0.0	0	0.0	0
All Ages		160	38.8	74	36.2	85
2008	0-9	0	0.0	0	0.0	0
	10-14	3	9.3	1	6.0	2
	15-19	30	98.8	7	44.8	23
	20-24	42	165.3	18	143.4	24
	25-29	27	101.2	11	83.6	16
	30-34	27	99.9	15	112.1	12
	35-39	10	32.6	7	45.3	3
	40-44	5	16.5	5	32.8	0
	45+	15	9.6	12	16.1	3
	Missing	1	0.0	1	0.0	0
All Ages		160	38.2	77	37.1	83
2009	0-9	1	1.6	0	0.0	1
	10-14	1	3.1	0	0.0	1
	15-19	26	85.8	5	32.1	21
	20-24	32	126.5	14	112.1	18
	25-29	27	99.9	11	82.6	16
	30-34	15	54.4	10	73.3	5
	35-39	10	33.0	6	39.5	4
	40-44	8	26.6	6	39.5	2
	45+	4	2.5	2	2.6	2
	Missing	0	0.0	0	0.0	0
All Ages		124	29.3	54	25.7	70

Continued Table 3. Gonorrhea

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2010	0-9	0	0.0	0	0.0	0
	10-14	1	3.0	0	0.0	1
	15-19	45	149.9	13	84.1	32
	20-24	47	192.8	17	139.6	30
	25-29	37	140.1	19	146.5	18
	30-34	22	77.3	15	107.6	7
	35-39	7	23.6	4	27.2	3
	40-44	3	10.0	3	19.9	0
	45+	7	4.3	6	7.7	1
	Missing	0	0.0	0	0.0	0
All Ages		169	39.7	77	36.7	92
2011	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	31	104.7	10	65.6	21
	20-24	42	172.1	21	172.4	21
	25-29	35	133.1	16	123.7	19
	30-34	17	58.5	9	63.2	8
	35-39	8	27.6	5	34.9	3
	40-44	7	22.9	5	32.6	2
	45+	20	12.1	17	21.5	3
	Missing	0	0.0	0	0.0	0
All Ages		160	37.4	83	39.3	77
2012	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	17	56.5	10	64.7	7
	20-24	41	156.2	21	159.4	20
	25-29	34	130.0	14	107.9	20
	30-34	21	73.7	8	57.4	13
	35-39	13	45.7	9	64.2	4
	40-44	6	19.3	5	32.0	1
	45+	19	11.2	15	18.5	4
	Missing	0	0.0	0	0.0	0
All Ages		151	35.0	82	38.5	69
2013	0-9	1	1.7	1	3.4	0
	10-14	1	3.1	0	0.0	1
	15-19	6	20.0	1	6.5	5
	20-24	42	154.1	19	139.5	23
	25-29	46	176.1	28	215.8	18
	30-34	20	69.9	16	113.6	4
	35-39	13	45.8	7	50.0	6
	40-44	7	22.7	7	45.1	0
	45+	13	7.5	12	14.5	1
	Missing	0	0.0	0	0.0	0
All Ages		149	34.2	91	42.3	58
2014	0-9	1	1.7	1	3.3	0
	10-14	0	0.0	0	0.0	0
	15-19	20	66.2	6	38.5	14
	20-24	50	181.0	26	188.5	24
	25-29	41	156.3	27	207.4	14
	30-34	39	134.6	31	216.9	8
	35-39	19	65.9	13	91.4	6
	40-44	20	65.3	18	117.3	2
	45+	18	10.0	16	18.6	2
	Missing	0	0.0	0	0.0	0
All Ages		208	47.0	138	63.1	70

Table 4. P&S Syphilis Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2005	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	1	4.1	1	8.3	0
	25-29	1	4.2	1	8.4	0
	30-34	0	0.0	0	0.0	0
	35-39	2	6.8	2	13.6	0
	40-44	0	0.0	0	0.0	0
	45+	1	0.7	1	1.5	0
	Missing	0	0.0	0	0.0	0
All Ages		5	1.3	5	2.6	0
2006	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	0	0.0	0	0.0	0
	30-34	1	3.8	1	7.6	0
	35-39	0	0.0	0	0.0	0
	40-44	0	0.0	0	0.0	0
	45+	1	0.7	1	1.4	0
	Missing	0	0.0	0	0.0	0
All Ages		2	0.5	2	1.0	0
2007	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	0	0.0	0	0.0	0
	45+	1	0.7	1	1.4	0
	Missing	0	0.0	0	0.0	0
All Ages		1	0.2	1	0.5	0
2008	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	1	3.7	1	7.6	0
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	0	0.0	0	0.0	0
	45+	0	0.0	0	0.0	0
	Missing	0	0.0	0	0.0	0
All Ages		1	0.2	1	0.5	0
2009	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	0	0.0	0	0.0	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	1	3.3	1	6.6	0
	40-44	1	3.3	1	6.6	0
	45+	2	1.3	2	2.6	0
	Missing	0	0.0	0	0.0	0
All Ages		4	0.9	4	1.9	0

Continued Table 4. P&S Syphilis

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2010	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	2	8.2	2	16.4	0
	25-29	1	3.8	0	0.0	1
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	1	3.3	1	6.6	0
	45+	2	1.2	2	2.6	0
	Missing	0	0.0	0	0.0	0
All Ages		6	1.4	5	2.4	1
2011	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	1	4.1	1	8.2	0
	25-29	0	0.0	0	0.0	0
	30-34	0	0.0	0	0.0	0
	35-39	0	0.0	0	0.0	0
	40-44	1	3.3	1	6.5	0
	45+	3	1.8	3	3.8	0
	Missing	0	0.0	0	0.0	0
All Ages		5	1.2	5	2.4	0
2012	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	3.3	0	0.0	1
	20-24	3	11.4	2	15.2	1
	25-29	3	11.5	3	23.1	0
	30-34	0	0.0	0	0.0	0
	35-39	4	14.1	4	28.5	0
	40-44	4	12.9	4	25.6	0
	45+	8	4.7	8	9.9	0
	Missing	0	0.0	0	0.0	0
All Ages		24	5.6	21	9.9	2
2013	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	1	3.3	1	6.5	0
	20-24	6	22.0	5	36.7	1
	25-29	3	11.5	3	23.1	0
	30-34	2	7.0	2	14.2	0
	35-39	3	10.6	3	21.4	0
	40-44	1	3.2	1	6.4	0
	45+	7	4.0	7	8.4	0
	Missing	0	0.0	0	0.0	0
All Ages		23	5.3	22	10.2	1
2014	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	0	0.0	0	0.0	0
	20-24	4	14.5	4	29.0	0
	25-29	1	3.8	1	7.7	0
	30-34	1	3.5	1	7.0	0
	35-39	3	10.4	3	21.1	0
	40-44	3	9.8	3	19.6	0
	45+	8	4.5	8	9.3	0
	Missing	0	0.0	0	0.0	0
All Ages		20	4.5	20	9.1	0

Table 5. Genital Herpes Cases and Incidence Rates by Gender and Age Group, 2005-2014

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2005	0-9	0	0	0.0	0	0.0	
	10-14	0	0	0.0	0	0.0	
	15-19	9	31.5	1	6.8	8	57.7
	20-24	25	102.4	9	74.6	16	129.6
	25-29	13	54.1	3	25.2	10	82.5
	30-34	7	26.1	4	30.0	3	22.3
	35-39	6	20.5	1	6.8	5	34.4
	40-44	1	3.2	0	0.0	1	6.4
	45+	9	6.4	2	3.0	7	9.5
	Missing	3	0.0	0	0.0	3	0.0
	All Ages	73	18.5	20	10.2	53	26.6
2006	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	8	27.2	0	0.0	8	56.0
	20-24	11	43.8	3	24.2	8	63.1
	25-29	9	35.8	1	8.1	8	62.9
	30-34	3	11.3	1	7.6	2	15.0
	35-39	3	9.9	1	6.6	2	13.3
	40-44	2	6.4	0	0.0	2	13.0
	45+	2	1.4	1	1.4	1	1.3
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	38	9.4	7	3.5	31	15.2
2007	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	10	33.3	0	0.0	10	68.5
	20-24	10	39.4	5	39.9	5	38.9
	25-29	6	23.1	3	23.4	3	22.8
	30-34	5	18.8	1	7.6	4	29.8
	35-39	3	9.8	1	6.5	2	13.1
	40-44	2	6.5	1	6.5	1	6.6
	45+	7	4.6	2	2.8	5	6.3
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	43	10.4	13	6.4	30	14.4
2008	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	3.1	1	6.0	0	0.0
	15-19	8	26.4	0	0.0	8	54.3
	20-24	28	110.2	6	47.8	22	171.1
	25-29	15	56.2	4	30.4	11	81.4
	30-34	5	18.5	1	7.5	4	29.3
	35-39	9	29.3	3	19.4	6	39.3
	40-44	2	6.6	2	13.1	0	0.0
	45+	12	7.7	5	6.7	7	8.6
	Missing	2	0.0	2	0.0	0	0.0
	All Ages	82	19.6	24	11.6	58	27.4
2009	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	7	23.1	2	12.8	5	34.0
	20-24	19	75.1	7	56.0	12	93.8
	25-29	12	44.4	3	22.5	9	65.6
	30-34	15	54.4	5	36.6	10	71.8
	35-39	7	23.1	5	32.9	2	13.3
	40-44	7	23.3	2	13.2	5	33.6
	45+	20	12.5	9	11.8	11	13.2
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	87	20.5	33	15.7	54	25.2

Continued Table 5. Genital Herpes

Age Group	Total		Males		Females	
	Cases	Rate	Cases	Rate	Cases	Rate
2010	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	12	40.0	4	25.9	8
	20-24	13	53.3	2	16.4	11
	25-29	16	60.6	5	38.6	11
	30-34	14	49.2	7	50.2	7
	35-39	4	13.5	3	20.4	1
	40-44	10	33.3	2	13.3	8
	45+	13	8.0	3	3.9	10
	Missing	0	0.0	0	0.0	0
All Ages		82	19.3	26	12.4	56
2011	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	10	33.8	2	13.1	8
	20-24	12	49.2	5	41.0	7
	25-29	13	49.4	2	15.5	11
	30-34	9	31.0	2	14.1	7
	35-39	5	17.3	2	13.9	3
	40-44	8	26.2	2	13.0	6
	45+	10	6.0	5	6.3	5
	Missing	0	0.0	0	0.0	0
All Ages		67	15.7	20	9.5	47
2012	0-9	1	1.7	0	0.0	1
	10-14	1	3.1	0	0.0	1
	15-19	9	29.9	2	12.9	7
	20-24	22	83.8	9	68.3	13
	25-29	17	65.0	6	46.3	11
	30-34	6	21.0	1	7.2	5
	35-39	5	17.6	1	7.1	4
	40-44	8	25.8	3	19.2	5
	45+	21	12.4	7	8.6	14
	Missing	0	0.0	0	0.0	0
All Ages		90	20.9	29	13.6	61
2013	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	4	13.3	1	6.5	3
	20-24	26	95.4	9	66.1	17
	25-29	25	95.7	7	54.0	18
	30-34	21	73.4	2	14.2	19
	35-39	16	56.4	8	57.2	8
	40-44	16	51.8	5	32.2	11
	45+	45	25.9	14	16.9	31
	Missing	0	0.0	0	0.0	0
All Ages		153	35.1	46	21.4	107
2014	0-9	0	0.0	0	0.0	0
	10-14	0	0.0	0	0.0	0
	15-19	6	19.8	0	0.0	6
	20-24	44	159.3	11	79.8	33
	25-29	30	114.4	9	69.1	21
	30-34	22	75.9	12	83.9	10
	35-39	21	72.8	4	28.1	17
	40-44	11	35.9	6	39.1	5
	45+	59	32.9	26	30.3	33
	Missing	0	0.0	0	0.0	0
All Ages		193	43.6	68	31.1	125
						55.8