
Sexually Transmitted Disease Profile

Grant County 2014



Disease Control and Health Statistics Infectious Disease Assessment Unit



Sexually Transmitted Disease Profile

Grant County 2014



For persons with disabilities, this document is available in other formats. To submit a request, please call 1-800-525-0127 (TDD/TYY 1-800-833-6388).

Washington State Department of Health Disease Control and Health Statistics Infectious Disease Assessment Unit Olympia, Washington (360) 236-3455

DOH Pub# 347-634, May 2015

Contents

Introduction
Data Sources, Definitions, and Limitations
County STD Trends Table 1. Washington State Reportable Sexually Transmitted Diseases, 20144
<u>Chlamydia</u> Figure 1. Chlamydia Cases and Incidence Rates per 100,000 population, 1995-2014
Gonorrhea Figure 4. Gonorrhea Cases and Incidence Rates per 100,000 population, 1995-2014
Genital Herpes Figure 7. Genital Herpes Cases and Incidence Rates per 100,000 population 1995-2014
<u>Data Tables</u> Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group, 2005-20149 Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group, 2005-201411 Table 4. Genital Herpes Cases and Incidence Rates by Gender and Age Group, 2005-2014 13

Introduction

Sexually transmitted infections continue to be the most frequently diagnosed and reported notifiable conditions in Washington State. This report describes the sexually transmitted infection burden in Grant County. Data are presented for the more commonly reported diseases of chlamydial infection, gonorrhea, primary and secondary syphilis, and genital herpes when at least twenty (20) cases of disease were diagnosed in 2014 and the corresponding rates are statistically stable. The incidence rates, including those by age and sex, are presented graphically. The report concludes with tables containing a decade of historical data by age group and gender.

Data Sources, Definitions and Limitations

Cases: Surveillance cases are the number of new episodes of disease (not unique persons) diagnosed in a given year. Cases are identified and submitted by health care providers to local health jurisdictions and entered into the Washington State Department of Health Public Health Information Management System – Sexually Transmitted Diseases (PHIMS-STD) data system. To be included in surveillance reporting, each case must meet disease definitions (see below). Data presented in this report represent new cases of infection diagnosed during a given year and reported as of March 2, 2015.

Confidence Intervals: A confidence interval is a range around a measurement that suggests how precise the measurement is. The confidence interval displays both the possible range around the estimate and also how stable the estimate is. Wider confidence intervals in relation to the estimate itself indicate instability. A confidence interval is always qualified by a particular confidence level, usually expressed as a percentage. In this report, 95% confidence intervals are used.

Disease Definitions:

Chancroid

- A sexually transmitted infection caused by the bacterium *Haemophilus* ducreyi that may include the symptoms of painful genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of H. ducreyi from a clinical specimen.

Chlamydia (CT)

- A sexually transmitted infection caused by the bacterium Chlamydia trachomatis that may include the symptoms of swelling and pain of internal sexual organs, however the infection often has no symptoms in women. Cases are defined by laboratory detection of *C. trachomatis* from a clinical specimen.

Genital Herpes (HSV) – A sexually transmitted infection caused by the herpes simplex viruses type 1 and type 2 that may include the symptoms of blisters or sores in the genital area. Cases are defined by laboratory detection of herpes simplex virus (HSV1 or HSV2) or positive antibody response from a clinical specimen. Reportable cases include only adult genital initial infection and neonatal infection.

Gonorrhea (GC)

- A sexually transmitted infection caused by the bacterium Neisseria gonorrhoeae that may include the symptoms of swelling and pain of internal sexual organs, however the infection sometimes has no symptoms. Cases are defined by laboratory detection of the bacterium N. gonorrhoeae from a clinical specimen.

Granuloma Inguinale (GI) – A sexually transmitted infection caused by the bacterium Klebsiella granulomatis that may include the symptoms of slowly increasing genital sores and swollen pelvic lymph nodes. Cases are defined by microscopic examination of a clinical specimen.

Lymphogranuloma Venereum (LGV) – A sexually transmitted infection caused by three strains of Chlamydia trachomatis that may include the symptoms of genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of the L1, L2 and L3 serovars of C. trachomatis from a clinical specimen.

Syphilis

- A sexually transmitted infection caused by the bacterium *Treponema* pallidum that may include many kinds of symptoms or none at all depending upon the stage of disease. Cases are defined and assigned a stage by a combination of positive blood tests, symptoms, and history of previous treatment. The U.S. Centers for Disease Control and Prevention (CDC) provides guidelines with additional details of surveillance definitions and staging criteria. The stages of primary and secondary (P&S) syphilis are grouped together for analysis in this report.

Primary – identified by the presence of one or many painless sores. Secondary – identified by the presence of a rash on one or more areas of the body, often with fever, fatigue or other symptoms at the same time. Other Stages – additional stages of syphilis include, early latent, latent, late latent, and congenital. See CDC guidelines for specific criteria: www.cdc.gov/std/

Incidence Rates: Incidence rates in this report are calculated as the number of new episodes of a disease (not unique persons) diagnosed in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates are not presented when there were fewer than 20 cases of disease reported and the calculated relative standard error was greater than thirty percent (>30%) as this indicates statistical instability.

Limitations: The data presented in this report may be subject to a number of limiting factors. Clinically diagnosed cases (without laboratory confirmation) may be missed through public health surveillance systems. Depending upon diagnosing practices, completeness of reporting may vary by the source of health care. In addition, the diagnosing practitioner is responsible for providing the case information including the patient demographic data items of age and gender upon which many of the analyses in this report depend. Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. Also, small increases or decreases in numbers from year to year can look large if the actual number of cases is small. Care should be taken in interpreting these data in light of known limitations.

<u>Population</u>: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM) April 1, 2014. This data was accessed March 3, 2015 from http://www.ofm.wa.gov/pop/

<u>Tabular Data</u>: The data tables are provided in hopes that community and local partners will use these historical data as a resource for future health planning. Users of these data should keep in mind that small numbers often produce rates that can be unstable from year to year and may not be reliable when comparing across strata within a particular year. Data tables for additional years previous are available upon request.

Anyone with specific questions about how these data should be interpreted is encouraged to contact the Infectious Disease Assessment Unit at 360-236-3455.

Grant County STD Disease Trends


Table 1. Washington State Reportable Sexually Transmitted Diseases, Grant County, 2014

Disease	County Cases	County Rate§	WA State Rate
Chlamydia	392	422.0	376.7
Gonorrhea	80	86.1	88.1
P&S Syphilis	4	*	4.8
Genital Herpes	14	15.1	29.9
Chancroid/GI/LGV	0		
Total	490		

[§] Crude incidence rate per 100,000 population.

Chlamydia


Figure 1. Chlamydia Cases and Incidence Rates* per 100,000 population, Grant County, 1995-2014



^{*} Crude incidence rate with 95% confidence intervals.


^{*} Rates are suppressed for counts under 20 with a corresponding RSE >30% due to statistical instability.

Figure 2. Age-specific Chlamydia Incidence Rates* by Gender, Grant County, 2014



^{*} Age-specific crude incidence rate with 95% confidence intervals.


Figure 3. Chlamydia Incidence Rates by Gender*, Grant County, 1995-2014



^{*} Gender-specific crude incidence with 95% confidence intervals.


Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates* per 100,000 population, Grant County, 1995-2014



^{*} Crude incidence rate with 95% confidence intervals.

Figure 5. Age-specific Gonorrhea Incidence Rates* by Gender, Grant County, 2014




^{*} Age-specific crude incidence rate with 95% confidence intervals.

Figure 6. Gonorrhea Incidence Rates by Gender*, Grant County, 1995-2014

160


— Male — Female



^{*} Gender-specific crude incidence with 95% confidence intervals.


Genital Herpes

Figure 7. Genital Herpes Cases and Incidence Rates* per 100,000 population, Grant County, 1995-2014




^{*} Crude incidence rate with 95% confidence intervals.

Figure 8. Age-specific Genital Herpes Incidence Rates* by Gender, Grant County, 2014



^{*} Age-specific crude incidence rate with 95% confidence intervals.

Figure 9. Genital Herpes Incidence Rates by Gender*, Grant County, 1995-2014



^{*} Gender-specific crude incidence with 95% confidence intervals.

Data Tables

 Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group, 2005-2014

	Age	ge Total		Males		Females		
	Group	Cases	Rate	Cases	Rate	Cases	Rate	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	5	71.4	0	0.0	5	146.5	
	15-19	74	1,088.7	8	222.6	66	2,060.6	
	20-24	58	1,008.9	17	545.6	41	1,557.2	
2	25-29	29	569.7	8	298.7	21	870.6	
2005	30-34	15	302.3	4	157.3	11	454.7	
2	35-39	7	137.3	2	76.3	5	201.9	
	40-44	0	0.0	0	0.0	0	0.0	
	45+	1	3.8	0	0.0	1	7.5	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	189	235.9	39	95.3	150	382.8	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	2	28.6	0	0.0	2	58.7	
	15-19	72	1,041.2	11	300.8	61	1,872.3	
	20-24	77	1,295.6	18	558.3	59	2,169.9	
9	25-29	36	673.0	5	177.9	31	1,221.4	
2006	30-34	10	201.7	2	78.9	8	330.3	
2	35-39	5	94.8	2	73.7	3	117.0	
	40-44	3	57.5	1	37.2	2	79.2	
	45+	2	7.4	0	0.0	2	14.6	
	Missing	5	0.0	3	0.0	2	0.0	
	All Ages	212	259.2	42	100.5	170	425.1	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	3	42.6	0	0.0	3	87.4	
	15-19	77	1,087.9	11	293.8	66	1,979.6	
	20-24	61	1,006.4	14	425.4	47	1,696.8	
2	25-29	27	482.7	9	305.6	18	679.8	
2007	30-34	15	299.2	4	156.6	11	447.5	
2	35-39	11	202.8	5	179.1	6	228.1	
	40-44	5	97.3	1	37.7	4	161.1	
	45+	0	0.0	0	0.0	0	0.0	
	Missing	12	0.0	5	0.0	7	0.0	
	All Ages	211	252.0	49	114.5	162	395.7	
	0-9	1	6.7	0	0.0	1	13.7	
	10-14	5	70.2	0	0.0	5	143.8	
	15-19	81	1,126.1	15	394.4	66	1,946.9	
	20-24	66	1,072.6	16	480.6	50	1,770.5	
800	25-29	41	700.4	11	355.6	30	1,086.6	
20(30-34	15	288.5	4	151.2	11	430.9	
•	35-39	5 5	90.9	0	0.0	5	187.1	
	40-44		97.6	3	113.0	2	81.1	
	45+	4	13.8	1	7.0	3	20.6	
	Missing	7 233	0.0 271.0	2 52	0.0 118.4	5 178	0.0 423.2	
	All Ages	233				1/0	13.3	
	0-9 10-14	2	6.6 27.7	0	0.0 27.1	1	28.4	
	15-14	83	1,152.6	14	367.9	69	2,031.8	
	20-24	92	1,132.0	16	475.3	76	2,666.7	
	25-29	43	713.5	9	282.0	34	1,199.3	
2009	30-34	17	315.5	4	145.9	13	490.9	
20	35-39	11	200.4	2	71.1	9	336.6	
	40-44	9	175.4	1	37.4	8	325.6	
	40-44 45+	4	173.4	1	6.8	3	20.0	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	262	298.7	48	107.1	214	498.8	
	All Ayes	202	230.7	40	107.1	۷۱۴	+30.0	

Continued Table 2. Chlamydia

	Age	Age Total		М	ales	Fen	nales
	Group	Cases	Rate	Cases	Rate	Cases	Rate
	0-9	0	0.0	0	0.0	0	0.0
0	10-14	1	13.6	0	0.0	1	27.9
	15-19	83	1,153.9	9	236.9	74	2,180.3
	20-24	112	1,835.8	15	449.4	97	3,510.7
	25-29	42	699.1	7	220.3	35	1,236.7
2010	30-34	31	545.7	8	280.2	23	813.9
7	35-39	9	163.9	4	143.8	5	184.4
	40-44	8	154.4	4	148.6	4	160.8
	45+	4	13.1	0	0.0	4	25.9
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	290	325.4	47	103.5	243	556.1
	0-9	0	0.0	0	0.0	0	0.0
	10-14	4	54.1	1	26.4	3	83.1
	15-19 20-24	90 123	1,261.8	13 22	345.1	77 101	2,288.0
	25-29	36	2,003.5 598.7	12	655.3 376.6	24	3,630.5 849.1
2011	30-34	14	240.1	4	136.5	10	344.7
70	35-39	13	240.1	2	73.5	11	413.5
	40-44	5	94.4	1	36.3	4	157.3
	45+	2	6.4	1	6.4	1	6.3
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	287	318.5	56	121.9	231	522.9
	0-9	0	0.0	0	0.0	0	0.0
	10-14	6	80.3	2	52.6	4	109.0
	15-19	112	1,669.0	15	423.3	97	3,062.9
	20-24	102	1,524.3	22	618.4	80	2,552.9
7	25-29	52	843.5	14	426.4	38	1,318.8
2012	30-34	34	566.7	9	291.3	25	858.9
2	35-39	18	329.4	3	108.3	15	556.9
	40-44	4	73.9	4	142.4	0	0.0
	45+	2	6.4	1	6.4	1	6.3
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	330	362.6	70	151.1	260	582.1
	0-9	0	0.0	0	0.0	0	0.0
	10-14 15-19	3	40.0	0	0.0	3	81.1
	20-24	78 154	1,174.7 2,300.7	17 26	490.7 718.9	61 128	1,920.8 4,160.1
	25-29	87	1,406.3	28	849.9	59	2,040.2
က	30-34	36	588.5	9	283.2	27	918.4
2013	35-39	19	343.6	10	355.7	9	331.1
7	40-44	7		2		5	
			127.7		70.6		188.8
	45+	1	3.1	1	6.3	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	385	419.4	93	199.0	292	647.9
	0-9	0	0.0	0	0.0	0	0.0
	10-14	2	26.4	0	0.0	2	53.5
	15-19	83	1,236.7	13	372.6	70	2,172.1
	20-24	144	2,146.9	35	974.1	109	3,500.0
4	25-29	83	1,332.0	26	780.9	57	1,964.3
2014	30-34	40	649.2	13	405.4	27	913.6
.,	35-39	27	480.1	5	174.0	22	799.7
	40-44	6	109.4	3	106.1	3	112.8
	45+	6	18.4	3	18.5	3	18.3
		0					
	Missing		0.0	0	0.0	0	0.0
	All Ages	392	422.0	98	207.4	293	642.0

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group, 2005-2014

	Age	Total		М	Males		Females	
	Group	Cases	Rate	Cases	Rate	Cases	Rate	
	0-9	0	0.0	0	0.0	0	0.0	
2	10-14	0	0.0	0	0.0	0	0.0	
	15-19	3	44.1	0	0.0	3	93.7	
	20-24	6	104.4	4	128.4	2	76.0	
	25-29	4	78.6	2	74.7	2	82.9	
2005	30-34	1	20.2	1	39.3	0	0.0	
N	35-39	1	19.6	1	38.1	0	0.0	
	40-44	0	0.0	0	0.0	0	0.0	
	45+	0	0.0	0	0.0	0	0.0	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	15	18.7	8	19.5	7	17.9	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	4	57.8	0	0.0	4	122.8	
	20-24	3	50.5	2	62.0	1	36.8	
9	25-29	0	0.0	0	0.0	0	0.0	
2006	30-34	1	20.2	1	39.4	0	0.0	
7	35-39	1	19.0	0	0.0	1	39.0	
	40-44	1	19.2	1	37.2	0	0.0	
	45+	0	0.0	0	0.0	0	0.0	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	10	12.2	4	9.6	6	15.0	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	3	42.4	0	0.0	3	90.0	
	20-24	3	49.5	1	30.4	2	72.2	
04	25-29	1	17.9	0	0.0	1	37.8	
2007	30-34	1	19.9	0	0.0	1	40.7	
	35-39 40-44	0	18.4 0.0	0	35.8 0.0	0	0.0 0.0	
	40-44 45+	0	0.0	0	0.0	0	0.0	
	Missing	1	0.0	0	0.0	1	0.0	
	All Ages	10	11.9	2	4.7	8	19.5	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	8	111.2	3	78.9	5	147.5	
	20-24	1	16.3	1	30.0	0	0.0	
ω	25-29	2	34.2	2	64.7	0	0.0	
2008	30-34	0	0.0	0	0.0	0	0.0	
7	35-39	0	0.0	0	0.0	0	0.0	
	40-44	1	19.5	0	0.0	1	40.5	
	45+	1	3.5	0	0.0	1	6.9	
	Missing	1	0.0	0	0.0	1	0.0	
	All Ages	14	16.3	6	13.7	8	19.0	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	2	27.8	0	0.0	2	58.9	
	20-24	2	32.2	1	29.7	1	35.1	
0	25-29	4	66.4	2	62.7	2	70.5	
2009	30-34	1	18.6	1	36.5	0	0.0	
N	35-39	0	0.0	0	0.0	0	0.0	
	40-44	0	0.0	0	0.0	0	0.0	
	45+	0	0.0	0	0.0	0	0.0	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	9	10.3	4	8.9	5	11.7	

Continued Table 3. Gonorrhea

	Age	Total Males		Fe	males		
	Group	Cases	Rate	Cases	Rate	Cases	Rate
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	3	41.7	1	26.3	2	58.9
	20-24	5	82.0	1	30.0	4	144.8
0	25-29	3	49.9	2	62.9	1	35.3
2010	30-34	3	52.8	1	35.0	2	70.8
	35-39	0	0.0	0	0.0	0	0.0
	40-44	2	38.6	2	74.3	0	0.0
	45+	1	3.3	1	6.6	0	0.0
	Missing All Ages	0 17	0.0 19.1	0	0.0 17.6	0	0.0 20.6
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	14.0	0	0.0	1	29.7
	20-24	9	146.6	1	29.8	8	287.6
_	25-29	6	99.8	1	31.4	5	176.9
2011	30-34	2	34.3	1	34.1	1	34.5
2	35-39	1	18.6	0	0.0	1	37.6
	40-44	2	37.7	2	72.6	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	21	23.3	5	10.9	16	36.2
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	7	104.3	1	28.2	6	189.5
	20-24	12	179.3	6	168.6	6	191.5
2012	25-29 30-34	15 11	243.3 183.3	6 7	182.7 226.6	9	312.3 137.4
8	30-3 4 35-39	9	164.7	2	72.2	7	259.9
	40-44	3	55.4	2	71.2	1	38.4
	45+	2	6.4	0	0.0	2	12.6
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	59	64.8	24	51.8	35	78.4
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	0	0.0	0	0.0	0	0.0
	20-24	11	164.3	3	82.9	8	260.0
~	25-29	10	161.6	5	151.8	5	172.9
201;	30-34	6	98.1	5	157.4	1	34.0
7	35-39	3	54.3	2	71.1	1	36.8
	40-44	1	18.2	0	0.0	1	37.8
	45+	3	9.4	3	19.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	34	37.0	18	38.5	16	35.5
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	12	178.8	4	114.7	8	248.2
	20-24	25	372.7	7	194.8	18	578.0
4	25-29	19	304.9	6	180.2	13	448.0
2014	30-34	7	113.6	3	93.6	4	135.4
, ,	35-39	7	124.5	0	0.0	7	254.5
	40-44	4	72.9	2	70.8	2	75.2
	45+	6	18.4	5	30.9	1	6.1
	Missing	0	0.0	0	0.0	0	0.0
	_						
	All Ages	80	86.1	27	57.1	53	116.1

Table 4. Genital Herpes Cases and Incidence Rates by Gender and Age Group, 2005-2014

	Age	Total		М	Males		Females	
	Group	Cases	Rate	Cases	Rate	Cases	Rate	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	5	73.6	1	27.8	4	124.9	
	20-24	8	139.2	1	32.1	7	265.9	
	25-29	2	39.3	1	37.3	1	41.5	
2005	30-34	2	40.3	0	0.0	2	82.7	
7	35-39	1	19.6	0	0.0	1	40.4	
	40-44	2	37.8	0	0.0	2	77.7	
	45+	2	7.6	1	7.6	1	7.5	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	22	27.5	4	9.8	18	45.9	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	2	28.9	0	0.0	2	61.4	
	20-24	1	16.8	1	31.0	0	0.0	
(0	25-29	4	74.8	1	35.6	3	118.2	
2006	30-34	0	0.0	0	0.0	0	0.0	
2(35-39	0	0.0	0	0.0	0	0.0	
	40-44	0	0.0	0	0.0	0	0.0	
	45+	2	7.4	0	0.0	2	14.6	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	9	11.0	2	4.8	7	17.5	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	2	28.3	0	0.0	2	60.0	
	20-24	3	49.5	2	60.8	1	36.1	
2	25-29	3	53.6	1	34.0	2	75.5	
2007	30-34	0	0.0	0	0.0	0	0.0	
7	35-39	0	0.0	0	0.0	0	0.0	
	40-44	2	38.9	1	37.7	1	40.3	
	45+	1	3.6	1	7.2	0	0.0	
	Missing	1	0.0	0	0.0	1	0.0	
	All Ages	12	14.3	5	11.7	7	17.1	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	2	27.8	1	26.3	1	29.5	
	20-24	2	32.5	0	0.0	2	70.8	
œ	25-29	3	51.2	1	32.3	2	72.4	
2008	30-34	1	19.2	0	0.0	1	39.2	
0	35-39	0	0.0	0	0.0	0	0.0	
	40-44	1	19.5	0	0.0	1	40.5	
	45+	1	3.5	1	7.0	0	0.0	
	Missing	0	0.0	0	0.0	0	0.0	
	All Ages	10	11.6	3	6.8	7	16.6	
	0-9	0	0.0	0	0.0	0	0.0	
	10-14	0	0.0	0	0.0	0	0.0	
	15-19	2	27.8	0	0.0	2	58.9	
	20-24	6	96.5	0	0.0	6	210.5	
6	25-29	3	49.8	2	62.7	1	35.3	
2009	30-34	1	18.6	1	36.5	0	0.0	
(A)	35-39	0	0.0	0	0.0	0	0.0	
	40-44	0	0.0	0	0.0	0	0.0	
	45+	1	3.4	0	0.0	1	6.7	
	Missing	1	0.0	0	0.0	1	0.0	
	All Ages	14	16.0	3	6.7	11	25.6	

Continued Table 4. Genital Herpes

	Age	To	otal	Males		Fer	males
	Group	Cases	Rate	Cases	Rate	Cases	Rate
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	13.9	0	0.0	1	29.5
	20-24	5	82.0	1	30.0	4	144.8
0	25-29	2	33.3	0	0.0	2	70.7
2010	30-34	1	17.6	1	35.0	0	0.0
	35-39 40-44	2	36.4 19.3	1 0	36.0	1	36.9 40.2
	40-44 45+	4	13.1	1	0.0 6.6	3	19.4
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	16	18.0	4	8.8	12	27.5
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	14.0	0	0.0	1	29.7
	20-24	4	65.2	0	0.0	4	143.8
7	25-29	0	0.0	0	0.0	0	0.0
201	30-34	4	68.6	1	34.1	3	103.4
,,,	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+ Missing	3	9.6 0.0	0	12.9 0.0	0	6.3 0.0
	All Ages	12	13.3	3	6.5	9	20.4
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	2	29.8	0	0.0	2	63.2
	20-24	6	89.7	0	0.0	6	191.5
2	25-29	1	16.2	0	0.0	1	34.7
2012	30-34	1	16.7	1	32.4	0	0.0
7	35-39	2	36.6	1	36.1	1	37.1
	40-44	0	0.0	0	0.0	0	0.0
	45+	4	12.7	1	6.4	3	18.9
	Missing All Ages	0 16	0.0 17.6	0 3	0.0 6.5	0 13	0.0 29.1
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	15.1	0	0.0	1	31.5
	20-24	4	59.8	1	27.6	3	97.5
	25-29	2	32.3	0	0.0	2	69.2
013	30-34	1	16.3	0	0.0	1	34.0
20	35-39	0	0.0	0	0.0	0	0.0
	40-44	1	18.2	0	0.0	1	37.8
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	9	9.8	1	2.1	8	17.8
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0		0	0.0	0	
			0.0				0.0
	15-19	2	29.8	0	0.0	2	62.1
	20-24	5	74.5	1	27.8	4	128.4
4	25-29	2	32.1	1	30.0	1	34.5
2014	30-34	2	32.5	1	31.2	1	33.8
.,	35-39	1	17.8	0	0.0	1	36.4
	40-44	2	36.5	2	70.8	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	14	15.1	5	10.6	9	19.7
	All Ages	14	13.1	J	10.0	9	19.7