
# Sexually Transmitted Disease Profile

Pend Oreille County 2012



Disease Control and Health Statistics Infectious Disease Assessment Unit



# Sexually Transmitted Disease Profile

Pend Oreille County 2012



For persons with disabilities, this document is available in other formats. To submit a request, please call 1-800-525-0127 (TDD/TYY 1-800-833-6388).

Washington State Department of Health Disease Control and Health Statistics Infectious Disease Assessment Unit Olympia, Washington (360) 236-3455

DOH Pub# 347-634, July 2013

# Contents

Introduction1
Data Sources, Definitions, and Limitations1
<u>County STD Trends</u> Table 1. Washington State Reportable Sexually Transmitted Diseases, 20124
<u>Chlamydia</u> Figure 1. Chlamydia Cases and Incidence Rates per 100,000 population, 1993-20124 Figure 2. Age-specific Chlamydia Incidence Rates by Gender, 20125 Figure 3. Chlamydia Incidence Rates by Gender, 1993-20125
Data Tables Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group6

### Introduction

Sexually transmitted infections continue to be the most frequently diagnosed and reported notifiable conditions in Washington State. This report describes the sexually transmitted disease burden in Pend Oreille County. Data are presented for the more commonly reported diseases of primary and secondary syphilis, chlamydial infection, gonorrhea and genital herpes when at least twenty (20) cases of disease were diagnosed in 2012 and the corresponding rates are statistically stable. The incidence rates, including those by age and sex, are presented graphically. The report concludes with tables containing a decade of historical data by age group and gender.

### **Data Sources, Definitions and Limitations**

<u>Cases</u>: Surveillance cases are the number of new episodes of disease (not unique persons) diagnosed in a given year. Cases are identified and submitted by health care providers to local health jurisdictions and entered into the Washington State Department of Health Public Health Information Management System – Sexually Transmitted Diseases (PHIMS-STD) data system. To be included in surveillance reporting, each case must meet disease definitions (see below). Data presented in this report represent new cases of infection diagnosed during a given year and reported as of March 4, 2013.

<u>Confidence Intervals</u>: A confidence interval is a range around a measurement that suggests how precise the measurement is. The confidence interval displays both the possible range around the estimate and also how stable the estimate is. Wider confidence intervals in relation to the estimate itself indicate instability. A confidence interval is always qualified by a particular confidence level, usually expressed as a percentage. In this report, 95% confidence intervals are used.

**Disease Definitions**:

Chancroid	- A sexually transmitted infection caused by the bacterium <i>Haemophilus ducreyi</i> that may include the symptoms of painful genital sores and
	swollen pelvic lymph nodes. Cases are defined by laboratory detection of
	<i>H. ducreyi</i> from a clinical specimen.
Chlamydia (CT)	- A sexually transmitted infection caused by the bacterium Chlamydia
	trachomatis that may include the symptoms of swelling and pain of
	internal sexual organs, however the infection often has no symptoms in
	women. Cases are defined by laboratory detection of <i>C. trachomatis</i> from
	a clinical specimen.
Genital Herpes (HSV	<i>Y</i> ) – A sexually transmitted infection caused by the herpes simplex viruses
	type 1 and type 2 that may include the symptoms of blisters or sores in the
	genital area. Cases are defined by laboratory detection of herpes simplex
	virus (HSV1 or HSV2) or positive antibody response from a clinical
	specimen. Reportable cases include only adult genital initial infection and
	neonatal infection.
Conomboo (CC)	
Gonorrhea (GC)	- A sexually transmitted infection caused by the bacterium Neisseria
	gonorrhoeae that may include the symptoms of swelling and pain of
	internal sexual organs, however the infection sometimes has no symptoms.
	Cases are defined by laboratory detection of the bacterium <i>N. gonorrhoeae</i>
	from a clinical specimen.

- Granuloma Inguinale (GI) A sexually transmitted infection caused by the bacterium *Klebsiella granulomatis* that may include the symptoms of slowly increasing genital sores and swollen pelvic lymph nodes. Cases are defined by microscopic examination of a clinical specimen.
- Lymphogranuloma Venereum (LGV) A sexually transmitted infection caused by three strains of *Chlamydia trachomatis* that may include the symptoms of genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of the L1, L2 and L3 serovars of *C. trachomatis* from a clinical specimen.
- Syphilis A sexually transmitted infection caused by the bacterium *Treponema pallidum* that may include many kinds of symptoms or none at all depending upon the stage of disease. Cases are defined and assigned a stage by a combination of positive blood tests, symptoms, and history of previous treatment. The U.S. Centers for Disease Control and Prevention (CDC) provides guidelines with additional details of surveillance definitions and staging criteria. The stages of primary and secondary (P&S) syphilis are grouped together for analysis in this report. *Primary* – identified by the presence of one or many painless sores.

*Secondary* – identified by the presence of a rash on one or more areas of the body, often with fever, fatigue or other symptoms at the same time.

*Other Stages* – additional stages of syphilis include, early latent, latent, late latent, and congenital. See CDC guidelines for specific criteria: www.cdc.gov/std/

<u>Incidence Rates</u>: Incidence rates in this report are calculated as the number of new episodes of a disease (not unique persons) diagnosed in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates were not presented for when the calculated relative standard error is greater than thirty percent (>30%) as this indicates statistical instability.

Limitations: The data presented in this report may be subject to a number of limiting factors. Clinically diagnosed cases (without laboratory confirmation) may be missed through public health surveillance systems. Depending upon diagnosing practices, completeness of reporting may vary by the source of health care. In addition, the diagnosing practitioner is responsible for providing the case information including the patient demographic data items of age and gender upon which many of the analyses in this report depend. Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. Also, small increases or decreases in numbers from year to year can look large if the actual number of cases is small. Care should be taken in interpreting these data in light of known limitations.

<u>Population</u>: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM) April 1, 2012.

<u>Tabular Data</u>: The data tables are provided in hopes that community and local partners will use these historical data as a resource for future health planning. Users of these data should keep in mind that small numbers often produce rates that can be unstable from year to year and may not be reliable when comparing across strata within a particular year. Data tables for additional years previous are available upon request.

Anyone with specific questions about how these data should be interpreted is encouraged to contact the Infectious Disease Assessment Unit at 360-236-3455.

## Pend Oreille County STD Disease Trends


Table 1.	Washington State Reportable Sexually Transmitted Diseases,
	Pend Oreille County, 2012

Disease	County Cases	County Rate§	WA State Rate
Chlamydia	22	167.9	360.8
Gonorrhea	4	*	48.1
P&S Syphilis	0	*	4.4
Genital Herpes	1	*	32.2
Chancroid/GI/LGV	0		
Total	27		

§ Crude incidence rate per 100,000 population.
\* Rates are suppressed for counts with RSE >30% due to statistical instability.

#### Chlamydia

### Figure 1. Chlamydia Cases and Incidence Rates\* per 100,000 population, Pend Oreille County, 1993-2012



\* Crude incidence rate with 95% confidence intervals.





Figure 2. Age-specific Chlamydia Incidence Rates\* by Gender, Pend Oreille County, 2012

\* Age-specific crude incidence rate with 95% confidence intervals.





\* Gender-specific crude incidence with 95% confidence intervals.

1	Age	Ace Total		Males		Females	
	Group	Cases	Rate	Cases	Rate	Cases	Rate
	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	99.5	0	0.0	1	204.9
	15-19	12	1,400.2	2	470.6	10	2,314.8
	20-24	3	688.1	0	470.0	3	1,463.4
	25-29	1	239.8	1	507.6	0	0.0
03	30-34	0	0.0	0	0.0	0	0.0
2003	35-39	1	132.6	0	0.0	1	251.9
	40-44	0	0.0	0	0.0	0	0.0
	40-44 45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	18	148.9	3	49.3	15	249.8
	0-9	0	0.0	0	0.0	0	240.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	5	590.3	0	0.0	5	1,176.5
	20-24	6	1,336.3	0	0.0	5	2,803.7
	20-24 25-29	3	702.6	1	492.6	2	2,803.7 892.9
2004	25-29 30-34	0	0.0	0	492.6	0	892.9 0.0
20	30-34 35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	40-44 45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	14	115.0	1	16.3	13	215.3
	0-9	0	0.0	0	0.0	0	213.3
	0-9 10-14	2	207.3	0	0.0	2	432.9
	10-14	3	353.4	0	0.0	2	432.9 710.9
	20-24	4	854.7	0	0.0	4	1,762.1
	20-24	4	0.0	0	0.0	4	0.0
2005	30-34	1	191.6	0	0.0	1	363.6
20	30-34	1	138.5	0	0.0	1	269.5
	40-44	0	0.0	0	0.0	0	209.5
	40-44 45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	11	89.0	0	0.0	11	179.8
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	10-14 15-19	3	352.9	0	0.0	3	0.0 717.7
	20-24	3	623.7	2	813.0	3	425.5
	20-24 25-29	2	434.8	∠ 1	458.7	1	425.5 413.2
006	25-29 30-34	0	434.0	0	456.7	0	413.2
20	35-39	1	138.9	0	0.0	1	274.0
	40-44	0	0.0	0	0.0	0	0.0
	40-44 45+	0	0.0	0	0.0	0	0.0
	45+ Missing	1	0.0	0	0.0	1	0.0
	All Ages	10	80.0	3	47.4	7	113.3
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	5	591.0	1	232.0	4	963.9
	20-24	9	1,297.2	2	813.0	7	2,904.6
	20-24	3	626.3	2 1	436.7	2	2,904.0
2007	25-29 30-34	3	597.6	2	826.4	1	384.6
20	30-34 35-39	0	0.0	2	0.0	0	304.0 0.0
	40-44	0	0.0	0	0.0	0	0.0
	40-44 45+	0	0.0	0	0.0	0	0.0
	45+ Missing	0		0		0	
			0.0		0.0		0.0
	All Ages	20	158.0	6	93.6	14	224.0

## Data Tables Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group

# Continued Table 2. Chlamydia

1	Age Total		M	ales	Females		
	Group	Cases	Rate	Cases	Rate	Cases	Rate
	0-9	0	0.0	0	0.0	0	0.0
8	10-14	0	0.0	0	0.0	0	0.0
	15-19	5	598.1	1	233.6	4	980.4
	20-24	5	1,010.1	2	806.5	3	1,214.6
	25-29	4	804.8	1	414.9	3	1,171.9
2008	30-34	0	0.0	0	0.0	0	0.0
5	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	2	0.0 124.5	0	0.0 61.3	2	0.0
	All Ages	16	-	4		12	189.5
	0-9 10-14	0	0.0 0.0	0	0.0 0.0	0	0.0 0.0
	10-14	7	0.0 867.4	3	0.0 724.6	4	0.0 1,017.8
	20-24	6	1,209.7	2	816.3	4	1,593.6
	25-29	4	781.3	1	400.0	4	1,145.0
2009	30-34	0	0.0	0	0.0	0	0.0
20	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	1	14.7	0	0.0	1	29.6
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	18	140.2	6	92.1	12	189.8
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	5	632.1	0	0.0	5	1,308.9
	20-24	15	3,073.8	5	2,057.6	10	4,081.6
0	25-29	0	0.0	0	0.0	0	0.0
2010	30-34	1	187.6	0	0.0	1	370.4
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	21	161.5	5	75.8	16	249.7
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	3	360.4	0	0.0	3	738.6
	20-24 25-29	5	994.0 1,212.8	2	776.8 822.6	3 4	1,221.9 1,589.9
-	30-34	0	0.0	0	0.0	0	0.0
01	35-39	0	0.0	0	0.0	0	0.0
2		2		2		0	
	40-44		278.3	1	539.3		0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	0	123.1	6	91.1	10	156.0
	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	8	969.4	2	478.7	6	1,472.3
	20-24	7	1,305.6	1	357.8	6	2,337.8
2	25-29	4	817.6	1	415.0	3	1,208.2
2012	30-34	3	531.3	1	358.7	2	699.6
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	-						
	All Ages	22	167.9	5	75.4	17	263.0