


Sexually Transmitted Disease Profile

Whitman County 2012



Disease Control and Health Statistics
Infectious Disease Assessment Unit



Sexually Transmitted Disease Profile

Whitman County 2012



For persons with disabilities, this document is available in other formats. To submit a request, please call 1-800-525-0127 (TDD/TYY 1-800-833-6388).

Washington State Department of Health
Disease Control and Health Statistics
Infectious Disease Assessment Unit
Olympia, Washington
(360) 236-3455

DOH Pub# 347-634, July 2013

Contents

Introduction.....	1
Data Sources, Definitions, and Limitations	1
<u>County STD Trends</u>	
Table 1. Washington State Reportable Sexually Transmitted Diseases, 2012.	4
<u>Chlamydia</u>	
Figure 1. Chlamydia Cases and Incidence Rates per 100,000 population, 1993-2012	4
Figure 2. Age-specific Chlamydia Incidence Rates by Gender, 2012	5
Figure 3. Chlamydia Incidence Rates by Gender, 1993-2012	5
<u>Gonorrhea</u>	
Figure 4. Gonorrhea Cases and Incidence Rates per 100,000 population, 1993-2012	6
Figure 5. Age-specific Gonorrhea Incidence Rates by Gender, 2012	6
Figure 6. Gonorrhea Incidence Rates by Gender, 1993-2012	7
<u>Data Tables</u>	
Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group	8
Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group	10

Introduction

Sexually transmitted infections continue to be the most frequently diagnosed and reported notifiable conditions in Washington State. This report describes the sexually transmitted disease burden in Whitman County. Data are presented for the more commonly reported diseases of primary and secondary syphilis, chlamydial infection, gonorrhea and genital herpes when at least twenty (20) cases of disease were diagnosed in 2012 and the corresponding rates are statistically stable. The incidence rates, including those by age and sex, are presented graphically. The report concludes with tables containing a decade of historical data by age group and gender.

Data Sources, Definitions and Limitations

Cases: Surveillance cases are the number of new episodes of disease (not unique persons) diagnosed in a given year. Cases are identified and submitted by health care providers to local health jurisdictions and entered into the Washington State Department of Health Public Health Information Management System – Sexually Transmitted Diseases (PHIMS-STD) data system. To be included in surveillance reporting, each case must meet disease definitions (see below). Data presented in this report represent new cases of infection diagnosed during a given year and reported as of March 4, 2013.

Confidence Intervals: A confidence interval is a range around a measurement that suggests how precise the measurement is. The confidence interval displays both the possible range around the estimate and also how stable the estimate is. Wider confidence intervals in relation to the estimate itself indicate instability. A confidence interval is always qualified by a particular confidence level, usually expressed as a percentage. In this report, 95% confidence intervals are used.

Disease Definitions:

- Chancroid – A sexually transmitted infection caused by the bacterium *Haemophilus ducreyi* that may include the symptoms of painful genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of *H. ducreyi* from a clinical specimen.
- Chlamydia (CT) – A sexually transmitted infection caused by the bacterium *Chlamydia trachomatis* that may include the symptoms of swelling and pain of internal sexual organs, however the infection often has no symptoms in women. Cases are defined by laboratory detection of *C. trachomatis* from a clinical specimen.
- Genital Herpes (HSV) – A sexually transmitted infection caused by the herpes simplex viruses type 1 and type 2 that may include the symptoms of blisters or sores in the genital area. Cases are defined by laboratory detection of herpes simplex virus (HSV1 or HSV2) or positive antibody response from a clinical specimen. Reportable cases include only adult genital initial infection and neonatal infection.
- Gonorrhea (GC) – A sexually transmitted infection caused by the bacterium *Neisseria gonorrhoeae* that may include the symptoms of swelling and pain of internal sexual organs, however the infection sometimes has no symptoms. Cases are defined by laboratory detection of the bacterium *N. gonorrhoeae* from a clinical specimen.

Granuloma Inguinale (GI) – A sexually transmitted infection caused by the bacterium *Klebsiella granulomatis* that may include the symptoms of slowly increasing genital sores and swollen pelvic lymph nodes. Cases are defined by microscopic examination of a clinical specimen.

Lymphogranuloma Venereum (LGV) – A sexually transmitted infection caused by three strains of *Chlamydia trachomatis* that may include the symptoms of genital sores and swollen pelvic lymph nodes. Cases are defined by laboratory detection of the L1, L2 and L3 serovars of *C. trachomatis* from a clinical specimen.

Syphilis – A sexually transmitted infection caused by the bacterium *Treponema pallidum* that may include many kinds of symptoms or none at all depending upon the stage of disease. Cases are defined and assigned a stage by a combination of positive blood tests, symptoms, and history of previous treatment. The U.S. Centers for Disease Control and Prevention (CDC) provides guidelines with additional details of surveillance definitions and staging criteria. The stages of primary and secondary (P&S) syphilis are grouped together for analysis in this report.

Primary – identified by the presence of one or many painless sores.

Secondary – identified by the presence of a rash on one or more areas of the body, often with fever, fatigue or other symptoms at the same time.

Other Stages – additional stages of syphilis include, early latent, latent, late latent, and congenital. See CDC guidelines for specific criteria: www.cdc.gov/std/

Incidence Rates: Incidence rates in this report are calculated as the number of new episodes of a disease (not unique persons) diagnosed in a given year divided by the total population (age and sex appropriate) for that year, expressed as a rate per 100,000. Incidence rates allow comparisons between two or more populations by standardizing the denominator and are the most appropriate statistic to use when investigating differences between groups. Rates were not presented for when the calculated relative standard error is greater than thirty percent (>30%) as this indicates statistical instability.

Limitations: The data presented in this report may be subject to a number of limiting factors. Clinically diagnosed cases (without laboratory confirmation) may be missed through public health surveillance systems. Depending upon diagnosing practices, completeness of reporting may vary by the source of health care. In addition, the diagnosing practitioner is responsible for providing the case information including the patient demographic data items of age and gender upon which many of the analyses in this report depend. Biases could exist in the data due to under-reporting, inability of certain populations to access medical services, error in laboratory reporting, or differential reporting or screening by disease and source of care. Also, small increases or decreases in numbers from year to year can look large if the actual number of cases is small. Care should be taken in interpreting these data in light of known limitations.

Population: Denominator population estimates for incidence rates are from Washington State Adjusted Population Estimates, Office of Financial Management (OFM) April 1, 2012.

Tabular Data: The data tables are provided in hopes that community and local partners will use these historical data as a resource for future health planning. Users of these data should keep in mind that small numbers often produce rates that can be unstable from year to year and may not be reliable when comparing across strata within a particular year. Data tables for additional years previous are available upon request.

Anyone with specific questions about how these data should be interpreted is encouraged to contact the Infectious Disease Assessment Unit at 360-236-3455.

Whitman County STD Disease Trends

Table 1. Washington State Reportable Sexually Transmitted Diseases, Whitman County, 2012


Disease	County Cases	County Rate§	WA State Rate
Chlamydia	173	376.5	360.8
Gonorrhea	26	56.6	48.1
P&S Syphilis	2	*	4.4
Genital Herpes	10	*	32.2
Chancroid/GI/LGV	0		
Total	211		

§ Crude incidence rate per 100,000 population.

* Rates are suppressed for counts with RSE >30% due to statistical instability.


Chlamydia

Figure 1. Chlamydia Cases and Incidence Rates* per 100,000 population, Whitman County, 1993-2012




* Crude incidence rate with 95% confidence intervals.

Figure 2. Age-specific Chlamydia Incidence Rates* by Gender, Whitman County, 2012



* Age-specific crude incidence rate with 95% confidence intervals.


Figure 3. Chlamydia Incidence Rates by Gender*, Whitman County, 1993-2012



* Gender-specific crude incidence with 95% confidence intervals.


Gonorrhea

Figure 4. Gonorrhea Cases and Incidence Rates* per 100,000 population, Whitman County, 1993-2012




* Crude incidence rate with 95% confidence intervals.

Figure 5. Age-specific Gonorrhea Incidence Rates* by Gender, Whitman County, 2012



* Age-specific crude incidence rate with 95% confidence intervals.

Figure 6. Gonorrhea Incidence Rates by Gender*, Whitman County, 1993-2012



* Gender-specific crude incidence with 95% confidence intervals.

Data Tables

Table 2. Chlamydia Cases and Incidence Rates by Gender and Age Group

Age Group	Total		Males		Females		
	Cases	Rate	Cases	Rate	Cases	Rate	
2003	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	41	785.3	6	243.7	35	1,268.6
	20-24	71	687.1	23	414.6	48	1,003.1
	25-29	13	443.1	5	312.5	8	599.7
	30-34	2	89.6	2	176.5	0	0.0
	35-39	2	104.0	1	104.4	1	103.6
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	2	0.0	0	0.0	2	0.0
	All Ages	131	319.9	37	178.0	94	466.0
2004	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	51.2	0	0.0	1	106.2
	15-19	40	744.3	8	315.7	32	1,126.8
	20-24	97	909.6	36	627.0	61	1,239.3
	25-29	11	363.4	5	304.0	6	434.2
	30-34	3	136.2	1	89.2	2	184.8
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	3	28.0	1	19.3	2	36.2
	Missing	1	0.0	0	0.0	1	0.0
	All Ages	156	374.2	51	240.8	105	512.0
2005	0-9	0	0.0	0	0.0	0	0.0
	10-14	2	103.6	0	0.0	2	215.1
	15-19	41	743.7	10	384.3	31	1,064.9
	20-24	82	751.2	26	442.4	56	1,111.3
	25-29	11	349.0	8	470.9	3	206.5
	30-34	3	137.9	1	89.4	2	189.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	1	47.5	0	0.0	1	94.2
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	140	329.4	45	208.3	95	454.6
2006	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	30	537.7	8	302.8	22	749.1
	20-24	80	715.9	21	349.6	59	1,141.9
	25-29	10	305.5	3	171.1	7	460.5
	30-34	2	94.2	1	91.1	1	97.5
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	2	17.9	2	37.1	0	0.0
	Missing	1	0.0	0	0.0	1	0.0
	All Ages	125	291.7	35	160.5	90	427.6
2007	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	20	354.7	5	186.2	15	508.1
	20-24	87	779.6	29	483.8	58	1,122.9
	25-29	8	238.0	3	167.1	5	319.1
	30-34	3	143.2	1	91.7	2	199.2
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	1	8.9	1	18.4	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	122	285.1	39	178.9	80	381.1

Continued Table 2. Chlamydia

	Age Group	Total		Males		Females	
		Cases	Rate	Cases	Rate	Cases	Rate
2008	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	36	611.6	7	248.4	29	945.2
	20-24	96	836.5	29	470.7	67	1,260.3
	25-29	9	251.8	4	210.2	5	299.2
	30-34	2	92.4	1	88.3	1	96.9
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	1	0.0	0	0.0	1	0.0
	All Ages	144	326.8	41	182.5	103	476.9
2009	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	39	653.3	7	244.4	32	1,030.3
	20-24	73	643.2	26	427.1	47	893.4
	25-29	15	419.2	7	370.2	8	474.2
	30-34	2	91.3	1	86.6	1	96.6
	35-39	2	109.3	0	0.0	2	223.2
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	131	297.7	41	182.7	90	417.4
2010	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	34	559.9	6	206.1	28	885.8
	20-24	83	728.5	25	405.7	58	1,108.6
	25-29	11	303.8	6	316.0	5	290.4
	30-34	2	86.1	1	82.0	1	90.6
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	1	8.3	1	17.1	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	131	292.6	39	171.0	92	418.8
2011	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	49	566.1	10	241.3	39	864.6
	20-24	93	829.9	38	635.3	55	1,052.9
	25-29	8	231.1	4	214.0	4	251.1
	30-34	4	192.7	1	94.5	3	294.6
	35-39	2	125.1	1	122.6	1	127.6
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	156	348.2	54	238.3	102	460.8
2012	0-9	0	0.0	0	0.0	0	0.0
	10-14	1	62.7	0	0.0	1	131.1
	15-19	49	535.5	14	318.7	35	735.7
	20-24	102	866.6	32	507.4	70	1,281.2
	25-29	15	405.7	8	400.4	7	412.0
	30-34	2	98.4	1	96.0	1	100.9
	35-39	1	64.4	0	0.0	1	130.1
	40-44	1	61.7	0	0.0	1	121.8
	45+	2	18.1	1	18.8	1	17.5
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	173	376.5	56	240.6	117	516.1

Table 3. Gonorrhea Cases and Incidence Rates by Gender and Age Group

	Age Group	Total		Males		Females	
		Cases	Rate	Cases	Rate	Cases	Rate
2003	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	19.2	0	0.0	1	36.2
	20-24	5	48.4	1	18.0	4	83.6
	25-29	1	34.1	1	62.5	0	0.0
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	7	17.1	2	9.6	5	24.8
2004	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	0	0.0	0	0.0	0	0.0
	20-24	3	28.1	1	17.4	2	40.6
	25-29	1	33.0	0	0.0	1	72.4
	30-34	0	0.0	0	0.0	0	0.0
	35-39	1	52.5	0	0.0	1	105.2
	40-44	1	46.9	0	0.0	1	92.9
	45+	1	9.3	1	19.3	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	7	16.8	2	9.4	5	24.4
2005	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	0	0.0	0	0.0	0	0.0
	20-24	1	9.2	0	0.0	1	19.8
	25-29	0	0.0	0	0.0	0	0.0
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	1	47.5	1	95.7	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	2	4.7	1	4.6	1	4.8
2006	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	17.9	0	0.0	1	34.0
	20-24	4	35.8	3	49.9	1	19.4
	25-29	1	30.6	1	57.0	0	0.0
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	6	14.0	4	18.3	2	9.5
2007	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	3	53.2	2	74.5	1	33.9
	20-24	4	35.8	1	16.7	3	58.1
	25-29	2	59.5	2	111.4	0	0.0
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	9	21.0	5	22.9	4	19.1

Continued Table 3. Gonorrhea

	Age Group	Total		Males		Females	
		Cases	Rate	Cases	Rate	Cases	Rate
2008	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	17.0	1	35.5	0	0.0
	20-24	13	113.3	5	81.2	8	150.5
	25-29	1	28.0	0	0.0	1	59.8
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	1	8.6	1	17.8	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	16	36.3	7	31.2	9	41.7
2009	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	2	33.5	1	34.9	1	32.2
	20-24	4	35.2	0	0.0	4	76.0
	25-29	4	111.8	3	158.6	1	59.3
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	10	22.7	4	17.8	6	27.8
2010	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	1	16.5	0	0.0	1	31.6
	20-24	7	61.4	0	0.0	7	133.8
	25-29	3	82.9	2	105.3	1	58.1
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	11	24.6	2	8.8	9	41.0
2011	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	2	23.1	2	48.3	0	0.0
	20-24	4	35.7	2	33.4	2	38.3
	25-29	5	144.4	3	160.5	2	125.5
	30-34	0	0.0	0	0.0	0	0.0
	35-39	0	0.0	0	0.0	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	11	24.6	7	30.9	4	18.1
2012	0-9	0	0.0	0	0.0	0	0.0
	10-14	0	0.0	0	0.0	0	0.0
	15-19	3	32.8	1	22.8	2	42.0
	20-24	12	102.0	7	111.0	5	91.5
	25-29	6	162.3	4	200.2	2	117.7
	30-34	4	196.7	2	192.0	2	201.8
	35-39	1	64.4	1	127.4	0	0.0
	40-44	0	0.0	0	0.0	0	0.0
	45+	0	0.0	0	0.0	0	0.0
	Missing	0	0.0	0	0.0	0	0.0
	All Ages	26	56.6	15	64.4	11	48.5